Production Note

Cornell University Library produced this volume to replace the irreparably deteriorated original. It was scanned using Xerox software and equipment at 600 dots per inch resolution and compressed prior to storage using CCITT Group 4 compression. The digital data were used to create Cornell's replacement volume on paper that meets the ANSI Standard Z39.48-1984. The production of this volume was supported in part by the Commission on Preservation and Access and the Xerox Corporation. 1990.

The Kelly Library.

PRESENTED TO

THE CORNELL UNIVERSITY, 1870,

BY

The Hon. William Kelly

OF RHINEBECK.

MATHEMATICS

Digitized by the Internet Archive
in 2026

https://archive.org/details/elementsofdiffer0000jryo

M. M. Newth
August 1844.

ELEMENTS

OF THE

DIFFERENTIAL CALCULUS;

COMPREHENDING THE

GENERAL THEORY OF CURVE SURFACES,

AND OF

CURVES OF DOUBLE CURVATURE.

SECOND EDITION, CORRECTED AND ENLARGED.

BY J. R. YOUNG,

PROFESSOR OF MATHEMATICS IN BELFAST COLLEGE.

LONDON:

PRINTED FOR JOHN SOUTER, 131, FLEET STREET;

By J. and C. Adlard, Bartholomew Close.

1836.

Kelly

An Edition of this Work is published also in 8vo., *for the use of the Universities, price* 12*s.*

PREFACE.

THE object of the present volume is to teach the principles of the *Differential Calculus*, and to shew the application of those principles to several interesting and important enquiries, more particularly to the general theory of Curve Surfaces, and of Curves of Double Curvature. Throughout these applications I have endeavoured to preserve the strictest rigour in the various processes employed, so that the student who may have hitherto been accustomed only to the pure reasoning of the ancient geometry, will not, I think, find in these higher order of researches any principle adopted, or any assumption made, inconsistent with his previous notions of mathematical accuracy. Logical strictness of deduction is an ingredient which every one has a right to expect who opens a work professedly mathematical; and to admit that either in the exposition of the theory of the Calculus, or in its practical applications to physical enquiries, there is necessarily involved any principle that will not bear the most scrupulous examination, is tantamount to a declaration that the subject has no claim to a place among the mathematical sciences.

Yet I suspect that it is the adoption of exceptionable principles, and even, in some cases, of contradictory theories, into the elements of this science, that has been the chief cause why it has hitherto been so little studied in a country

where the ancient geometry has been so extensively and so successfully cultivated. The student who proceeds from the works of *Euclid* or of *Apollonius* to study those of our modern analysts, will be naturally enough startled to find that in the *theory* of the differential calculus he is to consider that as absolutely *nothing* which, in the *Geometrical applications* of that theory, is to be considered as a quantity *infinitely small;* and he will very properly hesitate to receive a conclusion as general, when he is at the same time told that the process which led to that conclusion has failing cases; and yet one or both of these inconsistencies pervade more or less every book on the calculus which I have had an opportunity of examining.

The whole theory of what the French mathematicians vaguely call *consecutive points* and *consecutive elements* involves the first of these objectionable principles; for, if the abscissa of any point be represented by x, then the abscissa of the consecutive point, or that separated from the former by an infinitely small interval, is represented by $x + \mathrm{d}x$, although $\mathrm{d}x$, at the outset of the subject, is said to be 0. Again, the theory of tangents, the radius of curvature, principles of osculation, &c. are all made to depend upon Taylor's theorem, and therefore can strictly apply only at those points of the curve where this theorem does not fail: the conclusions, however, are to be received in all their generality.*

If the foregoing statement be true, it is not to be wondered

* I am anxious not to be misunderstood here, and shall therefore state specifically the nature of my objection. In establishing the theory of contact, &c. by aid of Taylor's theorem, it is assumed that a value may be given to the increment h so small as to render the term into which it enters greater than all the following terms of the series taken together. Now how can a function of absolutely indeterminate quantities be shewn to

at that students so often abandon the study of this science, less discouraged with its difficulties than disgusted with its apparent inconsistencies. To remove some of these harassing impediments to the student's progress, has been my object in the present volume; and, although my endeavours may not have entirely succeeded, I have still reason to hope that they have not entirely failed. The following brief outline will convey a notion of the extent and pretensions of the book; a more detailed enumeration of the various topics treated of will be found in the table of contents.

I have taken for the basis of the theory the method of *limits* first employed by *Newton*, although designated by foreign writers as the *method of D'Alembert;* and I have

be greater or less than a series of other functions of the same indeterminate quantities, without, at least, assuming some determinate relation among them? If we say that the assertion applies, whatever particular value we substitute for the indeterminate in the proposed functions or differential coefficients, we merely shift the dilemma, for an indefinite number of these particular values may render the functions all infinite; and we shall be equally at a loss to conceive how one of these infinite quantities can be greater or less than the others. It appears, therefore, that the usual process by which the theory of contact is established, applies rigorously only to those points of curves for which Taylor's development does not fail, and I cannot help thinking that on these grounds the *Analytical Theory of Functions, by Lagrange*, in its application to Geometry is defective, although I feel anxious to express my opinion of that celebrated performance with all becoming caution and humility. Indeed Lagrange himself has adverted to this defect and observes, (*Théorie des Fonctions, p.* 181,) "Quoique ces exceptions ne portent aucune atteinte à la théorie générale, il est nécessaire, pour ne rien laisser à desirer, de voir comment elle doit être modifier dans les cas particuliers dont il s'agit." (*See note* C *at the end.*) But he has not modified the expression deduced from this exceptionable theory for the radius of curvature, which indeed is always applicable, whether the differential coefficients become infinite or not, although, for reasons already assigned, the process which led to it restricts its application to particular points.

done so because I consider this method to be as unexceptionable as that of *Lagrange*, and, on account of its greater simplicity, better adapted to elementary instruction.

In the First chapter, which is devoted to an exposition of the fundamental principles of the Calculus, I have endeavoured to convey a clear and distinct notion of the true meaning of the expression $\frac{dy}{dx}$, the symbol employed by modern analysts to represent the limiting or ultimate ratio of the increment of any function y to that of the variable x, on which it depends.

The *general* expression for this ratio is $\frac{\textit{increment of } y}{\textit{increment of } x}$ and the ratio or quotient itself is the result of the division here implied, a result which will of course vary with the *increment of* x. In the Calculus, the *increment of* x is supposed to vary continually towards, and at length to merge into, zero; and the corresponding quotient, during these changes, is found continually to approach towards, and at length to merge into, a certain fixed and determinate value; which value is therefore the limiting, or ultimate, or extreme value of the entire series of values through which the varying quotient passes. It is this extreme value which the Calculus selects for its operations, and which is represented by $\frac{dy}{dx}$.

It is obvious that the expression $\frac{\textit{increment of } x}{\textit{increment of } y}$, which all along represents the varying quotient of which I have been speaking, takes the form $\frac{0}{0}$ when that quotient arrives at its extreme value. But it is known that in other enquiries this is often a symbol of dubious import: it is sometimes the indication of incompatible conditions in the analytical statements which have led to it; sometimes the symbol of multiple values; and sometimes wholly unintelligible, or admitting

of altogether an arbitrary interpretation. Although, therefore, in the Calculus, this symbol is limited to a distinct and definite interpretation, yet it would be most inconvenient to use a form that would be perpetually reminding us of vagueness and ambiguity; and thus the symbol $\frac{dy}{dx}$ is employed in its stead, the zeros being replaced by dy, dx. Some writers, however, are very averse to considering dy, dx, as in themselves of absolutely no value, and prefer to call them "infinitely small quantities," meaning by "an infinitely small quantity" that which continually oscillates between something and nothing without ever attaining to either. But as I can form no conception of such things, all mention of them is necessarily excluded from the present work.

The Second chapter is a practical illustration of the first. It commences by shewing that if $f(x)$ represent any function of x, and x be changed into $x+h$, the new state $f(x+h)$ of the function may always be developed according to the ascending integral powers of the increment h; and this leads to the important conclusion that the coefficient of the second term in the development of the function $f(x+h)$ is the differential coefficient derived from the function $f(x)$; a fact which *Lagrange* has made the foundation of his theory of analytical functions. The chapter then goes on to treat of the differentiation of the various kinds of functions, algebraic and transcendental, direct and inverse, and concludes with an article on successive differentiation.

The Third chapter is devoted to *Maclaurin's theorem*, and its application is shewn in the development of a great variety of functions. Occasion is taken, in the course of this chapter, to introduce to the student's attention some valuable analytical formulas and expressions from *Euler*, *Demoivre*, *Cotes*, and other celebrated analysts, together

with those curious properties of the circle discovered by *Cotes* and *Demoivre*.

The Fourth chapter is on *Taylor's theorem*, which makes known the actual development of the function $f(x + h)$ according to the form established in the second chapter. From this theorem are derived commodious expressions for the total differential coefficient when the function is complicated, and whether its form be explicit or implicit; the whole being illustrated by a variety of examples.

The Fifth chapter contains the complete theory of vanishing fractions, and points out the principal sources of error which have led some mathematicians to dispute the legitimacy of the conclusions arrived at in the ordinary expositions of this theory.

The charge of bad logic in the processes in question appears to me to arise from a complete misconception, on the part of the objectors, as to what those processes really are.

The Sixth chapter is on the maxima and minima values of functions of a single variable, and will, I think, be found to contain several original remarks and improved modes of solution.

Chapter the Seventh is on the differentiation and development of functions of two independent variables. The usual method of obtaining the development of a function of two variables according to the powers of the increments is to develop first on the supposition that x only varies, and that y is constant; and afterwards to consider y, which is *assumed* to enter into the coefficients, to be changed into $y + h$. But y may be so combined with x in the function $F(x, y)$ that it shall, when considered as a constant, disappear from all the differential coefficients, which circumstance should be pointed out, and be shewn not to affect the truth of the result: I have, however, avoided the necessity of shewing this, by

proceeding rather differently. The chapter concludes with *Lagrange's Theorem*, concisely demonstrated, and applied to several examples.

The Eighth chapter completes the theory of maxima and minima, by applying the principles delivered in Chap. VI. to functions of two independent variables, and it also contains an important article on changing the independent variable.

The Ninth chapter is devoted to a matter of considerable importance, viz. to the examination of the cases in which Taylor's theorem fails; and I have, I think, satisfactorily shewn, that these failing cases are always indicated by the differential coefficients becoming infinite, and that the theorem does not fail when these coefficients become imaginary, as *Lacroix*, and others after him, have asserted.

These nine chapters constitute the First Section of the work, and comprise the pure theory of the subject; the remaining part is devoted to the application of this to geometry, and is divided into two parts, the first containing the theory of plane curves, and the second the theory of curve surfaces, and of curves of double curvature.

The First chapter in the Second Section explains the method of tangents, and the general differential equation of the tangent to any plane curve is obtained by the same means that the equation is obtained in analytical geometry, and is therefore independent of the failing cases of Taylor's theorem. The method of tangents naturally leads to the consideration of rectilinear asymptotes, which is, therefore, treated of in this chapter, and several examples are given, as well when the curve is referred to polar as to rectangular coordinates; and a few passing observations are made on the circular asymptotes to spiral curves, the chapter terminating with the differential expression for the arc of any plane curve determined without the aid of Taylor's theorem.

The Second chapter contains the theory of osculation, which is shewn to be unaffected by the failing cases of Taylor's theorem, although this is employed to establish the theory. The expressions for the radius of curvature are afterwards deduced, and several examples of their application given principally to the curves of the second order, and an instance of their utility shewn in determining the ratio of the earth's diameters.

The Third chapter is on involutes, evolutes, and consecutive curves, and contains some interesting theorems and practical examples. Of what the French call consecutive curves, I have endeavoured to give a clear and satisfactory explanation, unmixed with any vague notions about infinity.

The Fourth chapter is on the singular points of curves, and contains easy rules for detecting them, from an examination of the equation of the curve. This chapter also contains the general theory of curvilinear asymptotes, and completes the Second Section, or that assigned to the consideration of plane curves.

The Third Section is devoted to the general theory of curve surfaces, and of curves of double curvature; in the First chapter of which are established the several forms of the equations of the tangent plane and normal line at any point of a curve surface, and of the linear tangent and normal plane at any point of a curve of double curvature.

In the Second chapter the theory of conical and cylindrical surfaces is discussed, as also that of surfaces of revolution; and that remarkable case is examined, where the revolution of a straight line produces the same surface as the revolution of the hyperbola, whose asymptote is parallel to that line. Throughout this chapter are interspersed many valuable and interesting applications of the calculus, chiefly from *Monge*.

The Third chapter embraces the theory of the curvature of

surfaces in general, and will be found to form a collection of very beautiful theorems, the results, principally, of the researches of *Euler*, *Monge*, and *Dupin*. Most of these theorems have, however, usually been established by the aid of the *Infinitesimal Calculus*, or by the use of some other equally objectionable principle; they are here fairly deduced from the principles of the differential calculus, without, in any instance, departing from those principles, as laid down in the preliminary chapter. Those who are familiar with these enquiries, will find that I have obtained some of these theorems in a manner much more simple and concise than has hitherto been done. I need only mention here, as instances of this simplicity, the theorems of *Euler* and of *Meusnier*, at pages 227, 233.

The Fourth chapter is on *twisted surfaces*, a class of surfaces which have never been treated of, to any extent, by any English author, although, as has been recently shewn, the English were the first who noticed the peculiarities of certain individual surfaces belonging to this extensive class.* For what is here given, I am indebted to the French mathematicians—to *Monge* principally, and also to the *Chevalier Le Roy*, who has recently published a very neat and comprehensive treatise on curves and surfaces.

The Fifth chapter treats on the developable surfaces, or those which, like the cone and cylinder, may, if flexible, be unrolled upon a plane, without being twisted or torn. The Sixth chapter is on curves of double curvature; and the Seventh, which concludes the volume, contains a few miscellaneous propositions intimately connected with the theory of surfaces.

* See a paper in *Leybourn's Repository*, No. 22, by T. S. Davies, Esq. F.R.S., &c., of the Royal Military Academy.

From the foregoing brief analysis, it will appear evident to those familiar with the present state of mathematical instruction in this country, that I have introduced into one small volume, a more comprehensive view of the theory and applications of the Differential Calculus than has yet appeared in the English language. But I have aimed at more than this; and have endeavoured to simplify and improve much that I have adopted from foreign sources; and, above all, to establish every thing here taught, upon principles free from inconsistency and logical objections; and if it be found, upon examination, that I have entirely failed in this endeavour, I shall certainly feel a proportionate disappointment.

May 21, 1831.

The forgeoing remarks sufficiently set forth the scope and pretensions of the present work.

In this new edition the same comprehensiveness of plan is preserved; the details, in some instances, have been remodelled and amplified; several new articles and illustrative remarks, have been interspersed through the text; and the typographical blemishes of the former edition, as far as a careful examination has led to their detection, have all been removed from this. I have exercised an unremitting watchfulness over the press; and trust that I have thus secured the work from that inaccuracy with which, I regret to say, the former edition was justly chargeable.

J. R. YOUNG.

Belfast College;
August 1, 1836.

CONTENTS.

SECTION I.

On the Differentiation of Functions in general.

Article		*Page*
1.	A Function defined	1
2.	Effect produced on the function by a change in the variable	2
3.	Differential coefficient determined	3
4.	General form of the development of $f(x+h)$	7
5.	The coefficient of the second term in the general development is the differential coefficient derived from the function $f(x)$	9
6.	To differentiate the product of two or more functions of the same variable	10
7.	To differentiate any power of a function	12
8.	To differentiate a fraction	13
9.	To differentiate an expression consisting of several functions of the same variable	14
10.	Application of the preceding rules to examples	ib.
11.	Transcendental functions	18
12.	To find the differential of a logarithm	ib.
13.	To differentiate an exponential function	19
	Examples on Transcendental functions	ib.
14.	To differentiate circular functions	22
15.	Differentiation of inverse functions	27
16.	Forms of the differentials when the radius is arbitrary	31
17.	Successive differentiation explained	32
18.	Illustrations of the process	33
19.	Investigation of *Maclaurin's Theorem*	35
20.	Application of Maclaurin's theorem to the development of functions	37

Article *Page*

21. Deduction of *Euler's* expressions for the sine and cosine of an arc, by means of imaginary exponentials . . 41
22. *Demoivre's* formula, and series for the sine and cosine of a multiple arc 42
23. Decomposition of the expression $y^{2m} - 2y \cos \theta + 1$ into its quadratic factors 43
24. *Demoivre's* property of the circle 45
5. *Cotes's* properties of the circle 46
26. *John Bernoulli's* development of $(\sqrt{-1})^{\sqrt{-1}}$ . . 47
Developments of tan x and $\tan^{-1} x$ . . . 49
27. *Euler's* series for approximating to the circumference of a circle 51
28. *Machin's* more convergent series 52
29. Examples for exercise 53
30. Investigation of *Taylor's Theorem* 54
31. *Maclaurin's* theorem deduced from Taylor's . . 56
32. Application of Taylor's theorem to the development of functions ib.
33. Commodious form for the differential coefficient when the function is complicated 58
34. Examples of the application of this form . . 59
35. Form of the differential coefficient derived from the function $u = F(p, q)$, where p and q are functions of the same variable 60
36. Form of the coefficient when the function is $v = F(p, q, r)$. 61
37. Distinction between *partial* and *total* differential coefficients 62
38. Examples 63
39. Differentiation and development of *implicit functions* . 64
40. On vanishing fractions 68
41. On the erroneous views sometimes taken of this subject . 70
Ib. Application of the calculus to determine the true value of a vanishing fraction 72
42. Determination of the value when Taylor's theorem fails 76
43. Determination of the value of a fraction, of which both numerator and denominator are infinite . . . 79
44. Determination of the value of the product of two factors, when one becomes 0 and the other ∞ 80
45. Determination of the value of the difference of two functions, when they both become infinite . . . ib.
46. Examples on the preceding theory . . . 81
47. On the maxima and minima values of functions of a single variable 83

Article *Page*

48. If the function $F(a+h)$ be developed according to the ascending powers of h, a value so small may be given to h that any proposed term in the series shall exceed the sum of all that follow 84
49. Determination of the maxima and minima values in those cases where Taylor's theorem is applicable . . ib.
50. Determination of the values when Taylor's theorem is not applicable 87
51. Maxima and minima values which satisfy the condition $\frac{dy}{dx} = \infty$ 89
52. Conditions of maxima and minima, when the function is implicitly given 90
53. Precepts to abridge the process of finding the maxima and minima values 91
54. Examples 92
55. On the cautions to be observed in applying the analytical theory of maxima and minima to Geometry . . 103
56. Differentiation of functions of two independent variables . 105
57. Form of the differential when the function is implicit . 107
58. The ratio of the two partial differential coefficients derived from $u = F(z)$, z being a function of x and y, is independent of F 108
59. Development of functions of two independent variables . 109
60. The partial differential coefficients composing the coefficient of any term in the general development are identical with those arising from differentiating the preceding term . . 112
61. Maclaurin's theorem extended to functions of two independent variables 114
62. *Lagrange's Theorem* 115
63. Applications of Lagrange's theorem . . 118
64. Maxima and minima values of functions of two variables . 120
65. Examples 123
66. On changing the independent variable . . . 126
67. On the failing cases of Taylor's theorem . . 128
68. Explanation of the cause and extent of these failing cases . ib.
69. Particular examination of them . . . 130
70. Inferences from this examination . . . 131
71. The converse of these inferences true . . 132
72. To obtain the true development when Taylor's theorem fails . 133

Article *Page*

73. Correction of the errors of some analysts with respect to the failing cases of Taylor's theorem . . . 135

74. On the multiple values of $\frac{dy}{dx}$ in implicit functions . 137

75. Determination of these multiple values . . 138

76. Determination of the multiple values of $\frac{d^2y}{dx^2}$ . . 140

SECTION II.

Application of the Differential Calculus to the Theory of plane Curves.

77. Explanation of the method of tangents . . . 143

78. Equation of the normal 145

79. Application to curves related to rectangular coordinates . 146

80. Formulas for polar curves . . . 148

81. Applications to spirals 150

82. Expression for the perpendicular from the pole upon the tangent 152

83. Rectilinear asymptotes 153

84. Examples on the determination of asymptotes . . 154

85. Rectilinear asymptotes to spirals . . . 156

86. Circular asymptotes to spirals . . . 157

87. Expression for the differential of an arc of a plane curve . 158

88. Principles of osculation 159

89. Different orders of contact . . . 161

90. Nature of the contact at those points for which Taylor's development holds 162

91. Of the contact at the points for which Taylor's development fails 163

92. Osculating circle . . . . . ib.

93. Determination of the radius and direction of curvature . 164

94. The centres of touching circles all on the normal . . 166

95. Examples on the radius of curvature . . ib.

96. Expression for the radius of curvature of an ellipse applied to determining the ratio of the polar and equatorial diameters of the earth 168

Article *Page*

97. To determine those points in a given curve at which the osculating circle shall have contact of the third order . 170
98. Expression for the radius of curvature when the independent variable is arbitrary 171
99. Particular forms derivable from this general expression . 172
100. Suitable formula for polar curves . . . 173
101. Expression for the radius in terms of the radius vector and perpendicular from the pole on the tangent . . 175
102. Involutes and evolutes 176
103. Determination of the evolutes of several curves . 177
104. Normals to the involute are tangents to the evolute . 179
105. The difference of any two radii of curvature is equal to the arc of the evolute comprehended between them . . 180
106. On consecutive lines and curves . . . 182
107. Determination of the points of intersection of consecutive curves . . . . . ib.
108. Determination of the envelope of a family of curves . 184
109. Examples of this theory 185
110. Multiple points of curves 188
111. Determination of these points from the equation of the curve 189
112. Conjugate points 191
113. The determination of these points does not depend on Taylor's theorem . . . . . ib.
114. Multiple points of the second species . . ib.
115. Cusps, or points of regression . . . 193
116. Cusps exist only at points for which Taylor's theorem fails ib.
117. To distinguish a limit from a cusp . . . ib.
118. Examples of the determination of cusps whose branches touch an ordinate or an abscissa 194
119. Cusps whose branches touch a line oblique to the axes . 196
120. Conditions fulfilled by such cusps . . . ib.
121. Distinction between cusps of the first and those of the second kind . . . . . ib.
122. Examples 197
123. On points of inflexion 198
124. On curvilinear asymptotes 201

SECTION III.

On the general Theory of Curve Surfaces, and of Curves of Double Curvature.

Article		*Page*
125.	To determine the equation of the tangent plane at any point on a curve surface	204
126.	Form of the equation when the equation of the surface is implicit	205
127.	To determine the equation of the normal line at any point of a curve surface	206
128.	Expressions for the inclinations of the normal to the axes	ib.
129.	Forms of these expressions when the equation of the surface is implicit	207
130.	To determine the equation of the linear tangent at any point of a curve of double curvature	ib.
131.	To determine the equation of the normal plane at any point of a curve of double curvature	208
132.	To determine the equation of cylindrical surfaces in general	209
133.	General differential equation of cylindrical surfaces	210
134.	The same determined otherwise	ib.
135.	Given the equation of the generatrix to determine the cylindrical surface which envelopes a given curve surface	211
136.	If the enveloped surface be of the second order the curve of contact will be a plane curve and of the second order	ib.
137.	To determine the general equation of conical surfaces	213
138.	General differential equation of conical surfaces	ib.
139.	The same determined otherwise	214
140.	Given the position of the vertex, to determine the equation of the conical surface that envelopes a given curve surface	ib.
141.	*Monge's* proof that when the given curve surface is of the second order, the curve of contact is a plane curve	215
142.	Shorter method of proof	ib.
143.	*Davies's* proof that there is one point and only one from which as a vertex, if tangent cones be drawn to two surfaces of the second order, their planes of contact shall coincide	216

Article *Page*

144. The plane through the curve of contact is always conjugate to the diameter through the vertex of the cone . 217
145. Surfaces of revolution . . . . ib.
146. To determine the equation of surfaces of revolution in general 218
147. Simplified form of the equation when the axis of revolution coincides with the axis of z 219
148. Remarkable case, in which the generatrix is a straight line . ib.
149. General differential equation of surfaces of revolution . 220
150. A given curve surface revolves round a given axis, to determine the surface which touches and envelopes the moveable surface in every position 221
151. Example in the case of the spheroid . . 223
152. Characteristic property of developable surfaces . . 223
153. ——————— twisted surfaces . . 224
154. Osculation of curve surfaces . . . ib.
155. At any point on a curve surface to find the radius of curvature of a normal section 226
156. *Euler's* theorem, viz. at every point on a curve surface the sections of greatest and least curvature are always perpendicular to each other 227
157. Values of the radii of curvature of *any* perpendicular normal sections . . . . . ib.
158. Expressions for the radii of *greatest* and *least* curvatures . ib.
159. Peculiarities of the surface at the point where the principal radii have the same sign . . . 229
160. Peculiarities at the point where the principal radii are the same in length and in sign 230
161. Principal radii of curvature in *rule surfaces* . ib.
162. Peculiarities of the surface at the point where the principal radii have different signs . . . . ib.
163. Means of determining when the signs are different . 231
164. A paraboloid may always be found that shall have at its vertex the same curvature as any surface whatever at a given point ib.
165. To determine the radius of curvature at any point in an oblique section. The *theorem of Meusnier* . . . 233
166. Lines of curvature 234
167. To determine the lines of curvature through any point on a curve surface . . . . . ib.
168. Lines of curvature through any point are always perpendicular to each other 237

Article *Page*

169. On the developable surfaces, edges of regression, &c. generated by the normals to lines of curvature 238
170. Radii of spherical curvature 239
171. Given the coordinates of a point on a curve surface to determine the radii of spherical curvature at that point . . 240
172. The radius of curvature of an oblique section any how situated with respect to the surface and to the axes of coordinates is now determinable 241
173. To determine the radii of curvature at any point in a paraboloid 242
174. Twisted surfaces 243
175. To determine the surfaces generated by a straight line moving parallel to a fixed plane and along two rectilinear directrices not situated in one plane . . . . 244.
176. Two straight lines shown to pass through every point on the surface of a hyperbolic paraboloid . . . 246
177. To determine the surface generated by the motion of a straight line along three others fixed in position, so that no two of them are in the same plane . . . . ib.
178. To determine the surface generated when the directrices are not all parallel to the same plane . . . 247
179. Two straight lines shown to pass through every point on the surface of a hyperboloid of a single sheet . . 248
180. On twisted surfaces having but one curvilinear directrix 250
181. To determine the general equation of conoidal surfaces . ib.
182. Equation of the right conoid 251
183. To find the equation of the inferior surface of a winding staircase 252
184. To determine the differential equation of conoidal surfaces 253
185. The same determined otherwise . . . . ib.
186. Twisted surfaces having curvilinear directrices only . 254
187. To determine the general equation of surfaces generated by a straight line which moves along any two directrices whatever, and continues parallel to a fixed plane . . ib.
188. Determination of the differential equation of these surfaces 255
189. To determine the general equation of surfaces generated by the motion of a straight line along three curvilinear directrices 256
190. Application of the preceding theory 258
191. Determination of the equations of the intersections of consecutive surfaces 260

Article *Page*

192. Determination of the general equations of developable surfaces 261
193. To determine the developable surface generated by the intersection of normal planes at every point in a curve of double curvature 262
194. To determine the developable surface which touches and embraces two given curve surfaces . . . 263
195. To determine the differential equation of developable surfaces in general 264
196. The same determined otherwise . . . 265
197. Envelopes, characteristics, and edges of regression . ib.
198. The centre of a sphere of given radius moves along a given plane curve to determine the surface enveloping the sphere in every position 267
199. On curves of double curvature . . . 269
200. Expression for the differential of an arc of double curvature ib.
201. Osculation of curves of double curvature . . 270
202. Equation of the tangent deduced from the principles of osculation 271
203. To determine the osculating circle at any point in a curve of double curvature 272
204. General expression for the radius of absolute curvature . 274
205. Conditions necessary for the circle to have contact of the first order only . . . . . ib.
206. Another method of determining the osculating plane . 275
207. Another method of determining the osculating circle . 276
208. Other and more simple expressions for the coordinates of the centre of the osculating circle . . . 277
209. To determine the centre and radius of spherical curvature at any point in a curve of double curvature . . ib.
210. Determination of the equations of the edge of regression of the developable surfaces generated by the intersections of consecutive normal planes to the curve . . 278
211. To determine the points of inflexion in a curve of double curvature 279
212. On the evolutes of curves of double curvature . 280
213. Lines of poles . . . . . ib.
214. The locus of the poles the same as the locus of the characteristics . . . . . ib.
215. Every curve has an infinite number of evolutes all situated on the developable surfaces which is the locus of the poles . 281

Article *Page*

216. Every curvilinear evolute of a plane curve is a helix described on the surface of the cylinder which is the locus of the poles of the plane curve 282
217. The shortest distance between two points of an evolute is the arc of that evolute 283
218. Given the equations of a curve of double curvature to determine those of any one of its evolutes . . . . ib.
219. To prove that the locus of all the linear tangents at any point of a curve surface is necessarily a plane . . . 285
220. Given the algebraical equation of a curve surface to determine whether or not the surface has a centre . . . 286
221. To determine the equation of the diametral plane in a surface of the second order which will be conjugate to a given system of parallel chords 288
222. A straight line moves so that three given points in it constantly rest on the same three rectangular planes; required the surface which is the locus of any other point in it . . . 290
223. To determine the line of greatest inclination . . . 291
224. The six edges of any irregular tetraedron are opposed two by two, and the nearest distance of two opposite edges is called *breadth*; so that the tetraedron has three breadths and four *heights*. It is required to demonstrate that in every tetraedron the sum of the reciprocals of the squares of the breadths is equal to the sum of the reciprocals of the squares of the heights 294

NOTES . . . . . 297—312

THE

DIFFERENTIAL CALCULUS.

SECTION I.

ON THE

DIFFERENTIATION OF FUNCTIONS IN GENERAL.

CHAPTER I.

EXPLANATION OF FIRST PRINCIPLES.

Article (1.) All quantities which enter into calculation may be divided into two principal classes, *constant* quantities and *variable* quantities; the former class comprehending those which undergo no change of value, but remain the same throughout the investigation into which they enter; while those quantities which have no fixed or determinate value, constitute the latter class.

In algebra we usually employ the first letters, a, b, c, &c. of the alphabet, to represent *known* quantities, and the latter letters, z, y, x, &c. as symbols of the *unknown* quantities; but, in the higher calculus, the early letters are adopted as the symbols of *constant* quantities, whether they be known or unknown, and the latter letters are used to represent *variables*.

Any analytical expression composed of constants and variables, is said to be a *function* of the variables. Thus, if $y = ax^2 + bx + c$, then is y a function of x, because x enters into the expression for y; y is also a function of x in the expressions $y = a^x + b$, $y = \log x + ax^2$, &c.

and, as in each of these cases the *form* of the function is exhibited, y is said to be an *explicit* function of x; but, in such equations as

$$ax^2 + by^2 + cxy + x + y + e = 0,$$
$$x^3 + x^2y - ay^2 = y^3 + bx + c, \text{ \&c.}$$

where the form of the function that y is of x, can be ascertained only by solving the equation, y is an *implicit* function of x.

Similar remarks apply to the equations

$$z = ax^2 + by^2 + cx + e, \quad az^2 + by^2 + cxz + e = 0, \text{ \&c.}$$

z being an explicit function of x and y in the first, and an implicit function of the same variables in the second equation.

If we wish merely to express that y is an explicit function of x, without writing the form of that function, we adopt the notation $y = \mathrm{F}(x)$, or $y = f(x)$, or $y = \phi(x)$, &c.; and, to denote an implicit function, we write $\mathrm{F}(x, y) = 0$, $f(x, y) = 0$, &c.

(2.) Let us now examine the effect produced on the function y, by a change taking place in the variable x; and, for a first example let us take the equation $y = mx^2$.

Changing, then, x into $x + h$, and representing the corresponding value of y by y', we have

$$y' = m(x + h)^2,$$

or, by developing the second member,

$$y' = mx^2 + 2mxh + mh^2.$$

As a second example, let us take the equation $y = x^3$; and putting, as before, y' for the value of the function, when x is changed into $x + h$, we have

$$y' = (x + h)^3 = x^3 + 3x^2h + 3xh^2 + h^3.$$

We thus see, in these two examples, the effect produced on the function by changing the value of the variable. On account of this dependence of the value of the function upon that of the variable the former, that is y, is called the *dependent* variable, and the latter, x, the *independent* variable.

Let us now ascertain the difference of the values of each of the above functions of x, in the two states y and y'. In the first example,

$$y' - y = 2mxh + mh^2.$$

In the second,

$$y' - y = 3x^2h + 3xh^2 + h^3;$$

so that, in the equation $y = mx^2$, if h be the increment of the variable x, we see that $2mxh + mh^2$ will be the corresponding increment of the function y; and, in the equation $y = x^3$, if x take the increment h, the corresponding increment of the function will be $3x^2h + 3xh^2 + h^3$.

We may, therefore, in each of these cases, readily find an expression for the ratio of the increment of the function to that of the variable, that is to say, the value of the fraction $\frac{y' - y}{h}$.

In the first case,

$$\frac{y' - y}{h} = 2mx + mh \ldots . (1).$$

In the second,

$$\frac{y' - y}{h} = 3x^2 + 3xh + h^2 \ldots . (2).$$

It is here worthy of remark, that, in both these expressions for the ratio, the first term is independent of h; so that, however we alter the value of h, this first term will remain unchanged. If, therefore, h be supposed to diminish continually, and, at length to become 0, the said first term will then express the value of the ratio in this extreme case. This first term, then, is the *limit* to which the ratio approaches as h diminishes, but which limit it cannot attain till h becomes absolutely 0

In the first of the foregoing examples, $2mx$ is the limit of the ratio $\frac{y' - y}{h}$; or it is the value towards which this ratio continually approaches when h is continually diminished, and at which it ultimately arrives when these continual diminutions bring it at length to $h = 0$. In the second example the limit is $3x^2$.

(3.) We may now understand what is meant by the *limit of the ratio of the increment of the function to that of the variable.* It is the

determination of this limit, in every possible form of the function, that is the principal object of the differential calculus. The limit itself is called the *differential coefficient*, derived from the function; so that, if the function be mx^2, the differential coefficient, as we have seen above, is $2mx$, and the differential coefficient, derived from the function x^3, is $3x^2$.

In both these cases, as indeed in every other, the respective differential coefficients are only so many particular values of the general symbol $\frac{0}{0}$, to which $\frac{y'-y}{h}$ always reduces when $h=0$. In the first example above, $\frac{0}{0}=2mx$; in the second, $\frac{0}{0}=3x^2$.

It appears, therefore, that whatever conflicting opinions may prevail among mathematicians, as to the meaning of the symbol $\frac{0}{0}$ when it occurs in an analytical result, they do not, in the least, affect the precision of the fundamental principles of the Differential Calculus. For here it has a distinct and definite interpretation: it is the symbolical expression for the limit of the value of $\frac{y'-y}{h}$; or, in other words, it represents, solely and exclusively, what the development of $\frac{y'-y}{h}$ becomes when $h=0$. It must not be forgotten that the expression $\frac{y'-y}{h}$ implies an *operation* to be performed, viz. division; and although, in conformity with ordinary usage, and to avoid circumlocution, we have called this expression a *ratio* or *quotient*, yet the thing really meant is the result of that operation; just as we call $\sqrt{a}$ "the square root of a," although $\sqrt{}$ indicates an unperformed process.

In each of the equations (1), (2), given above, the right hand member is the correct interpretation of the left; and although, for $h=0$, this left hand member takes the form $\frac{0}{0}$, which, in an isolated state, is altogether meaningless, or may mean anything, yet, being accompanied by its interpretation on the right, it becomes perfectly intelligible and unambiguous.

Instead, however, of the general symbol $\frac{0}{0}$, a particular notation is employed to represent the limiting ratio, or differential coefficient, in

each particular case; thus, if y is the function, and x the independent variable, the differential coefficient is represented thus, $\frac{dy}{dx}$. If z were the function, and y the independent variable, the differential coefficient would be $\frac{dz}{dy}$; the expressions $\frac{dy}{dx}$ and $\frac{dz}{dy}$ have, we see, the advantage over the symbol $\frac{0}{0}$, of particularizing the function and the independent variable under consideration, and of thus restricting its signification; and this, it must be remembered, is all that distinguishes $\frac{dy}{dx}$ or $\frac{dz}{dy}$ from $\frac{0}{0}$, for dy, dz, dx, are each absolutely 0.

This notation being agreed upon, we have, when $y = mx^2$,

$$\frac{dy}{dx} = 2mx,$$

and, when $y = x^3$,

$$\frac{dy}{dx} = 3x^2.$$

As a third example, let the function $y = a + 3x^2$ be proposed, then, changing x into $x + h$, and y into y', we have

$$y' = a + 3x^2 + 6xh + 3h^2,$$

$$\therefore \frac{y' - y}{h} = 6x + 3h;$$

and, making $h = 0$, we have, for the differential coefficient,

$$\frac{dy}{dx} = 6x.$$

If in this example the function had been $3x^2$, instead of $a + 3x^2$, the differential coefficient would obviously have been the same.

As a fourth example, let $y = ax^2 \pm b$,

$$\therefore y' = ax^2 \pm b + 2axh + ah^2,$$

$$\therefore \frac{y' - y}{h} = 2ax + ah \quad \therefore \frac{dy}{dx} = 2ax,$$

which would have been the same if the constant b had not entered the function.

As a last example, take the function $y = (a + bx)^2$ or $y = a^2 + 2abx + b^2x^2$, which, when x is changed into $x + h$, becomes

$$y' = a^2 + 2ab(x + h) + b^2(x + h)^2$$

$$= a^2 + 2abx + b^2x^2 + 2(ab + b^2x)h + b^2h^2,$$

$$\therefore \frac{y' - y}{h} = 2b(a + bx) + b^2h,$$

$$\therefore \frac{dy}{dx} = 2b(a + bx).$$

It should be remarked, that of the two parts dy, dx, of which the symbol $\frac{dy}{dx}$ consists, the former is called the differential of y, and the latter the differential of x. These differentials, although each $= 0$, have, nevertheless, as we have already seen, a determinate relation to each other; thus, in the last example, this relation is such, that $dy = 2b(a + bx)\,dx$, and, although this is the same as saying that $0 = 2b(a + bx) \times 0$, yet, as we can always immediately obtain from this form the true value of $\frac{0}{0}$ or $\frac{dy}{dx}$, we do not hesitate occasionally to make use of it.

From the expression for the differential of a function, we readily see the propriety of calling $\frac{dy}{dx}$ a *coefficient*, being, indeed, the coefficient of dx.

CHAPTER II.

DIFFERENTIATION OF FUNCTIONS OF ONE VARIABLE.

(4.) Let $f(x)$ represent any function of x whatever; then, if x be changed into $x + h$, the general form of the development of $f(x + h)$, arranged according to the powers of h, will be

$$f(x + h) = f(x) + Ah + Bh^2 + Ch^3 + Dh^4 + \&c.,$$

as may be proved as follows:

1. *The first term of the development must be* $f(x)$. This is obvious, for this first term is what the whole development reduces to when $h = 0$, but we must in this case have the identity $f(x) = f(x)$; hence, $f(x)$ is the first term.

2. *None of the exponents of* h *can be fractional.* For if any exponent were fractional, the term into which it enters would be irrational; so that the development of $f(x + h)$, containing an irrational term, $f(x + h)$ itself would be irrational, since it is impossible that there should be an equality between an irrational expression and one that is rational. But, if $f(x + h)$ were irrational, $f(x)$ would likewise be irrational; for the former function differs from the latter only in this, that $x + h$ occupies in it the place of x in the latter; and, therefore, as many values as $f(x)$ has, in consequence of the radicals that may enter into it, so many values, and no more, must $f(x + h)$ have. As, therefore, the first term of the development of $f(x + h)$ has the same number of values as $f(x + h)$ itself, none of the succeeding terms can contain a fractional power of h; for, if any term could, that term would have multiple values, each of which would combine with the values of $f(x)$, and thus the development would have more values than its equivalent function, which is absurd.

3. *None of the exponents of* h *can be negative* in the general development. For, on the supposition tbat a negative power of h enters any term, that term would become infinite when $h = 0$; but, when $h = 0$, the function is simply $f(x)$, which is not necessarily infinite; so that

that development into which a negative power of h enters, cannot be the *general development* of $f(x+h)$, understanding, by the general development, that which does not restrict x to any particular value or values. By supposing a negative power of h to appear in the development, we have just seen that such development would restrict x to the values determined, by the equation $f(x)=\infty$, or by the equation $\frac{1}{f(x)}=0$, to the exclusion of all other values, and is, therefore, not general.

As to the method of determining the coefficients A, B, C, &c., that will be investigated hereafter (in Chapter IV.)

4. It must be here remarked, that, although the general development of every function of x is of the above form, yet we are not to suppose that this form will remain unchanged whatever particular value we give to x; for particular values may be so chosen as to render this form of development, with *finite coefficients*, impossible, and such impossibility will be intimated by the assumed value of x rendering some of the coefficients A, B, C, &c. infinite. The well-known binomial theorem, which we already know to be of the above form, will afford an illustration of this. Thus the general development of $f(x)=\sqrt{x+a}$, when we replace x by $x+h$, is, by the above-mentioned theorem,

$$f(x+h)=\sqrt{(x+a)+h}=$$

$$(x+a)^{\frac{1}{2}}+\tfrac{1}{2}(x+a)^{-\frac{1}{2}}h-\tfrac{1}{8}(x+a)^{-\frac{3}{2}}h^2+\&c.$$

where, in the case $x=-a$, all the coefficients become infinite, and the development, according to the positive integral powers of h, is in this case impossible; for the function then becomes merely $\sqrt{h}$ or $h^{\frac{1}{2}}$, in which the exponent of h is fractional. The impossibility of the proposed form of development in such particular cases is always intimated, as in the example just adduced, by the circumstance of infinite coefficients entering it; for imaginary coefficients would imply merely that the function $f(x+h)$, for the assumed value of x, becomes imaginary, and not that the development failed. A particular examination of the cases in which the general form of the development fails to have place, will form the subject of a future chapter; at present it is sufficient to apprize the student that such failing cases may exist.

(5.) By transposing the first term in the general development of $f(x+h)$, at art. (4), we have

$$f(x+h)-f(x)=Ah+Bh^2+Ch^3+\&c.$$

$$\therefore \frac{f(x+h)-f(x)}{h}=A+Bh+Ch^2+\&c.$$

hence, when $h=0$,

$$\frac{df(x)}{dx}=A,$$

from which result we learn, that *the coefficient of the second term, in the development of the function* $f(x+h)$, *is the differential coefficient derived from the function* $f(x)$; so that the finding the differential coefficient from any proposed function, $f(x)$, reduces itself to the finding the coefficient of the second term in the general development of $f(x+h)$, or of the first term in the developed difference $f(x+h)-f(x)$.

Having obtained this general result, we may now proceed to apply it to functions of different forms; but it will be proper previously to observe, that those constants which are connected with the variable in the function $f(x)$, only by way of addition or subtraction, cannot appear in the coefficient A; because A, being multiplied by h, can contain no quantities which are not among those multiplied by $x+h$ in $f(x+h)$, or by x in $f(x)$.

Another obvious principle, which it may be proper here to mention, is, that if two functions of any variable, x, continue equal for all values of x, the differential coefficient derived from the one must be equal to that derived from the other. For the functions being always equal, they are equal when $x+h$ is substituted for x, whatever be the value of h; hence their developments are equal whatever be the value of h, and consequently, by the method of indeterminate coefficients, the two developments are identical, term and term; so that the coefficient of h, in the one development, is equal to the coefficient of h in the other. In *identical* equations, therefore, we may always equate the differential coefficient deduced from one member, with that deduced from the other. But in equations which are not identical, and which therefore restrict the variable to particular values, such a deduction would evidently be

erroneous. In fact, the so called *variable* is not, in the case supposed, a variable according to our definition, but a constant, which we can neither increase or diminish without violating the condition implied in the equation. It is important that the student keep this in mind, recollecting that although he may perform upon each side of an equation any of the ordinary algebraic operations, and still be led to an equation, yet the operation called *differentiation* must be excepted, unless indeed the proposed equation be something more than an equality :—an *identity*.

We shall now investigate the fundamental rules of differentiation, or methods by which we may rapidly determine the second term of the development of $f(x+h)$; a matter of great importance, because, as will be seen hereafter, this second term furnishes a ready key for the discovery of all the following terms of the development.

(6.) *To differentiate the product of two or more functions of the same variable.*

Let y, z, be functions of x, in the expression

$$u = ayz.$$

By changing x into $x+h$, the function y becomes

$$y' = y + Ah + Bh^2 + Ch^3 + \&c. \ldots\ldots (1);$$

and the function z becomes

$$z' = z + A'h + B'h^2 + C'h^3 + \&c. \ldots\ldots (2).$$

Hence, when $h=0$, we have, from (1),

$$\text{limit of } \frac{y'-y}{h} = \frac{dy}{dx} = A,$$

and from (2)

$$\text{limit of } \frac{z'-z}{h} = \frac{dz}{dx} = A'.$$

Multiplying the product of (1) and (2) by a, we have

$$u' = ayz + a\,(Az + A'y)\,h + \&c.$$

$$= ayz + a\,(\frac{dy}{dx}\,z + \frac{dz}{dx}\,y)\,h + \&c.$$

therefore, $a\,\frac{dy}{dx}\,z + a\,\frac{dz}{dx}\,y$ being the coefficient of the second term of the development of u', we have

$$\frac{du}{dx} = az\,\frac{dy}{dx} + ay\,\frac{dz}{dx}$$

$$\therefore\ du = azdy + aydz \ \ldots\ (3).$$

Hence, *to differentiate the product of two functions of the same variable, we must multiply each by the differential of the other, and add the results.*

It will be easy now to express the differential of a product of three functions of the same variable. Let

$$u = wyz$$

be the product of three functions of x; then, putting v for wy, the expression is

$$u = vz;$$

hence, by (3),

$$du = zdv + vdz,$$

but $v = wy$; therefore, by (3), $dv = ydw + wdy$; consequently, by substitution,

$$du = zydw + zwdy + wydz \ \ldots\ (4),$$

and it is plain that in this way the differential may be found, be the factors ever so many; so that, generally, *to differentiate a product of several functions of the same variable, we must multiply the several differentials of each factor by all the other factors, and add the results.*

If we suppose the factors to be all equal to each other, we shall obtain a rule to differentiate a positive integral power. Thus the differential of the function

$$u = x^m = x \cdot x \cdot x \cdot x \ldots .$$

is

$$du = x^{m-1}\,dx + x^{m-1}\,dx + x^{m-1}\,dx + \text{&c. to } m \text{ terms},$$

that is,

$$du = mx^{m-1}\,dx \quad \therefore \quad \frac{du}{dx} = mx^{m-1}.$$

The form of the differential is preserved whether m be integral or fractional, positive or negative; as appears from the following article:

(7.) *To differentiate any power of a function.*

The form of the differential when the power is whole and positive has been already established above. Let then

$$u = y^{\frac{m}{n}}$$

be proposed, y being a function of x, and $\frac{m}{n}$ being a positive fraction.

Since $u^n = y^m$,

$$\therefore \quad nu^{n-1}\,du = my^{m-1}\,dy,$$

$$\therefore \quad du = \frac{m}{n} \cdot \frac{y^{m-1}}{u^{n-1}}\,dy = \frac{m}{n} \cdot \frac{y^{m-1}}{y^{\frac{mn-m}{n}}}\,dy.$$

Now

$$(m-1) - \frac{mn-m}{n} = \frac{m}{n} - 1,$$

consequently,

$$du = \frac{m}{n}\, y^{\frac{m}{n}-1}\,dy.$$

Let now the exponent be negative, or

$$u = y^{-\frac{m}{n}}$$

$$\therefore \quad u^n = y^{-m} = \frac{1}{y^m}$$

$$\therefore \quad u^n y^m = 1 \quad \therefore \quad u^n\,dy^m + y^m\,du^n = 0,$$

$$\therefore\ mu^n y^{m-1}\,dy + ny^m u^{n-1}\,du = 0$$

$$\therefore\ du = -\frac{m}{n}\cdot\frac{u^n y^{m-1}}{y^m u^{n-1}}\,dy$$

$$= -\frac{m}{n}vy^{-1}\,dy$$

$$= -\frac{m}{n}y^{-\frac{m}{n}-1}\,dy.$$

Hence, generally, *to differentiate a power, we must multiply together these three factors, viz. the index of the power, the power itself diminished by unity, and the differential of the root.*

This rule might have been deduced with less trouble, by availing ourselves of the binomial theorem; for, supposing, in $u = y^p$, that the increment of the function y becomes k, when the increment of x becomes h, we have $u' = (y+k)^p$; and, by the binomial theorem, the coefficient of the second term of the expansion of $(y+k)^p$ is py^{p-1}, whether p be positive or negative, whole or fractional. As, however, we propose to demonstrate the binomial theorem by means of the differential calculus, we have thought it necessary to establish the fundamental principles of differentiation, independently of this theorem.

(8.) *To differentiate a fraction.* Let $u = \frac{y}{z}$, y and z being functions of x; therefore $uz = y$, $\therefore\ du\,z = dy$, that is, by article (6),

$$z\,du + u\,dz = dy\ \therefore\ du = \frac{dy - u\,dz}{z},$$

or, substituting $\frac{y}{z}$ for u,

$$du = \frac{z\,dy - y\,dz}{z^2}$$

Hence, to differentiate a fraction, the rule is this: *From the product of the denominator, and differential of the numerator, subtract the product of the numerator, and differential of the denominator, and divide the remainder by the square of the denominator.*

(9.) If it be required to differentiate an expression consisting of several functions of the same variable, combined by addition or subtraction, it will be necessary merely to differentiate each separately, and to connect together the results by their respective signs. For let the expression be

$$u = aw + by + cz + \&c.$$

in which w, y, z, are functions of x. Then, changing x into $x + h$ and developing,

$$\begin{aligned}
w &\text{ becomes } w + \mathrm{A}h + \mathrm{B}h^2 + \&c.\\
y &\qquad\quad y + \mathrm{A}'h + \mathrm{B}'h^2 + \&c.\\
z &\qquad\quad z + \mathrm{A}''h + \mathrm{B}''h^2 + \&c.\\
\therefore\ u &\qquad\quad u + (a\mathrm{A} + b\mathrm{A}' + c\mathrm{A}'' + \&c.)\,h + \&c.\\
\therefore\ \mathrm{d}u &= a\mathrm{A}\mathrm{d}x + b\mathrm{A}'\mathrm{d}x + c\mathrm{A}''\mathrm{d}x + \&c.
\end{aligned}$$

But

$$\mathrm{A}\mathrm{d}x = \mathrm{d}w,\ \mathrm{A}'\mathrm{d}x = \mathrm{d}y,\ \mathrm{A}''\mathrm{d}x = \mathrm{d}z,\ \&c.$$

therefore

$$\mathrm{d}u = a\mathrm{d}w + b\mathrm{d}y + c\mathrm{d}z + \&c.$$

that is, *the differential of the sum of any number of functions is equal to the sum of their respective differentials.*

(10.) We shall now apply the foregoing general rules to some examples, taking first those only which are purely algebraic, and afterwards extending the rules to the functions $\log x$, a^x, $\sin x$, &c.

EXAMPLES.

1. Let it be required to differentiate the function

$$y = 8x^4 - 3x^3 - 5x.$$

By the rule for powers (7) the differential of $8x^4$ is $8 \times 4x^3\mathrm{d}x$, and the differential of $-3x^3$ is $-3 \times 3x^2\mathrm{d}x$; also the differential of $-5x$ is $-5\mathrm{d}x$; hence (9),

$$\mathrm{d}y = 32x^3\mathrm{d}x - 9x^2\mathrm{d}x - 5\mathrm{d}x,$$

$$\therefore \frac{dy}{dx} = 32x^3 - 9x^2 - 5.$$

From this result we infer that whatever be the entire development of the function $8(x+h)^4 - 3(x+h)^3 - 5(x+h)$, the first two terms must be

$$(8x^4 - 3x^3 - 5x) + (32x^3 - 9x^2 - 5)h.$$

2. Let $y = (x^3 + a)(3x^2 + b)$.

By the rule for differentiating a product (6), we have

$$dy = (x^3 + a)\, d(3x^2 + b) + (3x^2 + b)\, d(x^3 + a),$$

and (7),

$$d(3x^2 + b) = 6x dx, \quad d(x^3 + a) = 3x^2 dx,$$

$$\therefore dy = (x^3 + a)\, 6x dx + (3x^2 + b)\, 3x^2 dx,$$

$$\therefore \frac{dy}{dx} = 15x^4 + 3bx^2 + 6ax.$$

Hence the development of the proposed function, when x is changed into $x + h$, is, as far as two terms,

$$y' = (x^3 + a)(3x^2 + b) + (15x^4 + 3bx^2 + 6ax)h + \&c.$$

3. Let $y = (ax + x^2)^2$.

The differential of the root $ax + x^2$ of this power, is $a dx + 2x dx$, therefore

$$dy = 2(ax + x^2)(a + 2x)\, dx,$$

$$\therefore \frac{dy}{dx} = 2(ax + x^2)(a + 2x).$$

4. Let $y = \sqrt{a + bx^2}$.

The differential of the root or function under the radical, is $2bx dx$; hence

$$dy = \tfrac{1}{2}(a + bx^2)^{-\frac{1}{2}} 2bx dx = \frac{bx}{\sqrt{a + bx^2}} dx,$$

$$\therefore \frac{dy}{dx} = \frac{bx}{\sqrt{a + bx^2}}.$$

5. Let $y = (a + bx^m)^n$.

The differential of the root or function within the parenthesis, is $mbx^{m-1}\,dx$; hence

$$dy = n\,(a + bx^m)^{n-1}\,mbx^{m-1}\,dx,$$

$$\therefore \frac{dy}{dx} = bmn\,(a + bx^m)^{n-1}\,x^{m-1}.$$

6. Let $y = \dfrac{x^2}{(a + x^3)^2}$.

The differential of the numerator of this fraction is $2x\,dx$, and the differential of $a + x^3$ is $3x^2dx$, therefore the differential of the denominator is $2\,(a + x^3)\,3x^2dx$; hence (8),

$$dy = \frac{(a + x^3)^2\,2x\,dx - 6x^4\,(a + x^3)\,dx}{(a + x^3)^4} = \frac{2ax - 4x^4}{(a + x^3)^3}\,dx,$$

$$\therefore \frac{dy}{dx} = \frac{2x\,(a - 2x^3)}{(a + x^3)^3}.$$

7. Let $y = \{a + \sqrt{(b + \frac{c}{x^2})}\}^4$.

The differential of the root $a + \sqrt{(b + \frac{c}{x^2})}$ is $\frac{1}{2}\,(b + \frac{c}{x^2})^{-\frac{1}{2}}\,d\,\frac{c}{x^2}$,

and $d\,\frac{c}{x^2} = -\frac{2c}{x^3}\,dx$; hence

$$\frac{dy}{dx} = -4\,\{a + \sqrt{(b + \frac{c}{x^2})}\}^3\,(b + \frac{c}{x^2})^{-\frac{1}{2}}\,\frac{c}{x^3} = -\frac{4c\,\{a + \sqrt{(b + \frac{c}{x^2})}\}^3}{x^3\sqrt{b + \frac{c}{x^2}}}$$

8. Let $y = \sqrt{x^2 + \sqrt{a + x^2}}$.

The differential of $x^2 + \sqrt{a + x^2}$ is $2x\,dx + (a + x^2)^{-\frac{1}{2}}\,x\,dx$,

$$\therefore \frac{dy}{dx} = \frac{x}{\sqrt{x^2 + \sqrt{a + x^2}}} + \frac{x}{2\sqrt{(a + x^2)\,(x^2 + \sqrt{a + x^2})}}.$$

9. Let $y = \dfrac{x}{\sqrt{a^2 + x^2} - x}$.

Multiplying numerator and denominator by $\sqrt{a^2 + x^2} + x$, the expression becomes

$$y = \frac{x^2}{a^2} + \frac{x}{a^2}\sqrt{a^2 + x^2},$$

$$\therefore \mathrm{d}y = \mathrm{d}\,\frac{x^2}{a^2} + \frac{\sqrt{a^2 + x^2}}{a^2}\,\mathrm{d}x + \frac{x}{a^2}\,\mathrm{d}\sqrt{a^2 + x^2},$$

$$\therefore \frac{\mathrm{d}y}{\mathrm{d}x} = \frac{2x}{a^2} + \frac{\sqrt{a^2 + x^2}}{a^2} + \frac{x^2}{a^2\sqrt{a^2 + x^2}}$$

$$= \frac{2x}{a^2} + \frac{a^2 + 2x^2}{a^2\sqrt{a^2 + x^2}}.$$

10. $y = a^2 - x^2$ $\therefore \dfrac{\mathrm{d}y}{\mathrm{d}x} = -2x.$

11. $y = 4x^3 - 2x^2 + 7x + 3$ $\therefore \dfrac{\mathrm{d}y}{\mathrm{d}x} = 12x^2 - 4x + 7.$

12. $y = (a + bx)\,x^3$ $\therefore \dfrac{\mathrm{d}y}{\mathrm{d}x} = 3ax^2 + 4bx^3.$

13. $y = (a + bx + cx^2 + \&c.)^m$ $\therefore \dfrac{\mathrm{d}y}{\mathrm{d}x} = m\,(a + bx + cx^2 + \&c.)^{m-1}\,(b + 2cx + \&c.)$

14. $y = (a + bx^2)^{\frac{5}{4}}$ $\therefore \dfrac{\mathrm{d}y}{\mathrm{d}x} = \dfrac{5bx}{2}\sqrt[4]{a + bx^2}.$

15. $y = a + \dfrac{4\sqrt{x}}{3 + x^2}$ $\therefore \dfrac{\mathrm{d}y}{\mathrm{d}x} = \dfrac{6\,(1 - x^2)}{(3 + x^2)^2\sqrt{x}}.$

16. $y = a + b\sqrt{x} - \dfrac{c}{x}$ $\therefore \dfrac{\mathrm{d}y}{\mathrm{d}x} = \dfrac{b}{2\sqrt{x}} + \dfrac{c}{x^2}.$

17. $y = (ax^3 + b)^2 + 2\sqrt{a^2 - x^2}\,(x - b)$ $\therefore \dfrac{\mathrm{d}y}{\mathrm{d}x} = 6ax^2(ax^3 + b) + \dfrac{2\,(a^2 - 2x^2 + bx)}{\sqrt{a^2 - x^2}}.$

$$18.\ y = \frac{x}{x + \sqrt{1 - x^2}} \therefore \frac{dy}{dx} = \frac{1}{\sqrt{1 - x^2}\,[x + \sqrt{1 - x^2}]^2}.$$

The functions in these examples are all *algebraic;* we shall now consider

Transcendental Functions.

(11.) Transcendental functions are those in which the variable enters in the form of an exponent, a logarithm, a sine, &c. Thus, a^x, $a \log x$, $\sin x$, &c., are transcendental functions: the first is an *exponential* function, the second a *logarithmic* function, and the third a *circular* function.

To find the Differential of a Logarithm.

(12.) Let it be required to differentiate $\log x$.

Put a for the base of the system of logarithms used, and let

$$M = \frac{1}{a - 1 - \frac{1}{2}(a-1)^2 + \frac{1}{3}(a-1)^3 - \&c.},$$

then, (*Algebra*, p. 186,)

$$\log(1 + n) = M(n - \tfrac{1}{2}n^2 + \tfrac{1}{3}n^3 - \&c.),$$

or, putting $\frac{h}{x}$ for n,

$$\log \frac{x+h}{x} = \log(x+h) - \log x = M\left(\frac{h}{x} - \frac{h^2}{2x^2} + \frac{h^3}{3x^3} - \&c.\right)$$

$$\therefore \frac{\log(x+h) - \log x}{h} = M\left(\frac{1}{x} - \frac{h}{2x^2} + \frac{h^2}{3x^3} - \&c.\right)$$

This is the general expression for the ratio of the increment of the function to that of the variable. Hence, taking the limit of this ratio, we have

$$\frac{d \log x}{dx} = \frac{M}{x} \quad \ldots\ (1).$$

If the logarithms employed be hyperbolic $M = 1$, and then

$$\frac{d \log x}{dx} = \frac{1}{x} \quad \ldots \quad (2).$$

If they are not hyperbolic, write Log instead of log, for distinction sake; then, since by putting a for $1 + n$ in the series for $\log (1 + n)$, we have

$$\log a = a - 1 - \tfrac{1}{2}(a-1)^2 + \tfrac{1}{3}(a-1)^3 - \&c. = \frac{1}{M},$$

it follows, from the expression (1), that

$$\frac{d \operatorname{Log} x}{dx} = \frac{1}{x \log a}.$$

Unless the contrary is expressed, the differential is always taken according to the hyperbolic system, because the expression is then simpler, $\log a$ being $= 1$.

From the preceding investigation we learn, that *the differential of a logarithm of a function is equal to the differential of the function divided by the function itself.*

(13.) *To differentiate an exponential function.*

1. Let $y = a^x$, then $\log y = x \log a$ $\therefore$ $d \log y = dx \log a$, that is, $\frac{dy}{y} = dx \log a$ $\therefore$ $dy = y \log a dx = \log a \cdot a^x dx$.

Hence, *to differentiate an exponential, we must multiply together the hyp. log of the base, the exponential itself, and the differential of the variable exponent.*

EXAMPLES.

1. Let $y = x(a^2 + x^2)\sqrt{a^2 - x^2}$ $\therefore$ $\log y = \log x + \log (a^2 + x^2) + \frac{1}{2} \log (a^2 - x^2)$,

$$\therefore \frac{dy}{y} = \frac{dx}{x} + \frac{2x dx}{a^2 + x^2} - \frac{x dx}{a^2 - x^2} = \frac{a^4 + a^2 x^2 - 4x^4}{x(a^2 + x^2)(a^2 - x^2)} dx,$$

therefore, substituting for y its value, we have

$$\frac{dy}{dx} = \frac{a^4 + a^2x^2 - 4x^4}{\sqrt{a^2 - x^2}}.$$

2. $y = \log \dfrac{\sqrt{a+x} + \sqrt{a-x}}{\sqrt{a+x} - \sqrt{a-x}}.$

Multiplying numerator and denominator by the denominator, the expression becomes

$$y = \log \frac{2x}{2a - 2\sqrt{a^2 - x^2}} = \log x - \log (a - \sqrt{a^2 - x^2})$$

$$\therefore \frac{dy}{dx} = \frac{1}{x} - \frac{x}{\sqrt{a^2 - x^2}\{a - \sqrt{a^2 - x^2}\}} = \frac{a\sqrt{a^2 - x^2} - a^2}{\sqrt{a^2 - x^2}\{a - \sqrt{a^2 - x^2}\}}$$

$$= \frac{-a}{\sqrt{a^2 - x^2}}.$$

3. $y = \dfrac{\sqrt{x^2 + 2ax}}{\sqrt[3]{x^3 + x^2 - x}}$ $\therefore \log y = \frac{1}{2} \log (x^2 + 2ax) - \frac{1}{3} \log (x^3 + x^2 - x)$,

$$\therefore \frac{dy}{y} = \frac{x + a}{x^2 + 2ax}\,dx - \frac{3x^2 + 2x - 1}{3(x^3 + x^2 - x)}\,dx =$$

$$\frac{(1 - 3a)x^2 - (a + 2)x - a}{3x(x^2 + x - 1)(x + 2a)}\,dx,$$

$$\therefore \frac{dy}{dx} = \frac{\{(1 - 3a)x^2 - (a + 2)x - a\}\sqrt{x}}{3x^{\frac{4}{3}}(x^2 + x - 1)^{\frac{4}{3}}(x + 2a)^{\frac{1}{2}}}.$$

4. $y = x^{m\sqrt{-1}}$ $\therefore \log y = m\sqrt{-1} \log x$ $\therefore \dfrac{dy}{y} = m\sqrt{-1}\,\dfrac{dx}{x}$,

$$\therefore \frac{dy}{dx} = m\frac{y}{x}\sqrt{-1} = m\sqrt{-1}\,.\,x^{m\sqrt{-1}-1}.$$

From this example it appears, that the rule at (7) applies when the exponent is imaginary.

5. $y = a^{x^x}$.

In this example the variable exponent is x^x; hence, calling it z and taking the logarithms, we have

$$\log z = x \log x \therefore \frac{dz}{z} = \log x dx + \frac{x dx}{x} \therefore dz = x^x (1 + \log x)\, dx;$$

hence, by the rule,

$$\frac{dy}{dx} = \log a \cdot a^{x^x} \cdot x^x (1 + \log x).$$

6. $y = e^{x\sqrt{-1}} + e^{-x\sqrt{-1}}$, where e is the base of the hyperbolic system,

$$\therefore dy = e^{x\sqrt{-1}} \sqrt{-1}\, dx - e^{-x\sqrt{-1}} \sqrt{-1}\, dx,$$

$$\therefore \frac{dy}{dx} = (e^{x\sqrt{-1}} - e^{-x\sqrt{-1}}) \sqrt{-1}.$$

7. $y = \log (\log x)$.* Put z for $\log x \therefore y = \log z \therefore dy = \frac{dz}{z}$; but $dz = d \log x = \frac{dx}{x} \therefore dy = \frac{dx}{x \log x} \therefore \frac{dy}{dx} = \frac{1}{x \log x}$.

8. $y = (\log x^n)^m \therefore \frac{dy}{dx} = \frac{mn (\log x^n)^{m-1}}{x}$.

9. $y = \log \{(a + x)^m (a' + x)^{m'} (a'' + x)^{m''}\} \therefore \frac{dy}{dx} = \frac{m}{a + x} + \frac{m'}{a' + x} + \frac{m''}{a'' + x}$.

10. $y = \log \frac{\sqrt{a} + \sqrt{x}}{\sqrt{a} - \sqrt{x}} \therefore \frac{dy}{dx} = \frac{\sqrt{a}}{(a - x)\sqrt{x}}$.

---

* This means the logarithm of the logarithm of x, but the notation we shall hereafter adopt will be $(\log)^2 x$, and which we shall extend to circular functions; thus, instead of sin (sin x), we shall write $(\sin)^2 x$, the *square* of the sine being written without the parenthesis, thus, $\sin^2 x$. We may call such expressions as $(\log)^n x$, $(\sin)^n x$, &c. the nth log of x, the nth sine of x, &c.

11. $y = e^{x^x} \therefore \frac{dy}{dx} = e^{x^x} \cdot x^x (1 + \log x)$.

12. $y = (\log)^n x \therefore \frac{dy}{dx} = \frac{1}{x \log x (\log)^2 x \ldots (\log)^{n-1} x}$.

13. $\log y = \frac{\sqrt{1+x^2}}{x} \therefore \frac{dy}{dx} = \frac{-y}{x^2\sqrt{1+x^2}} = \frac{-e^{\frac{\sqrt{1+x^2}}{x}}}{x^2\sqrt{1+x^2}}$.

14. $y = a^{b^z}$, z being a function of x, $\therefore \frac{dy}{dx} = \log a \log b \cdot a^{b^z} b^z \cdot \frac{dz}{dx}$.

15. $y = a^{b^{(x^2+x)}} \therefore \frac{dy}{dx} = \log a \cdot \log b \cdot a^{b^{(x^2+x)}} \cdot b^{x^2+x} (2x + 1)$.

16. $y = a^{\log x} \therefore \frac{dy}{dx} = \frac{\log a \cdot a^{\log x}}{x}$.

17. $y = e^{(\log)^n x} \therefore \frac{dy}{dx} = \frac{e^{(\log)^n x}}{x \log x (\log)^2 x \ldots (\log)^{n-1} x}$.

18. $y = x^{x^x} \therefore \frac{dy}{dx} = x^{x^x} \cdot x^x \{\frac{1}{x} + \log x (1 + \log x)\}$.

(14.) *To differentiate circular functions.*

In order to establish rules for the differentiation of circular functions, it will be necessary first to ascertain what ratio some two of the trigonometrical lines, as for instance sine and tangent, continually approach, as the arc to which they belong continually diminishes, and which ratio they actually attain only when the arc ultimately vanishes. For this purpose we may take the well known relation

$$\frac{\tan}{\sin} = \frac{\sec}{\text{rad}},$$

and, observing that as the arc diminishes the secant approaches nearer and nearer to the radius, with which it actually coincides when the arc vanishes, we may conclude that the value of $\frac{\tan}{\sin}$ continually ap-

proaches towards *unity* as the arc diminishes, and that unity must be the ratio with which the sine and tangent vanish simultaneously with the arc.* As, during the whole diminution of the sine and tangent,

---

* That 1 is the ultimate value of $\frac{\tan}{\sin}$ is evident; for, as in progressing towards this ultimate state, it approaches nearer and nearer to 1, from which it at length comes to differ by less than any assignable quantity, however small, the extreme value, or that which terminates this continuous series of results, can be no other than 1 itself.

It is no objection, therefore, to our conclusion to say that the geometrical relation $\frac{\tan}{\sin} = \frac{\sec}{\text{rad}}$ is not comprehensive of the extreme case; it is sufficient that it comprehends all but this, to enable us to infer the extreme boundary at which they terminate.

The student will not fail to observe that the ultimate or extreme state of the ratio here adverted to, is actually *attained* in the case before us. The terms of the ratio not only approach indefinitely near to evanescence, as the secant approaches the radius, but they actually vanish when the two latter lines coalesce. We direct attention to this circumstance because the terms "ultimate ratio" and "ultimately equal" appear to be used by Newton and most succeeding writers in a wider sense. Two quantities, for instance, are said to be ultimately in a ratio of equality when they approach continually towards equality; and, before the end of any finite time, come nearer to one another than by any given difference; although the quantities may be such as to never actually coalesce. Thus, the circle is the limit of all the circumscribed and inscribed polygons; and the three are therefore said to be ultimately equal, though they never actually coalesce. In such cases the limit, or ultimate state, is considered as never accurately reached, which is agreeably to the doctrine of the ancients; and the "ultimate equality" spoken of refers to the *fixed limits*, and not to the varying quantities themselves. When, however, the quantities compared really coalesce with their limits, then, of course, the ultimate equality, or the ultimate ratio, is that of the quantities themselves. Should these become evanescent, as in the case in the text, it may be said that the thing called the ultimate ratio can have no existence. Newton disposes of this objection as follows:

"Perhaps it may be objected that there is no ultimate proportion of evanescent quantities; because the proportion, before the quantities have

the arc itself continues of intermediate length, and as the former have, ultimately, a ratio of equality, the ultimate ratio of the intermediate arc to each must be that of equality; hence the ultimate ratios are as follow:

---

vanished, is not the ultimate, and when they are vanished, is none. But, by the same argument, it may be alleged, that a body arriving at a certain place and there stopping, has no ultimate velocity; because the velocity, before the body comes to the place, is not its ultimate velocity; when it has arrived, is none. But the answer is easy; for by the ultimate velocity is meant that with which the body is moved, neither before it arrives at its last place and the motion ceases, nor after, but at the very instant it arrives; that is, that velocity with which the body arrives at its last place, and with which the motion ceases. And, in like manner, by the ultimate ratio of evanescent quantities is to be understood the ratio of the quantities not before they vanish, nor afterwards, but with which they vanish. There is a limit which the velocity at the end of the motion may attain, but not exceed. This is the ultimate velocity. And there is the like limit in all quantities and proportions that begin and cease to be." See the Scholium at the end of the first section of the *Principia*.

With this quotation from Newton's celebrated Scholium we ought perhaps to terminate the present note; but having adverted above to the fixed limits, as distinct, in certain cases, from the varying quantities themselves, we feel inclined to ask whether in strictness such a distinction does not *always* exist? Are not the ultimate limits of varying geometrical quantities *always* things different in kind from the varying quantities themselves? And is there less impropriety in saying that a varying line ultimately *becomes* zero or 0, than in saying that a varying circumscribed polygon ultimately becomes, or merges into, the circle? If we admit that a line, continually decreasing, ultimately vanishes, we must admit that the side of a circumscribed polygon, continually decreasing, will ultimately vanish. As the diminution goes on, the entire polygon, of which this is a side, will furnish a series of points of contact, separated by intervals continually diminishing. But when the decreasing side actually vanishes, these intervals also vanish; and the hitherto interrupted series of points becomes, at the same instant, continuous; the polygon merges into the circle, and this circle is the last of the continuous series of magnitudes at which the varying polygon arrives. The limit, therefore, is actually reached, although not while the original figure continues a polygon.

$$\frac{\tan}{\sin} = 1, \quad \frac{\text{arc}}{\sin} = 1, \quad \frac{\text{arc}}{\tan} = 1,$$

and therefore,

$$\frac{\text{arc}}{\text{chord}} = 1, \quad \frac{\sin}{\text{chord}} = 1, \quad \frac{\tan}{\text{chord}} = 1.$$

1. Let it now be required to find the differential of $\sin x$. Changing x into $x + h$, we have (*Young's Trig.* p. 46, by putting $x + \frac{1}{2}h$ for A, and $\frac{1}{2}h$ for B, in form 4,)

$$\sin(x+h) = \sin x + 2\sin\tfrac{1}{2}h\cos(x+\tfrac{1}{2}h),$$

$$\therefore \frac{\sin(x+h) - \sin x}{h} = \frac{\sin\frac{1}{2}h}{\frac{1}{2}h}\cos(x+\tfrac{1}{2}h);$$

when $h = 0$, $\frac{\sin\frac{1}{2}h}{\frac{1}{2}h} = 1$, $\therefore \frac{\mathrm{d}\sin x}{\mathrm{d}x} = \cos x$ $\therefore \mathrm{d}\sin x = \cos x\,\mathrm{d}x$.

2. To differentiate $\cos x$,

$$\mathrm{d}\cos x = \mathrm{d}\sin(\tfrac{1}{2}\pi - x) = -\cos(\tfrac{1}{2}\pi - x)\,\mathrm{d}x = -\sin x\,\mathrm{d}x.$$

Cor. As $\mathrm{d}\cos = -\mathrm{d}\,\text{ver sin}$ $\therefore \mathrm{d}\,\text{ver sin}\, x = \sin x\,\mathrm{d}x$.

3. To differentiate $\tan x$,

$$\mathrm{d}\tan x = \mathrm{d}\frac{\sin x}{\cos x} = \frac{\cos x\,\mathrm{d}\sin x - \sin x\,\mathrm{d}\cos x}{\cos^2 x},$$

$$= \frac{\cos^2 x + \sin^2 x}{\cos^2 x}\,\mathrm{d}x = \frac{1}{\cos^2 x}\,\mathrm{d}x = \sec^2 x\,\mathrm{d}x.$$

4. To differentiate $\cot x$,

$$\mathrm{d}\cot x = \mathrm{d}\tan(\tfrac{1}{2}\pi - x) = -\sec^2(\tfrac{1}{2}\pi - x)\,\mathrm{d}x = -\operatorname{cosec}^2 x\,\mathrm{d}x.$$

5. To differentiate $\sec x$,

$$\mathrm{d}\sec x = \mathrm{d}\frac{1}{\cos x} = \frac{\sin x}{\cos^2 x}\,\mathrm{d}x = \tan x\sec x\,\mathrm{d}x.$$

6. To differentiate $\operatorname{cosec} x$,

$$\mathrm{d}\,\mathrm{cosec}\,x = \mathrm{d}\frac{1}{\sin x} = -\frac{\cos x}{\sin^2 x}\,\mathrm{d}x = -\cot x\,\mathrm{cosec}\,x\,\mathrm{d}x.$$

These six forms the student should endeavour to preserve in his memory.

EXAMPLES.

1. $y = \sin^2 x \;\therefore\; \mathrm{d}y = 2\sin x\,\mathrm{d}\sin x = 2\sin x\cos x\,\mathrm{d}x = \sin 2x\,\mathrm{d}x,$

$$\therefore \frac{\mathrm{d}y}{\mathrm{d}x} = \sin 2x.$$

2. $y = \sin^n x \;\therefore\; \mathrm{d}y = n\sin^{n-1} x\,\mathrm{d}\sin x = n\sin^{n-1} x\cos x\mathrm{d}x,$

$$\therefore \frac{\mathrm{d}y}{\mathrm{d}x} = n\sin^{n-1} x\cos x.$$

3. $y = \cos mx \;\therefore\; \mathrm{d}y = -\sin mx\mathrm{d}mx = -m\sin mx\mathrm{d}x,$

$$\therefore \frac{\mathrm{d}y}{\mathrm{d}x} = -m\sin mx.$$

4. $u = y\tan x^n$, y being a function of x,

$$\therefore \mathrm{d}u = \tan x^n\,\mathrm{d}y + y\,\mathrm{d}\tan x^n,$$

now $\mathrm{d}\tan x^n = \sec^2 x^n\,\mathrm{d}.x^n = nx^{n-1}\sec^2 x^n\,\mathrm{d}x,$

$$\therefore \frac{\mathrm{d}u}{\mathrm{d}x} = \tan x^n\frac{\mathrm{d}y}{\mathrm{d}x} + ynx^{n-1}\sec^2 x^n.$$

5. $u = \cot x^y \;\therefore\; \mathrm{d}u = -\mathrm{cosec}^2 x^y\,\mathrm{d}x^y.$

Put $z = x^y \;\therefore\; \log z = y\log x,$

$$\therefore \frac{\mathrm{d}z}{z} = y\frac{\mathrm{d}x}{x} + \log x\mathrm{d}y \;\therefore\; \mathrm{d}z = \mathrm{d}x^y = (y\frac{\mathrm{d}x}{x} + \log x\,\mathrm{d}y)x^y,$$

$$\therefore \frac{\mathrm{d}u}{\mathrm{d}x} = -\mathrm{cosec}^2 x^y\;(\frac{y}{x} + \log x\frac{\mathrm{d}y}{\mathrm{d}x})\,x^y.$$

6. $y = xe^{\cos x}$ $\therefore$ $dy = e^{\cos x}\,dx + xe^{\cos x}\,d\cos x = e^{\cos x}(1 - x\sin x)\,dx$,

$$\therefore \frac{dy}{dx} = e^{\cos x}(1 - x\sin x).$$

7. $y = \log(xe^{\cos x})$ $\therefore$ $dy = \frac{d(xe^{\cos x})}{xe^{\cos x}}$, and $d(xe^{\cos x}) = e^{\cos x}(1 - x\sin x)\,dx$,

$$\therefore \frac{dy}{dx} = \frac{1 - x\sin x}{x}.$$

8. $y = \cos x + \sqrt{-1}\sin x$, $\therefore$ $\frac{dy}{dx} = -\sin x + \sqrt{-1}\cos x$.

9. $y = \cos x + \cos 2x + \cos 3x + \&c.$

$$\therefore \frac{dy}{dx} = -(\sin x + 2\sin 2x + 3\sin 3x + \&c.)$$

10. $y = xe^{\tan x}$ $\therefore$ $\frac{dy}{dx} = \{1 + x\sec^2 x\}e^{\tan x}$.

11. $y = \frac{\sin^m x}{\cos^n x}$ $\therefore$ $\frac{dy}{dx} = m\frac{\sin^{m-1}x}{\cos^{n-1}x} + n\frac{\sin^{m+1}x}{\cos^{n+1}x}$.

12. $u = \sec x^y$ $\therefore$ $\frac{du}{dx} = x^y \tan x^y \sec x^y \left\{\frac{y}{x} + \log x\frac{dy}{dx}\right\}$.

(15.) In the preceding trigonometrical expressions, the arc is considered as the independent variable, and the lines sine, cosine, &c. as functions of it; we shall now consider the *inverse functions* as they are called, that is, those in which the arc is considered as a function of the sine, the cosine, &c. A particular notation has been proposed for inverse functions: thus, if $y = F(x)$ be the direct function, then $x = F^{-1}(y)$ is the inverse function, that is, if we represent the function that y is of x by $y = F(x)$, the function that x is of y will be denoted by $x = F^{-1}(y)$. By thus representing these inverse functions, we may return immediately to the direct functions, considering, for the moment, F^{-1} in the light of a negative power of F, or an equivalent to $\frac{1}{F}$; for then $x = F^{-1}(y)$ immediately leads to $y = F(x)$. Thus, if $x = \log^{-1} y$ $\therefore$ $y = \log x$, the

inverse function $\log^{-1} y$ meaning *the number whose log is* y. In like manner, $y = \sin^{-1} x$ means that y is the arc whose sine is x; that is, returning to the direct function, $\sin y = x$.

1. To differentiate $y = \sin^{-1} x$.

Here the direct function is $\sin y = x \therefore \mathrm{d} \sin y = \mathrm{d}x$, that is,

$$\cos y \mathrm{d}y = \mathrm{d}x \therefore \frac{\mathrm{d}y}{\mathrm{d}x} = \frac{1}{\cos y} = \frac{1}{\sqrt{1 - \sin^2 y}} = \frac{1}{\sqrt{1 - x^2}}.$$

2. To differentiate $y = \cos^{-1} x$.

$$\cos y = x \therefore -\sin y \mathrm{d}y = \mathrm{d}x,$$

$$\therefore \frac{\mathrm{d}y}{\mathrm{d}x} = -\frac{1}{\sin y} = -\frac{1}{\sqrt{1 - \cos^2 y}} = \frac{-1}{\sqrt{1 - x^2}}.$$

3. To differentiate $y = \mathrm{versin}^{-1} x$.

$$\mathrm{versin}\, y = x \therefore \sin y \mathrm{d}y = \mathrm{d}x,$$

$$\therefore \frac{\mathrm{d}y}{\mathrm{d}x} = \frac{1}{\sin y} = \frac{1}{\sqrt{2x - x^2}}.$$

4. To differentiate $y = \tan^{-1} x$.

$$\tan y = x \therefore \sec^2 y \mathrm{d}y = \mathrm{d}x,$$

$$\therefore \frac{\mathrm{d}y}{\mathrm{d}x} = \frac{1}{\sec^2 y} = \frac{1}{1 + x^2}.$$

5. To differentiate $y = \cot^{-1} x$.

$$\cot y = x \therefore -\mathrm{cosec}^2 y \mathrm{d}y = \mathrm{d}x,$$

$$\therefore \frac{\mathrm{d}y}{\mathrm{d}x} = -\frac{1}{\mathrm{cosec}^2 y} = \frac{-1}{1 + x^2}.$$

6. To differentiate $y = \sec^{-1} x$.

$$\sec y = x, \therefore \tan y \sec y \mathrm{d}y = \mathrm{d}x,$$

$$\therefore \frac{\mathrm{d}y}{\mathrm{d}x} = \frac{1}{\tan y \sec y} = \frac{1}{x\sqrt{x^2 - 1}}.$$

7. To differentiate $y = \operatorname{cosec}^{-1} x$.

$$\operatorname{cosec} y = x \therefore -\cot y \operatorname{cosec} y dy = dx,$$

$$\therefore \frac{dy}{dx} = \frac{-1}{\cot y \operatorname{cosec} y} = -\frac{1}{x\sqrt{x^2-1}}.$$

EXAMPLES.

1. $y = \sin^{-1} mx \therefore \frac{dy}{dmx} = \frac{1}{\sqrt{1-m^2x^2}}$,

$$\therefore \frac{dy}{dx} = \frac{m}{\sqrt{1-m^2x^2}}.$$

2. $y = x \sin^{-1} x^2 \therefore \frac{dy}{dx} = \sin^{-1} x^2 + x \frac{d \sin^{-1} x^2}{dx}$,

and

$$\frac{d \sin^{-1} x^2}{dx} = \frac{2x}{\sqrt{1-x^4}}.$$

$$\therefore \frac{dy}{dx} = \sin^{-1} x^2 + \frac{2x^2}{\sqrt{1-x^4}}.$$

3. $y = \cos^{-1} x\sqrt{1-x^2}$. Put $x\sqrt{1-x^2} = z \therefore dy = -\frac{dz}{\sqrt{1-z^2}}$,

but $dz = (\sqrt{1-x^2} - \frac{x^2}{\sqrt{1-x^2}})\,dx$, and $\sqrt{1-z^2} = \sqrt{1-x^2+x^4}$,

$$\therefore \frac{dy}{dx} = \frac{2x^2-1}{\sqrt{(1-x^2+x^4)(1-x^2)}}.$$

4. $y = \tan^{-1} \frac{x}{2}$.

$$\frac{dy}{\frac{1}{2}dx} = \frac{1}{1+(\frac{x}{2})^2} \therefore \frac{dy}{dx} = \frac{8}{4+x^2}.$$

5. $y = \cot^{-1}(a+mx)^2 \therefore dy = -\frac{1}{1+(a+mx)^4} d(a+mx)^2$,

and $d(a+mx)^2 = 2(a+mx)\,m dx$,

$$\therefore \frac{dy}{dx} = \frac{2m\,(a+mx)}{1+(a+mx)^4}.$$

6. $y = \sec^{-1}\frac{a}{x^m}$ $\therefore dy = \dfrac{x^m}{a\sqrt{(\frac{a}{x^m})^2-1}}\, d\,\frac{a}{x^m}$,

and

$$d\,\frac{a}{x^m} = -\frac{am\,dx}{x^{m+1}}.$$

$$\therefore \frac{dy}{dx} = -\frac{mx^{m-1}}{\sqrt{a^2-x^{2m}}}.$$

7. $y = \operatorname{cosec}^{-1}\dfrac{\sqrt{1+x^2}}{x}$,

$$\therefore dy = -d\,\frac{\sqrt{1+x^2}}{x} \div \frac{\sqrt{1+x^2}}{x}\,(\frac{1+x^2}{x^2}-1)^{\frac{1}{2}},$$

$$= -d\,\frac{\sqrt{1+x^2}}{x} \div \frac{\sqrt{1+x^2}}{x^2};$$

but $d\,\dfrac{\sqrt{1+x^2}}{x} = -\dfrac{1}{x^2\sqrt{1+x^2}}\,dx$,

$$\therefore \frac{dy}{dx} = \frac{1}{1+x^2}.$$

8. $y = (\sin^{-1}x)^2$ $\therefore \dfrac{dy}{dx} = 2\sin^{-1}x\dfrac{1}{\sqrt{1-x^2}}$.

9. $y = \cos^{-1}\dfrac{1}{\sqrt{1+x^2}}$ $\therefore \dfrac{dy}{dx} = \dfrac{1}{1+x^2}$.

10. $y = \tan^{-1}\sqrt{\dfrac{1-x}{1+x}}$ $\therefore \dfrac{dy}{dx} = -\dfrac{1}{2\sqrt{1-x^2}}$.

11. $y = (\cot^{-1}x)^2$ $\therefore \dfrac{dy}{dx} = -\dfrac{2}{1+x^2}\cot^{-1}x$.

12. $y = \sec^{-1} x^n \therefore \frac{dy}{dx} = \frac{n}{x\sqrt{x^{2n} - 1}}$.

13. $y = \operatorname{cosec}^{-1} mx^2 \therefore \frac{dy}{dx} = -\frac{2}{x\sqrt{m^2 x^4 - 1}}$.

14. $y = \operatorname{versin}^{-1} e^x \therefore \frac{dy}{dx} = \frac{e^{\frac{x}{2}}}{\sqrt{2 - e^x}}$.

(16.) In the preceding expressions the radius of the arc is always represented by unity, but, as the differentials are frequently required to radius r, we shall terminate this chapter with the several formulas in (15) accommodated to this radius. We must observe, that as y and x are homogeneous in each of those forms, $\frac{y}{x}$, and therefore $\frac{y' - y}{h}$, is always a number; so that this ratio in the limit, that is $\frac{dy}{dx}$, is a number. Hence, r must be introduced so as to render the numerator and denominator of each expression of the same dimensions. The formulas, therefore, become

$$\text{d} \sin^{-1} x = \frac{r\text{d}x}{\sqrt{r^2 - x^2}}.$$

$$\text{d} \cos^{-1} x = -\frac{r\text{d}x}{\sqrt{r^2 - x^2}}.$$

$$\text{d} \operatorname{versin}^{-1} x = \frac{r\text{d}x}{\sqrt{2rx - x^2}}.$$

$$\text{d} \tan^{-1} x = \frac{r^2\,\text{d}x}{r^2 + x^2}.$$

$$\text{d} \cot^{-1} x = -\frac{r^2\,\text{d}x}{r^2 + x^2}.$$

$$\text{d} \sec^{-1} x = \frac{r^2\,\text{d}x}{x\sqrt{x^2 - r^2}}.$$

$$\text{d} \operatorname{cosec}^{-1} x = -\frac{r^2\,\text{d}x}{x\sqrt{x^2 - r^2}}.$$

On successive Differentiation.

(17.) Since the differential coefficient derived from any function of a variable may contain that variable, this coefficient itself may be differentiated, and we thus derive a *second differential coefficient*. In like manner, by differentiating this second coefficient, if the variable still enters it, we obtain a *third differential coefficient*, and in this way we may continue the successive differentiation till we arrive at a coefficient without the variable, when the process must terminate.

Thus, taking the function $y = ax^4$, we have, for the first differential coefficient, $\frac{dy}{dx} = 4ax^3$, as this coefficient contains x, we have, by differentiating it, the second differential coefficient $= 12ax^2$; continuing the process, we have $24ax$ for the third differential coefficient, and $24a$ for the fourth, which being constant its differential coefficient is 0.

If we were to express these several coefficients agreeably to the notation hitherto adopted, they would be

$$\text{first diff. coef. } \frac{dy}{dx} = 4ax^3,$$

$$\text{second diff. coef. } \frac{d\frac{dy}{dx}}{dx} = 12\,ax^2,$$

$$\text{third diff. coef. } \frac{d\frac{d\frac{dy}{dx}}{dx}}{dx} = 24\,ax,$$

&c.

But this mode of expressing the successive coefficients is obviously very inconvenient, and they are accordingly written in the following more commodious manner:

$$\text{first diff. coef. } \frac{dy}{dx},$$

$$\text{second diff. coef. } \frac{d^2y}{dx^2},$$

$$\text{third diff. coef. } \frac{d^3y}{dx^3},$$

$$n\text{th diff. coef. } \frac{d^ny}{dx^n}.$$

It must be observed here, and carefully borne in mind by the student, that this notation is nothing more than a brief and compact mode of representing, symbolically, the results of certain *operations* performed upon a function, y, of x. The operation implied in $\frac{dy}{dx}$ has been already explained; a like operation, performed on the result, is expressed by $\frac{d^2y}{dx^2}$, a third such operation by $\frac{d^3y}{dx^3}$, and so on; hence d^2, d^3, &c. are not powers, but symbols, standing in place of the words *second differential of*, *third differential of*, &c. There is no impropriety in calling the expressions dx^2, dx^3, &c. powers, not, however, of x, but of dx: to distinguish the differential of a power from the power of a differential, a dot is placed in the former case between d and the power.

(18.) The following are a few illustrations of the process of successive differentiation:

1. $y = x^m$.

$$\therefore \frac{dy}{dx} = mx^{m-1},$$

$$\frac{d^2y}{dx^2} = m(m-1)x^{m-2},$$

$$\frac{d^3y}{dx^3} = m(m-1)(m-2)x^{m-3},$$

$$\frac{d^4y}{dx^4} = m(m-1)(m-2)(m-3)x^{m-4},$$

&c. &c.

2. $u = yz$, both y and z being functions of x,

$$\therefore \frac{du}{dx} = y\frac{dz}{dx} + z\frac{dy}{dx}$$

$$\frac{d^2u}{dx^2} = y\frac{d^2z}{dx^2} + \frac{dydz}{dx^2} + \frac{zd^2y}{dx^2} + \frac{dzdy}{dx^2}$$

$$= y\frac{d^2z}{dx^2} + 2\frac{dydz}{dx^2} + \frac{zd^2y}{dx^2}$$

$$\frac{d^3u}{dx^3} = y\frac{d^3z}{dx^3} + 3\frac{dyd^2z}{dx^3} + 3\frac{dzd^2y}{dx^3} + \frac{zd^3y}{dx^3}$$

&c. &c.

3. $y = \log x$.

$$\therefore \frac{dy}{dx} = \frac{1}{x} \qquad \frac{d^4y}{dx^4} = -\frac{2\cdot 3}{x^4}$$

$$\frac{d^2y}{dx^2} = -\frac{1}{x^2} \qquad \frac{d^5y}{dx^5} = \frac{2\cdot 3\cdot 4}{x^5}$$

$$\frac{d^3y}{dx^3} = \frac{2}{x^3} \qquad \frac{d^6y}{dx^6} = -\frac{2\cdot 3\cdot 4\cdot 5}{x^6}$$

&c.

4. $y = e^x$.

$$\therefore \frac{dy}{dx} = e^x \qquad \frac{d^3y}{dx^3} = e^x$$

$$\frac{d^2y}{dx^2} = e^x \qquad \frac{d^4y}{dx^4} = e^x$$

&c.

If instead of e the base were a, the several coefficients would be

$$\log a \cdot a^x, \quad \log^2 a \cdot a^x, \quad \log^3 a \cdot a^x, \quad \log^4 a \cdot a^x, \quad \&c.$$

It appears, therefore, that exponential functions possess this property, viz. that $\frac{d^n y}{dx^n} \div y$ is always constant.

5. $y = \sin x$.

$$\therefore \frac{dy}{dx} = \cos x \qquad \frac{d^3y}{dx^3} = -\cos x$$

$$\frac{d^2y}{dx^2} = -\sin x \qquad \frac{d^4y}{dx^4} = \sin x$$

&c.

We need not multiply examples here, as the process of successive differentiation will be very frequently employed in the next two chapters.

CHAPTER III.

ON MACLAURIN'S THEOREM.

(19.) If y represent a function of x, which it is possible to develop in a series of positive ascending powers of that variable, then will that development be

$$y = [y] + [\frac{dy}{dx}]\, x + \frac{1}{2}\, [\frac{d^2y}{dx^2}]\, x^2 + \frac{1}{2 \cdot 3}\, [\frac{d^3y}{dx^3}]\, x^3 + \&c.$$

where the brackets are intended to intimate that the functions which they enclose are to be taken in that particular state, arising from putting $x = 0$.*

For, since by hypothesis,

---

* This plan of enclosing the differential coefficient in brackets we shall usually adopt, when we wish to express not the *general* state of the function, but that state which arises from the variable taking a *particular* value. What that value is will generally be made known by the nature of the enquiry.

$$y = A + Bx + Cx^2 + Dx^3 + Ex^4 + \&c. \ldots . \quad (1),$$

$$\therefore \frac{dy}{dx} = B + 2Cx + 3Dx^2 + 4Ex^3 + \&c.$$

$$\frac{d^2y}{dx^2} = 2C + 2 \cdot 3Dx + 3 \cdot 4Ex^2 + \&c.$$

$$\frac{d^3y}{dx^3} = 2 \cdot 3D + 2 \cdot 3 \cdot 4Ex + \&c.$$

&c. &c.

Let, now, $x = 0$, then

$$[y] = A$$

$$[\frac{dy}{dx}] = B$$

$$[\frac{d^2y}{dx^2}] = 2C \therefore C = \frac{1}{2} [\frac{d^2y}{dx^2}]$$

$$[\frac{d^3y}{dx^3}] = 2 \cdot 3D \therefore D = \frac{1}{2 \cdot 3} [\frac{d^3y}{dx^3}]$$

&c. &c.

Hence, by substitution, equation (1) becomes

$$y = [y] + [\frac{dy}{dx}] x + \frac{1}{2} [\frac{d^2y}{dx^2}] x^2 + \frac{1}{2 \cdot 3} [\frac{d^3y}{dx^3}] x^3 + \&c. \ldots . \quad (2),$$

which is Maclaurin's theorem for the development of a function, according to the ascending powers of the variable.

From this important theorem a rule may be deduced, by the aid of which we may obtain the development of a function very readily, from merely knowing the development of the first differential coefficient. Thus, by the theorem just established, the development of the function $\frac{dy}{dx}$ is

$$\frac{dy}{dx} = [\frac{dy}{dx}] + [\frac{d^2y}{dx^2}] x + \frac{1}{2} [\frac{d^3y}{dx^3}] x^2 + \frac{1}{2 \cdot 3} [\frac{d^4y}{dx^4}] x^3 + \&c.$$

Comparing this with the development of the function y above, we see that y must be equal to $[y]$ plus the several terms in the development of $\frac{dy}{dx}$, multiplied respectively by x, $\frac{x}{2}$, $\frac{x}{3}$, &c. Hence we have this rule, viz.

Develop $\frac{dy}{dx}$ in a series of ascending powers of x, increase the exponent of x in each term by unity, and divide the term by the exponent thus increased.

In many cases where the successive differentiation would be laborious, the development of $\frac{dy}{dx}$ may be easily obtained by common algebra, and thus the labour adverted to entirely avoided. An example of this may be seen in the development of $\tan^{-1} x$ at page 50.*

EXAMPLES.

(20.) 1. Let it be required to develop $(a+x)^n$, the exponent n being any number whatever, either positive or negative, whole or fractional, rational or irrational.

Put $y = (a+x)^n$ . . therefore . $[y] = a^n$

$$\therefore \frac{dy}{dx} = n(a+x)^{n-1} \quad . \quad . \quad . \quad . \quad . \quad . \quad [\frac{dy}{dx}] = na^{n-1}$$

$$\frac{d^2y}{dx^2} = n(n-1)(a+x)^{n-2} \quad . \quad . \quad . \quad [\frac{d^2y}{dx^2}] = n(n-1)a^{n-2}$$

$$\frac{d^3y}{dx^3} = n(n-1)(n-2)(a+x)^{n-3}, \quad [\frac{d^3y}{dx^3}] =$$

$$n(n-1)(n-2)a^{n-3}$$

&c. &c.

Substituting these values for the coefficients in the foregoing theorem, there results

---

* The preceding rule was first given, incidentally, by the author of this work in his "Elements of the Integral Calculus."

$$(a+x)^n = a^n + na^{n-1}x + \frac{n(n-1)}{2}a^{n-2}x^2 + \frac{n(n-1)(n-2)}{2\cdot 3}a^{n-3}x^3 + \&c.$$

and thus the truth of the *Binomial Theorem* is established in its utmost generality.

2. To develop $\log(a+x)$.*

Put $y = \log(a+x)$, therefore $[y] = \log a$

$$\frac{dy}{dx} = \frac{1}{a+x} \quad \ldots \quad [\frac{dy}{dx}] = \frac{1}{a}$$

$$\frac{d^2y}{dx^2} = -\frac{1}{(a+x)^2} \quad \ldots \quad [\frac{d^2y}{dx^2}] = -\frac{1}{a^2}$$

$$\frac{d^3y}{dx^3} = \frac{2}{(a+x)^3} \quad \ldots \quad [\frac{d^3y}{dx^3}] = \frac{2}{a^3}$$

$$\frac{d^4y}{dx^4} = -\frac{2\cdot 3}{(a+x)^4} \quad \ldots \quad [\frac{d^4y}{dx^4}] = -\frac{2\cdot 3}{a^4}$$

&c. &c.

$$\therefore \log(a+x) = \log a + \frac{x}{a} - \frac{x^2}{2a^2} + \frac{x^3}{3a^3} - \frac{x^4}{4a^4} + \&c.$$

If $y = \log x$ were proposed, then, since $[y]$, $[\frac{dy}{dx}]$, &c. are infinite, we infer, for reasons similar to those assigned at page 8, that the development in the proposed form, with finite coefficients, is impossible.

The rule which we have given at the close of last article may be applied with some advantage to the present example. For, by common division, we readily get

$$\frac{dy}{dx} = \frac{1}{a+x} = \frac{1}{a} - \frac{x}{a^2} + \frac{x^2}{a^3} - \frac{x^3}{a^4} + \&c.$$

* This example must be regarded merely as an exercise in successive differentiation. The development of $\log(a+x)$ has already been supposed known from common algebra, (see p. 18.)

Hence, by the rule,

$$y = \log a + \frac{x}{a} - \frac{x^2}{2a^2} + \frac{x^3}{3a^3} - \frac{x^4}{4a^4} + \&c.$$

3. To develop $\sin x$.

$$y = \sin x \quad . \quad . \quad . \quad . \quad [y] = 0$$

$$\frac{dy}{dx} = \cos x \quad . \quad . \quad . \quad . \quad [\frac{dy}{dx}] = 1$$

$$\frac{d^2y}{dx^2} = -\sin x \quad . \quad . \quad . \quad [\frac{d^2y}{dx^2}] = 0$$

$$\frac{d^3y}{dx^3} = -\cos x \quad . \quad . \quad . \quad [\frac{d^3y}{dx^3}] = -1$$

$$\frac{d^4y}{dx^4} = \sin x \quad . \quad . \quad . \quad . \quad [\frac{d^4y}{dx^4}] = 0$$

$$\frac{d^5y}{dx^5} = \cos x \quad . \quad . \quad . \quad . \quad [\frac{d^5y}{dx^5}] = 1$$

&c. &c.

$$\therefore \sin x = x - \frac{x^3}{1 \cdot 2 \cdot 3} + \frac{x^5}{1 \cdot 2 \cdot 3 \cdot 4 \cdot 5} - \&c.$$

4. To develop $\cos x$.

$$y = \cos x \quad . \quad . \quad . \quad . \quad [y] = 1$$

$$\frac{dy}{dx} = -\sin x \quad . \quad . \quad . \quad [\frac{dy}{dx}] = 0$$

$$\frac{d^2y}{dx^2} = -\cos x \quad . \quad . \quad . \quad [\frac{d^2y}{dx^2}] = -1$$

&c. &c.

$$\therefore \cos x = 1 - \frac{x^2}{1 \cdot 2} + \frac{x^4}{1 \cdot 2 \cdot 3 \cdot 4} - \&c.$$

5. To develop a^x.

$$y = a^x \quad \text{therefore} \quad [y] = 1$$

$$\frac{dy}{dx} = {}^{*}Aa^x \quad . \quad . \quad . \quad . \quad [\frac{dy}{dx}] = A$$

$$\frac{d^2y}{dx^2} = A^2a^x \quad . \quad . \quad . \quad . \quad [\frac{d^2y}{dx^2}] = A^2$$

$$\frac{d^3y}{dx^3} = A^3a^x \quad . \quad . \quad . \quad . \quad [\frac{d^3y}{dx^3}] = A^3$$

&c. &c.

$$\therefore a^x = 1 + Ax + \frac{A^2x^2}{1 \cdot 2} + \frac{A^3x^3}{1 \cdot 2 \cdot 3} + \&c.$$

which is the *Exponential Theorem.*

Since $A = \log a$, we may give to the development the form

$$a^x = 1 + x \log a + \frac{1}{2}(x \log a)^2 + \frac{1}{2 \cdot 3}(x \log a)^3 + \&c.$$

For $x = 1$, we have the following expression for any number, a, in terms of its Napierian logarithm:

$$a = 1 + \log a + \frac{1}{2}\log^2 a + \frac{1}{2 \cdot 3}\log^3 a + \&c.$$

changing a into the Napierian base, e, we have

$$e^x = 1 + x + \frac{x^2}{2} + \frac{x^3}{2 \cdot 3} + \&c.$$

which, when $x = 1$, gives, for the base e, the value

$$e = 1 + 1 + \frac{1}{2} + \frac{1}{2 \cdot 3} + \&c.$$

---

* A is put here for the hyperbolic logarithm of the base a, that is, for the expression

$$(a-1) - \tfrac{1}{2}(a-1)^2 + \tfrac{1}{3}(a-1)^3 - \&c. \quad \text{(See p. 18.)}$$

(21.) From the development of e^x may be immediately derived several very curious and useful analytical formulas, and we shall avail ourselves of this opportunity to present the principal of them to the notice of the student.

In the development of e^x put $-x$ for x, and we shall have

$$e^{-x} = 1 - x + \frac{x^2}{2} - \frac{x^3}{2 \cdot 3} + \&c.$$

Add this to the development of e^x, and there results

$$\frac{e^x + e^{-x}}{2} = 1 + \frac{x^2}{2} + \frac{x^4}{2 \cdot 3 \cdot 4} + \frac{x^6}{2 \cdot 3 \cdot 4 \cdot 5 \cdot 6} + \&c.$$

Subtract e^{-x} from e^x, and we have

$$\frac{e^x - e^{-x}}{2} = x + \frac{x^3}{2 \cdot 3} + \frac{x^5}{2 \cdot 3 \cdot 4 \cdot 5} + \frac{x^7}{2 \cdot 3 \cdot 4 \cdot 5 \cdot 6 \cdot 7} + \&c.$$

In each of these developments substitute $z\sqrt{-1}$ for x, and there will result the following remarkable expressions, viz.

$$\frac{e^{z\sqrt{-1}} + e^{-z\sqrt{-1}}}{2} = 1 - \frac{z^2}{1 \cdot 2} + \frac{z^4}{1 \cdot 2 \cdot 3 \cdot 4} - \&c.$$

$$\frac{e^{z\sqrt{-1}} - e^{-z\sqrt{-1}}}{2\sqrt{-1}} = z - \frac{z^3}{1 \cdot 2 \cdot 3} + \frac{z^5}{1 \cdot 2 \cdot 3 \cdot 4 \cdot 5} - \&c.$$

Now it has been seen (examples 4 and 3) that these two series are also the respective developments of $\cos z$ and $\sin z$; hence, putting x instead of z, we may conclude that

$$\sin x = \frac{e^{x\sqrt{-1}} - e^{-x\sqrt{-1}}}{2\sqrt{-1}} \quad \ldots\ldots (1)$$

$$\cos x = \frac{e^{x\sqrt{-1}} + e^{-x\sqrt{-1}}}{2} \quad \ldots\ldots (2),$$

where the sine and cosine of a *real* arc are expressed by imaginary exponentials.

These expressions were first deduced by Euler, and are considered by Lagrange as among the finest analytical discoveries of the age. (*Calcul des Fonctions*, page 114.)

(22.) If for the real arc, x, we substitute the imaginary arc, $x\sqrt{-1}$, we shall have

$$\sin(x\sqrt{-1}) = \frac{e^{-x} - e^{x}}{2\sqrt{-1}} \quad \ldots\ (3)$$

$$\cos(x\sqrt{-1}) = \frac{e^{-x} + e^{x}}{2} \quad \ldots\ (4),$$

Also, since $\frac{\sin}{\cos} = \tan$, it follows, from (1) and (2), that

$$\sqrt{-1}\tan x = \frac{e^{x\sqrt{-1}} - e^{-x\sqrt{-1}}}{e^{x\sqrt{-1}} + e^{-x\sqrt{-1}}} = \frac{e^{2x\sqrt{-1}} - 1}{e^{2x\sqrt{-1}} + 1}.$$

By multiplying equation (1) by $\sqrt{-1}$, and adding the result to (2), we have

$$\cos x \pm \sqrt{-1}\sin x = e^{\pm x\sqrt{-1}} \quad \ldots\ (5);$$

or, if we change x into mx,

$$\cos mx \pm \sqrt{-1}\sin mx = e^{\pm mx\sqrt{-1}} \quad \ldots\ (6),$$

but $e^{\pm mx\sqrt{-1}}$ is $e^{\pm x\sqrt{-1}}$ raised to the mth power. Hence this singular property, viz.

$$(\cos x \pm \sqrt{-1}\sin x)^m = \cos mx \pm \sqrt{-1}\sin mx \quad \ldots\ (7),$$

which was discovered by De Moivre, and is hence called *De Moivre's formula*.

If the first side of this equation be developed by the binomial theorem, it becomes

$$\cos^m x \pm m \cos^{m-1} xp + \frac{m(m-1)}{2} \cos^{m-2} xp^2 \pm \&c.$$

p being put for the imaginary $\sqrt{-1} \sin x$.

Now in any equation, the imaginaries on one side are equal to those on the other, (*Alg.* p. 147); hence, expunging from this expression all the imaginaries, that is, all the terms containing an odd power of p, we have, in virtue of equation (7),

$$\cos mx = \cos^m x - \frac{m(m-1)}{2} \cos^{m-2} x \sin^2 x +$$

$$\frac{m(m-1)(m-2)(m-3)}{2 \cdot 3 \cdot 4} \cos^{m-4} x \sin^4 x - \&c.$$

In like manner, equating $\sqrt{-1} \sin mx$ with the imaginary part of the above development, and then dividing by $\sqrt{-1}$, we have

$$\sin mx = m \cos^{m-1} x \sin x - \frac{m(m-1)(m-2)}{2 \cdot 3} \cos^{m-3} x \sin^3 x + \&c.$$

From these two series the sine and cosine of a multiple arc may be determined from the sine and cosine of the arc itself.

(23.) If in the formula (2) we represent $e^{x\sqrt{-1}}$ by y, then $e^{-x\sqrt{-1}} = \frac{1}{y}$; therefore,

$$2 \cos x = y + \frac{1}{y},$$

or, if in the same formula mx be put x, we have

$$2 \cos mx = y^m + \frac{1}{y^m},$$

and from these two equations we deduce the following, viz.

$$y^2 - 2y \cos x + 1 = 0 \ldots\ldots (1)$$

$$y^{2m} - 2y^m \cos mx + 1 = 0 \ldots\ldots (2).$$

Since these equations exist simultaneously, the latter must have two of its roots or values of y equal to the two roots of the former, and therefore the first member of (2) must be divisible by the first member of (1), or, putting θ for mx, we have

$$y^{2m} - 2y^m \cos\theta + 1 \ldots\ldots (3),$$

divisible by

$$y^2 - 2y \cos\frac{\theta}{m} + 1 \ldots\ldots (4).$$

But $\cos\theta = \cos(\theta + 2n\pi)$, n being any whole number, and $\pi = 180°$; hence, making successively $n = 0$, $= 1$, $= 2$, &c. to $n = m - 1$, we have, since the first equation continues to be divisible by the second in all these cases,

$$y^{2m} - 2y^m \cos\theta + 1 = \left(y^2 - 2y \cos\frac{\theta}{m} + 1\right)$$

$$\times \left(y^2 - 2y \cos\frac{\theta + 2\pi}{m} + 1\right)$$

$$\times \left(y^2 - 2y \cos\frac{\theta + 4\pi}{m} + 1\right)$$

$$\times \left(y^2 - 2y \cos\frac{\theta + 6\pi}{m} + 1\right) \text{ &c. to } m \text{ factors.}$$

The truth of this equation is obvious, for, while the substitution of $\theta + 2n\pi$ for θ causes no alteration in the expression (3), the same substitution in (4) gives to that expression a new value, for every value of n, from $n = 0$ to $n = m - 1$; for the arcs $\frac{\theta}{m}$, $\frac{\theta + 2\pi}{m}$, &c. are all different. As, therefore, the expression (3) is divisible by (4) under all these m changes of value, it is plain that these are its m quadratic factors.

In this way may any trinomial of the form $y^{2m} - 2ky^m + 1$ be decomposed into its quadratic factors, provided k does not exceed unity, for then k may always be replaced by the cosine of an arc.

(24.) The geometrical interpretation of the foregoing equation, presents a curious property of the circle, first discovered by De Moivre. To exhibit this property, let P be any point either within or without the circle whose centre is O, and let the circumference be divided into any number, m, of equal parts, commencing at any point A. Join the points of division, A, B, C, &c. to P, then, since in the foregoing analytical expression the radius OA is expressed by unity, we shall have, by introducing the radius itself, so as to render the terms homogeneous, the following geometrical values of the above factors, where it is to be observed that





$$\angle \mathrm{POA} = \frac{\theta}{m} \text{ and } \mathrm{OP} = y,$$

$$y^{2m} - 2y^m \cos\theta + 1 = \mathrm{OP}^{2m} - 2\mathrm{OP}^m \times \mathrm{OA}^m \cos m\,(\mathrm{AOP}) + \mathrm{AO}^{2m}$$

$$y^2 - 2y \cos\frac{\theta}{m} + 1 = \mathrm{OP}^2 - 2\mathrm{OP} \times \mathrm{OA} \cos \mathrm{AOP} + \mathrm{AO}^2 = \mathrm{PA}^2$$

$$y^2 - 2y \cos\frac{\theta + 2\pi}{m} + 1 = \mathrm{OP}^2 - 2\mathrm{OP} \times \mathrm{OA} \cos \mathrm{BOP} + \mathrm{BO}^2 = \mathrm{PB}^2$$

$$y^2 - 2y \cos\frac{\theta + 4\pi}{m} + 1 = \mathrm{OP}^2 - 2\mathrm{OP} \times \mathrm{OA} \cos \mathrm{COP} + \mathrm{CO}^2 = \mathrm{PC}^2$$

&c. &c. &c.

Hence,

$$\mathrm{OP}^{2m} - 2\mathrm{OP}^m \times \mathrm{OA}^m \cos m\,(\mathrm{AOP}) + \mathrm{OA}^{2m} = \mathrm{PA}^2 \times \mathrm{PB}^2 \times \mathrm{PC}^2 \times \&c.$$

and this is *De Moivre's property of the circle.*

(25.) If $\mathrm{AOP} = 0$, that is, if P be upon the radius through one of the points of division A, then $\cos m\,(\mathrm{AOP}) = 1$. Hence,

$$\mathrm{OP}^{2m} - 2\mathrm{OP}^m \times \mathrm{OA}^m + \mathrm{OA}^{2m} = \mathrm{PA}^2 \times \mathrm{PB}^2 \times \mathrm{PC}^2 \times \&c.$$

---

* Young's Trigonometry, p. 19.

consequently, extracting the square root of each member,

$$OP^m \sim OA^m = PA \times PB \times PC \times \&c.$$

If with centre O and radius OP a circle were to be described, the first member of this equation would remain the same wherever, in the circumference, P be taken; it follows, therefore, that *if to the vertices of a regular polygon lines be drawn from any point in the circumference of a concentric circle, their product will be constant.*

Moreover, since, when P is on OA, $OP \sim OA = PA$, we have

$$\frac{OP^m \sim OA^m}{OP \sim OA} = PB \times PC \times PD \times \&c.,$$

that is, by actual division,

$$OP^{m-1} + OP^{m-2}\,OA + \ldots . OA^{m-1} = PB \times PC \times PD \times \&c.$$

Let $OP = OA$; that is, let P be at A, then this equation becomes

$$m\,OP^{m-1} = PB \times PC \times PD \times \&c.;$$

so that, *if from one of the vertices of a regular polygon, lines be drawn to all the other vertices, the continued product of these* $m - 1$ *lines will be equal to* m *times the* $m - 1$*th power of the radius of the circumscribed circle.*

If the arcs AB, BC, &c. be bisected in A′, B′, &c. the circumference will be divided into $2m$ equal parts, and, by the equation deduced above,

$$OP^{2m} \sim OA^{2m} = PA \times PA' \times PB \times PB' \times \&c.$$

that is,

$$OP^{2m} \sim OA^{2m} = (OP^m \sim OA^m)\ PA' \times PB' \times \&c.$$

therefore,

$$\frac{OP^{2m} \sim OA^{2m}}{OP^m \sim OA^m} = OP^m + OA^m = PA' \times PB' \times PC' \times \&c.$$

and these are *Cotes's* properties of the circle.

(26.) If now we return to the expression (5), page 42, and suppose $x = \frac{\pi}{2}$, it becomes

$$\sqrt{-1} = e^{\frac{\pi}{2}\sqrt{-1}} \quad \therefore \log \sqrt{-1} = \frac{\pi}{2}\sqrt{-1}$$

and

$$(\sqrt{-1})^{\sqrt{-1}} = e^{-\frac{\pi}{2}}.$$

From the second of these equations we get

$$\pi = \frac{2 \log \sqrt{-1}}{\sqrt{-1}} = \frac{\log (\sqrt{-1})^2}{\sqrt{-1}} = \frac{\log -1}{\sqrt{-1}} = -\sqrt{-1} \log -1.$$

From the third

$$(\sqrt{-1})^{\sqrt{-1}} = 1 - \frac{\pi}{2} + \frac{1}{1 \cdot 2} \cdot \frac{\pi^2}{4} - \frac{1}{1 \cdot 2 \cdot 3} \cdot \frac{\pi^3}{8} + \&c.$$

$$= 1 - \frac{\pi}{2} + \frac{\pi^2}{2 \cdot 4} - \frac{\pi^3}{2 \cdot 4 \cdot 6} + \&c.$$

Two very singular results; first obtained by *John Bernoulli*.

Such conclusions are something more than mere matters of analytical curiosity; they show that however unintelligible certain mathematical forms may be in an undeveloped state, yet, when actually subjected to the operations symbolized, a clear and satisfactory interpretation may be evolved. The form $(\sqrt{-1})^{\sqrt{-1}}$ we see is nothing more than a brief and compact expression for certain operations which terminate in the series on the right hand; and we may therefore always substitute for such an expression this its interpretation. In like manner, the form $\frac{2}{\sqrt{-1}} \log \sqrt{-1}$ above, must be regarded as the symbolical expression for certain processes which, when performed, furnish the length of a semicircular arc of radius 1. To render this expression more readily developable, let us put $\frac{1+\sqrt{-1}}{1-\sqrt{-1}}$ for its equal $\sqrt{-1}$,

in $\log \sqrt{-1}$, and the symbol for π will then be

$$\pi = \frac{2}{\sqrt{-1}} \{\log (1 + \sqrt{-1}) - \log (1 - \sqrt{-1})\}$$

$$= \frac{2}{\sqrt{-1}} \{\sqrt{-1} - \frac{1}{2} (\sqrt{-1})^2 + \frac{1}{3} (\sqrt{-1})^3 -$$

$$\frac{1}{4} (\sqrt{-1})^4 + \&c.\} +$$

$$\frac{2}{\sqrt{-1}} \{-\sqrt{-1} - \frac{1}{2} (\sqrt{-1})^2 - \frac{1}{3} (\sqrt{-1})^3 -$$

$$\frac{1}{4} (\sqrt{-1})^4 - \&c.\}$$

that is,

$$\pi = 4 \{1 - \frac{1}{3} + \frac{1}{5} - \frac{1}{7} + \&c.\}$$

6. To develop $\tan x$.

$$y = \tan x, \text{ therefore } \ldots [y] = 0$$

$$\frac{dy}{dx} = \sec^2 x \ldots [\frac{dy}{dx}] = 1$$

$$\frac{d^2y}{dx^2} = 2 \sec^2 x \tan x \ldots [\frac{d^2y}{dx^2}] = 0$$

$$\frac{d^3y}{dx^3} = 2 \sec^2 x (1 + 3 \tan^2 x) \ldots [\frac{d^3y}{dx^3}] = 2.$$

We thus see that the first two terms of the development are $x + \frac{2x^3}{1 \cdot 2 \cdot 3}$, but we shall not continue the differentiation, since it does not make known the law of the series. The development will be more readily obtained by means of those already given for $\sin x$ and $\cos x$, as follows:

$$\tan x = \frac{x - \frac{x^3}{1 \cdot 2 \cdot 3} + \frac{x^5}{1 \cdot 2 \cdot 3 \cdot 4 \cdot 5} - \&c.}{1 - \frac{x^2}{2} + \frac{x^4}{1 \cdot 2 \cdot 3 \cdot 4} - \&c.}$$

therefore the development, found by actual division, is

$$\tan x = x + \frac{2x^3}{1\cdot 2\cdot 3} + \frac{16x^5}{1\cdot 2\cdot 3\cdot 4\cdot 5} + \&c.$$

but, to obtain the *law* of the terms and thus be enabled to continue them at pleasure, it will be best to apply the method of indeterminate coefficients. Assume therefore this fraction, equal to the series

$$A_1 x + A_3 x^3 + A_5 x^5 + A_7 x^7 + \&c.$$

Multiplying this by the denominator, we have this expression for the numerator, viz.

$$x - \frac{x^3}{1\cdot 2\cdot 3} + \frac{x^5}{1\cdot 2\cdot 3\cdot 4\cdot 5} - \&c. =$$

$$\begin{array}{l|l|l} A_1 x + A_3 & x^3 + & A_5 \\ -\frac{A_1}{1\cdot 2} & - & \frac{A_3}{1\cdot 2} \\ & + & \frac{A_1}{1\cdot 2\cdot 3\cdot 4} \end{array} \quad x^5 + \&c.$$

Hence, equating the coefficients of the like terms, there results

$A_1 = 1$ therefore $A_1 = 1$

$$A_3 - \frac{A_1}{1\cdot 2} = -\frac{1}{1\cdot 2\cdot 3} \quad \ldots \quad A_3 = \frac{A_1}{1\cdot 2} - \frac{1}{1\cdot 2\cdot 3} = \frac{2}{1\cdot 2\cdot 3}$$

$$A_5 - \frac{A_3}{1\cdot 2} + \frac{A_1}{1\cdot 2\cdot 3\cdot 4} = \frac{1}{1\cdot 2\cdot 3\cdot 4\cdot 5}, \quad A_5 = \frac{A_3}{1\cdot 2} - \frac{A_1}{1\cdot 2\cdot 3\cdot 4} +$$

$$\frac{1}{1\cdot 2\cdot 3\cdot 4\cdot 5} = \frac{16}{1\cdot 2\cdot 3\cdot 4\cdot 5}$$

&c. &c.

the law of these coefficients being such, that

$$A_{2n+1} = \frac{A_{2n-1}}{1\cdot 2} - \frac{A_{2n-3}}{1\cdot 2\cdot 3\cdot 4} + \ldots \pm \frac{A_1}{1\cdot 2\cdot 3 \ldots 2n} \ldots \mp$$

$$\frac{1}{1\cdot 2\cdot 3\ldots(2n+1)}.$$

7. To develop $\tan^{-1} x$.

$y = \tan^{-1} x$ . . . . therefore . . . . $[y] = 0$*

$$\frac{dy}{dx} = \frac{1}{1+x^2} \quad \ldots\ldots\ldots\ldots \quad [\frac{dy}{dx}] = 1$$

$$\frac{d^2y}{dx^2} = -\frac{2x}{(1+x^2)^2} \quad \ldots\ldots\ldots\ldots \quad [\frac{d^2y}{dx^2}] = 0$$

$$\frac{d^3y}{dx^3} = -\frac{2}{(1+x^2)^2} + \frac{2\cdot 4x^2}{(1+x^2)^3} \quad \ldots\ldots \quad [\frac{d^3y}{dx^3}] = -2$$

$$\frac{d^4y}{dx^4} = \frac{2^3x}{(1+x^2)^3} + \frac{2^4x}{(1+x^2)^3} - \frac{2^4\cdot 3x^3}{(1+x^2)^4} \quad \ldots \quad [\frac{d^4y}{dx^4}] = 0$$

$$\frac{d^5y}{dx^5} = \frac{2^3\cdot 3}{(1+x^2)^3} - \frac{2^5\cdot 3^2x^2}{(1+x^2)^4} + \frac{2^7\cdot 3x^4}{(1+x^2)^5} \quad \ldots \quad [\frac{d^5y}{dx^5}] = 2^3\cdot 3$$

&c. &c.

$$\therefore\ y = \tan y - \tfrac{1}{3}\tan^3 y + \tfrac{1}{5}\tan^5 y - \tfrac{1}{7}\tan^7 y + \&c.$$

This development is obtainable with far less trouble by means of the rule at page 37. Thus, since by division

$$\frac{1}{1+x^2} = 1 - x^2 + x^4 - x^6 + \&c.,$$

we have, by the rule referred to,

$$y = \tan y - \tfrac{1}{3}\tan^3 y + \tfrac{1}{5}\tan^5 y - \tfrac{1}{7}\tan^7 y + \&c.$$

tan y being put for x, as before.

---

* The arc whose tangent is 0 may be either 0 or any positive or negative multiple of π. We shall, however, suppose here that the arc to be developed does not exceed a quadrant; it must, therefore, become 0 when x becomes 0.

If $y = 45°$, then $\tan y = 1$;

$$\therefore \text{ arc } 45° = 1 - \tfrac{1}{3} + \tfrac{1}{5} - \tfrac{1}{7} + \&c.$$

(27.) From this series an approximation may be made to the circumference of a circle, but, from its very slow convergency, it is not eligible for this purpose.* *Euler* has obtained from the above general development a series much more suitable, by help of the known formula, (*Young's Trig.* page 48,)

$$\tan(a + b) = \frac{\tan a + \tan b}{1 - \tan a \tan b},$$

for, when $a + b = 45°$, $\tan(a + b) = 1$; therefore,

$$\tan a + \tan b = 1 - \tan a \tan b.$$

If either $\tan a$ or $\tan b$ were given, the other would be determinable from this equation. Thus, if we suppose,

$$\tan a = \frac{1}{n}, \text{ then } \frac{1}{n} + \tan b = 1 - \frac{\tan b}{n}, \therefore \tan b = \frac{n-1}{n+1}.$$

Now the value of n is arbitrary, and our object is to assume it so that the sum of the series, expressing the arcs a, b, in terms of their tangents, may be the most convergent. This value appears to be $n = 2$ or $n = 3$; therefore, taking $n = 2$, we have

$$\tan a = \tfrac{1}{2}, \quad \tan b = \tfrac{1}{3}.$$

Hence, substituting in the general development a for y, and $\frac{1}{2}$ for $\tan y$, and then again b for y, and $\frac{1}{3}$ for $\tan b$, the sum of the resulting series will express the length of the arc $a + b = 45°$; that is,

$$\text{arc } 45° = \begin{cases} \dfrac{1}{2} - \dfrac{1}{3 \cdot 2^3} + \dfrac{1}{5 \cdot 2^5} - \dfrac{1}{7 \cdot 2^7} + \&c. \\ + \dfrac{1}{3} - \dfrac{1}{3 \cdot 3^3} + \dfrac{1}{5 \cdot 3^5} - \dfrac{1}{7 \cdot 3^7} + \&c. \end{cases}$$

* See Note (A) at the end, where an easy and tolerably rapid mode of approximation is given.

(28.) Another form of development, still more convergent than this, was obtained by *Machin* from the formula

$$\tan 2a = \frac{2\tan a}{1-\tan^2 a}.$$

For put $\tan a = \frac{1}{5}$, then $\tan 2a = \frac{5}{12}$, therefore $2a < 45°$, because $\tan 45° = 1$: from this value of $2a$ we deduce

$$\tan 4a = \frac{2\tan 2a}{1-\tan^2 2a} = \frac{120}{119}$$

$$\therefore\ 4a > 45°.$$

Let now $4a = A$, $45° = B$, $A - B = b$ = excess of $4a$ above $45°$; then we have $45° = A - b$. But (*Trig.* p. 48,)

$$\tan(A-B) = \tan b = \frac{\tan A - \tan B}{1+\tan A \tan B} = \frac{1}{239}.$$

Consequently, if in the general development we replace y by a, and $\tan y$ by $\frac{1}{5}$, and then multiply by 4, we shall have the length of the arc $4a$; and, since this arc exceeds $45°$ by the arc b, if we subtract the development of this latter, which is given by substituting $\frac{1}{239}$ for $\tan y$, the remainder will be the true development of $45°$. Thus

$$45° = \begin{cases} 4\left(\frac{1}{5} - \frac{1}{3\cdot 5^3} + \frac{1}{5\cdot 5^5} - \frac{1}{7\cdot 5^7} + \&c.\right) \\ -\left(\frac{1}{239} - \frac{1}{3\cdot 239^3} + \frac{1}{5\cdot 239^5} - \&c.\right) \end{cases}$$

This series is very convergent, and, by taking about 8 terms in the first row and 3 in the second, we find, for the length of the semicircle, the following value, viz.

$$\pi = 3\cdot 141592653589793.$$

If we take but three terms of the first and only one of the second, we shall have $\pi = 3\cdot 1416$, the approximation usually employed in practice.

(29.) The following examples are subjoined for the exercise of the student:

8. To develop $y = \sin^{-1} x$.

$$y = \sin y + \frac{\sin^3 y}{2 \cdot 3} + \frac{1 \cdot 3 \sin^5 y}{2 \cdot 4 \cdot 5} + \frac{1 \cdot 3 \cdot 5 \sin^7 y}{2 \cdot 4 \cdot 6 \cdot 7} + \&c.$$

9. To develop $y = \cos^{-1} x$.

$$y = \frac{\pi}{2} - \cos y - \frac{\cos^3 y}{2 \cdot 3} - \frac{1 \cdot 3 \cos^5 y}{2 \cdot 4 \cdot 5} - \frac{1 \cdot 3 \cdot 5 \cos^7 y}{2 \cdot 4 \cdot 6 \cdot 7} - \&c.$$

10. To develop $y = \cot x$ by the method of indeterminate coefficients, as in example 6.

$$\cot x = \frac{1}{x} - \frac{x}{3} - \frac{x^3}{3^2 \cdot 5} - \frac{2x^5}{3^2 \cdot 5 \cdot 7} - \&c.$$

11. To develop $y = (a + bx + cx^2 + \&c.)^n$

$$(a + bx + cx^2 + \&c.)^n =$$

$$a^n + na^{n-1}bx + \frac{n(n-1)}{2} a^{n-2} b^2 \left| x^2 + \frac{n(n-1)(n-2)}{2 \cdot 3} a^{n-3} b^3 \right| x^3 + \&c.$$

$$+ na^{n-1}c \qquad + n(n-1) a^{n-2} bc$$

$$+ na^{n-1} d$$

This is the multinomial theorem of De Moivre. It is given in a very convenient practical form in the Treatise on Algebra, page 235.

CHAPTER IV.

ON TAYLOR'S THEOREM, AND ON THE DIFFERENTIATION AND DEVELOPMENT OF IMPLICIT FUNCTIONS.

(30.) In the second chapter we established the *form* of the general development of the function $F(x+h)$. We here propose to investigate Taylor's theorem, which is an expression exhibiting the *actual* development of the same function. The following lemma, must, however, be premised, viz. that if in any function of $p+q$ one of the quantities p, q, is variable, and the other constant, we may determine the several differential coefficients, without enquiring which is the constant and which the variable; for these coefficients will be the same, whichever be variable. This principle is almost axiomatic. For as the function contains but *one* variable, we may put $p+q=x$ or $F(p+q)=F(x)$, and whichever of the parts p, q, takes the increment h, the result $F(x+h)$ is necessarily the same; hence the development of this function is the same on either hypothesis, and therefore the second term of that development, and hence also the differential coefficient. The first differential coefficient being the same, the succeeding must be the same; therefore, generally,

$$\frac{d^n F(p+q)}{dp^n} = \frac{d^n F(p+q)}{dq^n},$$

whatever be the value of n.

Let now $y=F(x)$, and $Y=F(x+h)$, and assume, agreeably to art. (4),

$$Y = y + Ah + Bh^2 + Ch^3 + \&c.$$

A, B, C, &c. being unknown functions of x, which it is now required to determine.

Suppose, first, h to be variable and x constant; then, differentiating on that supposition, we have

$$\frac{dY}{dh} = A + 2Bh + 3Ch^2 + \&c.$$

Suppose, secondly, that x is variable and h constant; then the differential coefficient is

$$\frac{dY}{dx} = \frac{dy}{dx} + \frac{dA}{dx}h + \frac{dB}{dx}h^2 + \frac{dC}{dx}h^3 + \&c.$$

But, by the lemma, these two differential coefficients are identical; hence, equating the coefficients of the like powers of h, there results

$$A = \frac{dy}{dx}, \quad B = \frac{dA}{2dx}, \quad C = \frac{dB}{3dx}, \quad \&c.$$

that is,

$$A = \frac{dy}{dx}, \quad B = \frac{d^2y}{dx^2}\,\frac{1}{2}, \quad C = \frac{d^3y}{dx^3}\,\frac{1}{2\cdot 3}, \quad \&c.$$

Hence the required development is

$$Y = y + \frac{dy}{dx}\,\frac{h}{1} + \frac{d^2y}{dx^2}\,\frac{h^2}{1\cdot 2} + \frac{d^3y}{dx^3}\,\frac{h^3}{1\cdot 2\cdot 3} + \&c.$$

If h is negative, the signs of the alternate terms will be negative.

When we wish for the development of the function $Y = F(x + h)$ in any particular state; that is, when x takes a given value, we have only to substitute this value for x in the general expressions for the coefficients previously determined, and we shall have the development according to the above form; that is, provided, of course, that the development in such form is possible. But if the value chosen for x render the development impossible, the impossibility will be intimated to us from the circumstance of some of the terms becoming infinite, as explained in *art.* (4). It may, however, be proper here to remark, that even in these cases of impossibility, the leading terms of the development, as given by Taylor's theorem, are still true as far as the first term that becomes infinite. But as we propose to devote hereafter an entire chapter to the examination of the *failing cases* of Taylor's theorem, we shall not enter into the enquiry here.

(31.) The theorem of Maclaurin may be easily deduced from that of Taylor thus:

Let x take the particular value $x = 0$, then

$$[Y] = F(h) = [F(x)] + [\frac{dF(x)}{dx}]\ \frac{h}{1} + [\frac{d^2F(x)}{dx^2}]\ \frac{h^2}{1\cdot 2} +$$

$$[\frac{d^3F(x)}{dx^3}]\ \frac{h^3}{1\cdot 2\cdot 3} + \&c.$$

Now each of these coefficients is constant, and therefore independent of the value of h; hence h may take any value whatever, without affecting these coefficients; we may therefore call it x, it being observed that although x appears in the *notation* of the coefficients, it does not appear in the coefficients themselves. It follows, therefore, that

$$F(x) = [F(x)] + [\frac{dF(x)}{dx}]\ x + [\frac{d^2F(x)}{dx^2}]\ \frac{x^2}{1\cdot 2} + [\frac{d^3F(x)}{dx^3}]\ \frac{x^3}{1\cdot 2\cdot 3} + \&c.$$

which is Maclaurin's theorem, before investigated.

EXAMPLES.

(32.) 1. To develop $\sin(x+h)$ in a series of powers of the arc h. Let $y = \sin x$ $\therefore$ $\frac{dy}{dx} = \cos x$, $\frac{d^2y}{dx^2} = -\sin x$, $\frac{d^3y}{dx^3} = -\cos x$, &c; hence, by Taylor's theorem,

$$\sin(x+h) = \sin x + \cos x\, h - \sin x\, \frac{h^2}{1\cdot 2} - \cos x\, \frac{h^3}{1\cdot 2\cdot 3} + \&c.$$

$$= \sin x\,(1 - \frac{h^2}{1\cdot 2} + \frac{h^4}{1\cdot 2\cdot 3\cdot 4} - \&c.)$$

$$+ \cos x\,(h - \frac{h^3}{1\cdot 2\cdot 3} + \frac{h^5}{1\cdot 2\cdot 3\cdot 4\cdot 5} - \&c.)$$

The series within the parentheses are respectively equal to $\cos h$ and $\sin h$ (p. 39,) so that the preceding development merely expresses the known property

$$\sin(x+h) = \sin x \cos h + \cos x \sin h.$$

2. To develop $\cos(x+h)$.

$$y = \cos x \;\therefore\; \frac{dy}{dx} = -\sin x, \quad \frac{d^2y}{dx^2} = -\cos x, \quad \frac{d^3y}{dx^3} = \sin x, \;\&c.$$

Hence

$$\cos(x+h) = \cos x - \sin x\, h - \cos x \frac{h^2}{1\cdot 2} + \sin x \frac{h^3}{1\cdot 2\cdot 3} + \&c.$$

$$= \cos x \left(1 - \frac{h^2}{1\cdot 2} + \frac{h^4}{1\cdot 2\cdot 3\cdot 4} - \&c.\right)$$

$$- \sin x \left(h - \frac{h^3}{1\cdot 2\cdot 3} + \frac{h^5}{1\cdot 2\cdot 3\cdot 4\cdot 5} - \&c.\right)$$

which merely brings us back to the other fundamental property

$$\cos(x+h) = \cos x \cos h - \sin x \sin h.$$

By putting $h = (m-1)x$, these two properties become

$$\sin mx = \sin x \cos(m-1)x + \sin(m-1)x \cos x,$$

$$\cos mx = \cos x \cos(m-1)x - \sin x \sin(m-1)x,$$

two equations which will be employed to abridge the expressions for the differential coefficients in the next Example.

3. To develop $\tan^{-1}(x+h)$.

$$y = \tan^{-1} x$$

$$\frac{dy}{dx} = \frac{1}{\sec^2 y} = \cos^2 y$$

$$\frac{d^2y}{dx^2} = -2 \sin y \cos y \frac{dy}{dx} = -\sin 2y \cos^2 y$$

$$\frac{d^3y}{dx^3} = -2(\cos 2y \cos^2 y - \sin 2y \sin y \cos y)\frac{dy}{dx}$$

$$= -2 \cos 3y \cos y \frac{dy}{dx}$$

$$= -2 \cos 3y \cos^3 y$$

$$\frac{d^4y}{dx^4} = 2 \cdot 3 \,(\sin 3y \cos^3 y + \cos 3y \sin y \cos^2 y) \frac{dy}{dx}$$

$$= 2 \cdot 3 \sin 4y \cos^2 y \frac{dy}{dx} = 2 \cdot 3 \sin 4y \cos^4 y$$

$$\frac{d^5y}{dx^5} = 2 \cdot 3 \cdot 4 \,(\cos 4y \cos^4 y - \sin 4y \sin y \cos^3 y) \frac{dy}{dx}$$

$$= 2 \cdot 3 \cdot 4 \cos 5y \cos^3 y \frac{y}{dx} = 2 \cdot 3 \cdot 4 \cos 5y \cos^5 y$$

&c. &c.

Hence

$$\tan^{-1}(x + h) = y + \cos^2 y\, h - \frac{\sin 2y \cos^2 y}{2} h^2 -$$

$$\frac{\cos 3y \cos^3 y}{3} h^3 + \frac{\sin 4y \cos^4 y}{4} h^4 + \frac{\cos 5y \cos^5 y}{5} h^5 - \&c.$$

4. To develop $\tan(x + h)$ according to the powers of h.

$$\tan(x + h) = \tan x + \sec^2 x \,.\, h + 2 \sec^2 x \tan x \frac{h^2}{1 \cdot 2} +$$

$$2 \sec^2 x (1 + 3 \tan^2 x) \frac{h^3}{1 \cdot 2 \cdot 3} + \&c.$$

(33.) By means of the theorem of Taylor may be obtained a very commodious and useful form for the representation and subsequent determination of the differential coefficient, when the function is complicated, thus:

Let $u = F(y)$, y being any function of x, which we may represent by $y = f(x)$, and let it be required to find the expression for $\frac{du}{dx}$. Let x take the increment h; then, since $y = f(x)$, the corresponding increment of y will, by Taylor's theorem, be

$$\frac{dy}{dx} h + \frac{d^2y}{dx^2} \frac{h^2}{1 \cdot 2} + \frac{d^3y}{dx^3} \frac{h^3}{1 \cdot 2 \cdot 3} + \&c.$$

Call this increment k, then the corresponding increment of u will be

$$F(y+k)-F(y)=\frac{du}{dy}k+\frac{d^2u}{dy^2}\frac{k^2}{1\cdot 2}+\&c.$$

that is, restoring the value of k,

$$F(y+k)-F(y)=\frac{du}{dy}\left\{\frac{dy}{dx}h+\frac{d^2y}{dx^2}\frac{h^2}{1\cdot 2}+\&c.\right\}$$
$$+\frac{1}{1\cdot 2}\frac{d^2u}{dy^2}\left\{\frac{dy}{dx}h+\frac{d^2y}{dx^2}\frac{h^2}{1\cdot 2}+\&c.\right\}^2$$
$$+\&c.$$

Hence, dividing by the increment h of the independent variable, and taking the limit as usual, we have

$$\frac{dF(y)}{dx}=\frac{du}{dx}=\frac{du}{dy}\frac{dy}{dx}\quad\therefore du=\frac{du}{dy}\frac{dy}{dx}dx.$$

It appears, therefore, that the differential coefficient $\frac{du}{dx}$ is found by differentiating the function, on the hypothesis that y is the independent variable, and then multiplying the coefficient thus obtained, by that derived from y considered as a function of x.

(34.) The following examples will suffice to illustrate this mode of finding the differential coefficient.

1. Let $u=a^y$ where $y=b^x$.

$$\frac{du}{dy}=a^y\log a,\quad\frac{dy}{dx}=b^x\log b$$

$$\therefore\frac{du}{dx}=\frac{du}{dy}\frac{dy}{dx}=a^{b^x}b^x\log a\log b.$$

2. Let $u=\log y$, where $y=\log x$.

$$\therefore\frac{du}{dy}=\frac{1}{y},\quad\frac{dy}{dx}=\frac{1}{x}$$

$$\therefore\frac{du}{dx}=\frac{du}{dy}\frac{dy}{dx}=\frac{1}{x\log x}.$$

3. Let $u = \sin\left(\frac{x}{a+x}\right)^2$.

Put $\left(\frac{x}{a+x}\right)^2 = y$ $\therefore \frac{du}{dy} = \frac{d \sin y}{dy} = \cos y$,

$$\frac{dy}{dx} = \frac{2ax}{(a+x)^3} \quad \therefore \frac{du}{dx} = \cos\left(\frac{x}{a+x}\right)^2 \cdot \frac{2ax}{(a+x)^3}.$$

4. Let $u = \cot a^y$, y being $= \log \frac{x}{\sqrt{a^2+x^2}}$

$$\therefore \frac{du}{dy} = -\operatorname{cosec}^2 a^y \cdot a^y \log a, \quad \frac{dy}{dx} = \frac{a^2}{x(a^2+x^2)}$$

$$\therefore \frac{du}{dx} = -\operatorname{cosec}^2 a^y \cdot a^y \log a \cdot \frac{a^2}{x(a^2+x^2)}$$

(35.) Let us now take the more general function $u = \mathrm{F}(p, q)$, p and q being functions of x, and suppose that when x becomes $x + h$, p and q become $p + k$, and $q + k'$. Call this latter q'; then, in consequence of the proposed change in the independent variable, the function will become $\mathrm{F}(q', p + k)$, in which it is to be observed q' enters as if it were a constant, since it is unaffected by the increment k of the variable p.

Hence, by Taylor's theorem,

$$\mathrm{F}(q' p + k) = \mathrm{F}(q', p) + \frac{d\,\mathrm{F}(q', p)}{dp} k + \frac{d^2\,\mathrm{F}(q', p)}{dp^2} \frac{k^2}{1 \cdot 2} + \&c. \ldots . (1).$$

But if in $\mathrm{F}(q', p)$ we substitute for q' its equal $q + k'$, we have

$$\mathrm{F}(q', p) = \mathrm{F}(p, q + k') = u + \frac{du}{dq} k' + \frac{d^2u}{dq^2} \frac{k'^2}{1 \cdot 2} + \&c. \ldots . (2)$$

$$\therefore \frac{d\,\mathrm{F}(q', p)}{dp} = \frac{du}{dp} + \frac{d\frac{du}{dq}}{dp} k' + \&c. \ldots . (3).$$

and thus by continuing the differentiation may all the coefficients in (1)

be developed according to the powers of k'; but this first will be sufficient for our purpose.

Substitute in the first two terms of (1) their developments (2), (3); and we have

$$\mathrm{F}(q+k',\ p+k) = u + \frac{\mathrm{d}u}{\mathrm{d}q}k' + \&\mathrm{c.} + \frac{\mathrm{d}u}{\mathrm{d}p}k + \&\mathrm{c.} \ldots, \quad (4).$$

But k being the increment of the function p, arising from x taking the increment h; and k' being the increment which the function q takes from the same cause, it follows that

$$k = \frac{\mathrm{d}p}{\mathrm{d}x}h + \frac{\mathrm{d}^2p}{\mathrm{d}x^2}\,\frac{h^2}{1\cdot 2} + \&\mathrm{c.}, \quad k' = \frac{\mathrm{d}q}{\mathrm{d}x}h + \frac{\mathrm{d}^2q}{\mathrm{d}x^2}\,\frac{h^2}{1\cdot 2} + \&\mathrm{c.}$$

Hence, by substitution in (4), we have finally

$$\mathrm{F}(q+k',\ p+k) = u + \left\{\frac{\mathrm{d}u}{\mathrm{d}q}\,\frac{\mathrm{d}q}{\mathrm{d}x} + \frac{\mathrm{d}u}{\mathrm{d}p}\,\frac{\mathrm{d}p}{\mathrm{d}x}\right\}h + \&\mathrm{c.}$$

$$\therefore \frac{\mathrm{d}u}{\mathrm{d}x} = \frac{\mathrm{d}u}{\mathrm{d}q}\,\frac{\mathrm{d}q}{\mathrm{d}x} + \frac{\mathrm{d}u}{\mathrm{d}p}\,\frac{\mathrm{d}p}{\mathrm{d}x}.$$

(36.) Again: let there be three functions of x, viz. $v = \mathrm{F}(p, q, r)$; then, when x becomes $x+h$, let p, q, r become $p+k$, $q+k'$, $r+k''$, respectively; and put r' for the latter; then, in the function $\mathrm{F}(p+k, q+k', r')$, r' enters as a constant; hence, as above,

$$\mathrm{F}(p+k,\ q+k',\ r') = u + \left\{\frac{\mathrm{d}u}{\mathrm{d}q}\,\frac{\mathrm{d}q}{\mathrm{d}x} + \frac{\mathrm{d}u}{\mathrm{d}p}\,\frac{\mathrm{d}p}{\mathrm{d}x}\right\}h + \&\mathrm{c.}$$

where $u = \mathrm{F}(p, q, r')$ $\therefore$ putting $r + k''$ for r',

$$u = v + \frac{\mathrm{d}v}{\mathrm{d}r}k'' + \&\mathrm{c.}, \quad \frac{\mathrm{d}u}{\mathrm{d}q} = \frac{\mathrm{d}v}{\mathrm{d}q} + \frac{\mathrm{d}\,\frac{\mathrm{d}v}{\mathrm{d}r}}{\mathrm{d}q}k'' + \&\mathrm{c.}$$

$$\frac{\mathrm{d}u}{\mathrm{d}p} = \frac{\mathrm{d}v}{\mathrm{d}p} + \frac{\mathrm{d}\,\frac{\mathrm{d}v}{\mathrm{d}r}}{\mathrm{d}p}k'' + \&\mathrm{c.}$$

But $k'' = \frac{\mathrm{d}r}{\mathrm{d}x}h + \&\mathrm{c.}$, consequently,

$$\mathrm{F}(p+k, q+k', r+k'') = v + \left\{\frac{\mathrm{d}v}{\mathrm{d}r}\frac{\mathrm{d}r}{\mathrm{d}x} + \frac{\mathrm{d}v}{\mathrm{d}q}\frac{\mathrm{d}q}{\mathrm{d}x} + \frac{\mathrm{d}v}{\mathrm{d}p}\frac{\mathrm{d}p}{\mathrm{d}x}\right\} h$$

$$+ \&c.$$

$$\therefore \frac{\mathrm{d}v}{\mathrm{d}x} = \frac{\mathrm{d}v}{\mathrm{d}r}\frac{\mathrm{d}r}{\mathrm{d}x} + \frac{\mathrm{d}v}{\mathrm{d}q}\frac{\mathrm{d}q}{\mathrm{d}x} + \frac{\mathrm{d}v}{\mathrm{d}p}\frac{\mathrm{d}p}{\mathrm{d}x},$$

and so on for any number of functions. Hence the rule is to *differentiate the expression with regard to each of its constituent functions severally, as if all the others were constants; their sum will be the required differential.*

Cor. If p is simply x, then, in the function $u = \overset{\bullet}{\mathrm{F}}(x, q)$,

$$\frac{\mathrm{d}u}{\mathrm{d}x} = \frac{\mathrm{d}u}{\mathrm{d}x} + \frac{\mathrm{d}u}{\mathrm{d}q}\frac{\mathrm{d}q}{\mathrm{d}x};$$

and in the function $u = \mathrm{F}(x, q, r)$,

$$\frac{\mathrm{d}u}{\mathrm{d}x} = \frac{\mathrm{d}u}{\mathrm{d}x} + \frac{\mathrm{d}u}{\mathrm{d}q}\frac{\mathrm{d}q}{\mathrm{d}x} + \frac{\mathrm{d}u}{\mathrm{d}r}\frac{\mathrm{d}r}{\mathrm{d}x}.$$

(37.) We must not confound here the $\frac{\mathrm{d}u}{\mathrm{d}x}$ on the left, with that on the right, in these equations; for the former denotes the *total* differential coefficient, of which the latter forms but a part, and is therefore called a *partial* differential coefficient. It is to be regretted, however, that analysts are not agreed as to the best means of distinguishing total, from partial, differential coefficients; and accordingly, in most works on the calculus, the same symbol is applied indiscriminately to both; a circumstance likely to prove a frequent source of perplexity to the learner; and to avoid which we shall, throughout this volume, always distinguish the total differential coefficient by enclosing it in brackets; so that the two equations above will be written thus:

$$\left\{\frac{\mathrm{d}u}{\mathrm{d}x}\right\} = \frac{\mathrm{d}u}{\mathrm{d}x} + \frac{\mathrm{d}u}{\mathrm{d}q}\frac{\mathrm{d}q}{\mathrm{d}x}$$

$$\left\{\frac{\mathrm{d}u}{\mathrm{d}x}\right\} = \frac{\mathrm{d}u}{\mathrm{d}x} + \frac{\mathrm{d}u}{\mathrm{d}q}\frac{\mathrm{d}q}{\mathrm{d}x} + \frac{\mathrm{d}u}{\mathrm{d}r}\frac{\mathrm{d}r}{\mathrm{d}x}.$$

(38.) We shall now add a few examples, showing the application of the rule deduced in last article.

EXAMPLES.

1. Let $u = \cot x^y$ $\therefore \{\frac{du}{dx}\} = \frac{du}{dx} + \frac{du}{dy}\,\frac{dy}{dx}$.

Now

$$\frac{du}{dx} = -\operatorname{cosec}^2 x^y \frac{dx^y}{dx} = -\operatorname{cosec}^2 x^y \cdot y x^{y-1}$$

$$\frac{du}{dy} = -\operatorname{cosec}^2 x^y \frac{dx^y}{dy} = -\operatorname{cosec}^2 x^y \cdot x^y \log x$$

$$\therefore \{\frac{du}{dx}\} = -x^y \operatorname{cosec}^2 x^y \left(\frac{y}{x} + \log x \frac{dy}{dx}\right).$$

2. Let $u = \dfrac{a(x + \sqrt{x} + \sqrt[3]{x} + \sqrt[4]{x})}{\sqrt{x^2 + (x + \sqrt{x} + \sqrt[3]{x} + \sqrt[4]{x})^2}}$.

Put $x + \sqrt{x} + \sqrt[3]{x} + \sqrt[4]{x} = q$ $\therefore$ $u = \mathrm{F}(x, q)$, and

$$\frac{du}{dx} = -\frac{aqx}{(x^2 + q^2)^{\frac{3}{2}}}$$

$$\frac{du}{dq} = \frac{ax^2}{(x^2 + q^2)^{\frac{3}{2}}}, \text{ and } \frac{dq}{dx} = 1 + \frac{1}{2\sqrt{x}} + \frac{1}{3x^{\frac{2}{3}}} + \frac{1}{4x^{\frac{3}{4}}};$$

hence

$$\{\frac{du}{dx}\} = \frac{ax^2}{(x^2 + q^2)^{\frac{3}{2}}}\left(1 + \frac{1}{2x^{\frac{1}{2}}} + \frac{1}{3x^{\frac{2}{3}}} + \frac{1}{4x^{\frac{3}{4}}}\right) - \frac{aqx}{(x^2 + q^2)^{\frac{3}{2}}}.$$

3. Let $u = \log \tan \dfrac{x}{y}$, y being a function of x.

$$\frac{du}{dx} = \frac{\mathrm{d} \tan \frac{x}{y}}{\tan \frac{x}{y}\, dx} = \frac{\sec^2 \frac{x}{y} \cdot \frac{1}{y}}{\tan \frac{x}{y}} = \frac{1}{y \sin \frac{x}{y} \cos \frac{x}{y}}$$

$$\frac{du}{dy} = \frac{d \tan \frac{x}{y}}{\tan \frac{x}{y} \, dy} = - \frac{\sec^2 \frac{x}{y} \cdot \frac{x}{y^2}}{\tan \frac{x}{y}} = - \frac{x}{y^2 \sin \frac{x}{y} \cos \frac{x}{y}};$$

hence

$$\{\frac{du}{dx}\} = \frac{y - x \frac{dy}{dx}}{y^2 \sin \frac{x}{y} \cos \frac{x}{y}} \cdot \frac{dy}{dx}.$$

4. Let $u = \log (x - a + \sqrt{x^2 - 2ax})$

$$\therefore \{\frac{du}{dx}\} = \frac{1}{\sqrt{x^2 - 2ax}}.$$

5. Let $u = (\cos x)^{\sin x}$

$$\therefore \{\frac{du}{dx}\} = (\cos x)^{\sin x} \, (\cos x \log \cos x - \frac{\sin^2 x}{\cos x}).$$

Implicit Functions.

(39.) Hitherto we have treated of *explicit* functions only, or those whose forms are supposed to be given. We shall now consider *implicit* functions, or those in which the relation between the independent variable x, and function y, is *implied* in an equation between the two, and which may be generally expressed by

$$F(x, y) = 0.$$

The deductions in article (36) will enable us very readily to find the coefficient $\frac{dy}{dx}$ from such equations, without being under the necessity of solving them, a thing indeed often impossible.

If we turn to the corollary in the article just referred to, and substitute y for q, we find

$$\{\frac{du}{dx}\} = \frac{du}{dx} + \frac{du}{dy} \frac{dy}{dx}.$$

But here $u = \mathrm{F}(x, y) = 0$, therefore $\{\frac{\mathrm{d}u}{\mathrm{d}x}\} = 0$, for $u' - u$ being always 0, $\frac{u' - u}{h}$ is always 0; hence

$$\frac{\mathrm{d}u}{\mathrm{d}x} + \frac{\mathrm{d}u}{\mathrm{d}y}\,\frac{\mathrm{d}y}{\mathrm{d}x} = 0.$$

This equation shows that if we differentiate the several terms contained in the function u with respect to x only, and then with respect to y regarded as a function of x, we may equate the sum of the results to zero, and thus determine $\frac{\mathrm{d}y}{\mathrm{d}x}$; or we may employ for the same purpose the following formula immediately deduced from that above, viz.

$$\frac{\mathrm{d}y}{\mathrm{d}x} = -\frac{\mathrm{d}u}{\mathrm{d}x} \div \frac{\mathrm{d}u}{\mathrm{d}y};$$

which shows that after having transposed the terms all to one side of the equation, we must differentiate the expression as if y were a constant, and then divide the resulting coefficient, taken with a contrary sign, by that derived from the same expression, on the supposition that x is a constant.

EXAMPLES.

1. Let $y^2 - 2mxy + x^2 - a = 0$.

Differentiating the several terms according to the first formula, we have

$$2y\,\frac{\mathrm{d}y}{\mathrm{d}x} - 2my - 2mx\,\frac{\mathrm{d}y}{\mathrm{d}x} + 2x = 0$$

$$\therefore\ 2(y - mx)\,\frac{\mathrm{d}y}{\mathrm{d}x} = 2(my - x)$$

$$\therefore\ \frac{\mathrm{d}y}{\mathrm{d}x} = \frac{my - x}{y - mx}$$

By the second formula we have

$$-\frac{du}{dx}=2my-2x,\ \frac{du}{dy}=2y-2mx\ \therefore\ \frac{dy}{dx}=\frac{my-x}{y-mx}.$$

2. Let $x^3+3axy+y^3=0$.

$$\therefore\ -\frac{du}{dx}=-3x^2-3ay,\ \frac{du}{dy}=3ax+3y^2\ \therefore\ \frac{dy}{dx}=-\frac{x^2+ay}{ax+y^2}.$$

If the second differential coefficient be required,* we have

$$\frac{d^2y}{dx^2}=-\frac{(ax+y^2)\,(2x+a\frac{dy}{dx})-(x^2+ay)\,(a+2y\frac{dy}{dx})}{(ax+y^2)^2};$$

or, substituting for $\frac{dy}{dx}$ its value just found,

$$\frac{d^2y}{dx^2}=-\frac{(ax+y^2)(2ax^2+2xy^2-ax^2-a^2y)-(x^2+ay)(a^2x+ay^2-2x^2y-2ay^2)}{(ax+y^2)^3}$$

$$=-\frac{2xy^4+6ax^2y^2+2x^4y-2a^3xy}{(ax+y^2)^3}$$

$$=-\frac{2xy\,(y^3+3axy+x^3)-2a^3xy}{(ax+y^2)^3},$$

that is, since $x^3+3axy+y^3=0$,

$$\frac{d^2y}{dx^2}=\frac{2a^3xy}{(ax+y^2)^3}.$$

3. Let $my^3-xy=m$, to develop y, according to the ascending powers of x,

$$-\frac{du}{dx}=y,\ \frac{du}{dy}=3my^2-x\ \therefore\ \frac{dy}{dx}=\frac{y}{3my^2-x},$$

therefore, calling the successive differential coefficients by p, q, r, &c.

---

* As the original equation is identically 0, that is, it is satisfied for *every* value of x, the successive differential expressions deduced from it are severally 0, (see p. 9.)

$$q = \frac{y - 3my^2p - xp}{(3my^2 - x)^2},$$

$$r = -\frac{3my^2q + 2 \cdot 3\, myp^2 + xq}{(3my^2 - x)^2} - \frac{(y - 3my^2p - xp)\,(12myp - 2)}{(3my^2 - x)^3}$$

&c. &c.

$$\therefore\ [y] = 1,\quad [p] = \frac{1}{3m},\quad [q] = 0,\quad [r] = -\frac{2}{3^3 m^3}\ \&c.$$

Therefore, by Maclaurin's theorem,

$$y = 1 + \frac{x}{3m} - \frac{x^3}{3^4 m^3} + \&c.$$

4. Let $y^3x - m^3(y + x) = 0$, to develop y according to the ascending powers of x.

Representing, as in last example, the successive differential coefficients by p, q, r, &c. we have

$$p = -\frac{y^3 - m^3}{3xy^3 - m^3}\ \therefore\ [p] = -1$$

$$q = -\frac{3y^2p}{3xy^2 - m^3} + (y^3 - m^3) \cdot \frac{3y^2 + 2 \cdot 3\, xyp}{(3xy^2 - m^3)^2}\ \therefore\ [q] = 0,$$

$$\therefore\ [r] = -\frac{2 \cdot 3\, yp^2}{3xy^2 - m^3} - \frac{2 \cdot 3\, m^3(2yp + xp^2)}{(3xy^2 - m^3)^3} = 0,$$

$$\therefore\ [s] = -\frac{2 \cdot 3p^3}{3xy^2 - m^3} - \frac{2 \cdot 3 \cdot 3\, m^3p^2}{(3xy^2 - m^3)^2} = -\frac{2 \cdot 3 \cdot 4}{m^3}$$

&c. &c.

Hence, by Maclaurin's theorem,

$$y = -x - \frac{x^4}{m^3} - \frac{3x^7}{m^6} - \&c.$$

5. Let $y^2 + 2xy + x^2 = a^2$, to find $\frac{dy}{dx}$.

$$\therefore\ \frac{dy}{dx} = -1.$$

6. Let $\frac{y^2}{(x-a)^2} = 2y\,\frac{\sqrt{x-b}}{x-a} - x + b$, to find $\frac{dy}{dx}$

$$\therefore \frac{dy}{dx} = \frac{3x-2b-a}{2\sqrt{x-b}}.$$

7. Let $y^3 - 3y + x = 0$, to develop y according to the ascending powers of x,

$$y = \frac{x}{3} + \frac{x^3}{3^4} + \frac{x^5}{3^6} + \&c.$$

8. Let $my^3x - y = m$, to develop y according to the ascending powers of x.

$$y = -m - m^4x - 3m^7x^2 - \&c.$$

9. Let $my^3 - x^3y = mx^3$, to develop y in a series of *descending** powers of x,

$$y = -m - \frac{m^4}{x^3} - \frac{3m^7}{x^6} - \&c.$$

CHAPTER V.

ON VANISHING FRACTIONS.

(40.) It is here proposed to determine what the value of a fraction $\frac{F(x)}{f(x)}$ becomes in the case in which, by giving a particular value, a, to the variable, both numerator and denominator vanish, the fraction then taking the form

$$\frac{F(a)}{f(a)} = \frac{0}{0}.$$

---

* This will be effected by substituting $\frac{1}{z}$ for x^3, which will transform the equation into $my^3z - y = m$, then developing y according to the ascending powers of z, and afterwards restoring the value of x.

As such a form can thus arise only from the circumstance of the same factor $x - a$ being common to both numerator and denominator, it is plain that if we can by any means eliminate this factor before our substitution of a for x, we shall then obtain the true value of the fraction.

Sometimes the vanishing factor is manifest at sight, and may be immediately expunged, as, for instance, in the fractions

$$\frac{(x-a)x}{bx-ab}, \quad \frac{x^2-a^2}{(a-x)^2}, \quad \frac{a^2-2ax+x^2}{a^2-x^2}, \text{ \&c.}$$

which each becomes $\frac{0}{0}$ when $x = a$, and which each obviously contain the factor $x - a$ in both numerator and denominator. In these cases, therefore, we at once see that the values of the fractions when $x = a$ are, severally,

$$\frac{a}{b}, -\infty, 0, \text{ \&c.}$$

In certain other cases the value, although not so easily seen as in the foregoing instances, may, nevertheless, be soon ascertained, by performing a few obvious transformations on the proposed fractions. Take the following example:

$$\frac{\sqrt{x-\sqrt{a}+\sqrt{(x-a)}}}{\sqrt{x^2-a^2}}$$

which becomes $\frac{0}{0}$ when $x = a$.

This fraction is the same as

$$\frac{\sqrt{x-\sqrt{a}}}{\sqrt{x^2-a^2}}+\frac{1}{\sqrt{x+a}}$$

and the first of these is the same as

$$\frac{x-a}{\sqrt{x^2-a^2}\,(\sqrt{x}+\sqrt{a})}=\frac{\sqrt{x-a}}{\sqrt{x+a}\,(\sqrt{x}+\sqrt{a})}$$

and this, when $x = a$, is $= 0$, therefore the value of the proposed fraction when $x = a$ is $\frac{1}{\sqrt{2a}}$.

(41) Before we proceed to discuss the general theory of these fractions it may be proper to apprize the student that some able mathematicians dispute the legitimacy of the conclusions to which we have arrived in the examples we have just been considering; and maintain that the elimination of the vanishing factor, as for instance $x - a$, in such a case as $\frac{x^2 - a^2}{x - a}$, is a mere artificial contrivance, to bring out, in the hypothesis of $x = a$, *one* only of the innumerable values which the expression is competent to give, seeing that it assumes the indeterminate form $\frac{a^2 - a^2}{a - a}$. It is overlooked, however, that the questions "What is the value of $\frac{x^2 - a^2}{x - a}$ when $x = a$?" and "What is the value of $\frac{a^2 - a^2}{a - a}$?" (which seem to have been confounded by the late Professor Woodhouse,) are perfectly distinct. The *value* of $\frac{x^2 - a^2}{x - a}$, when $x = a$, is not $\frac{a^2 - a^2}{a - a}$; this is only the symbolical *form* which $\frac{x^2 - a^2}{x - a}$ assumes in that case. The *value* of $\frac{x^2 - a^2}{x - a}$ is $x + a$; and this value, when $x = a$, is $2a$. The expression $\frac{x^2 - a^2}{x - a}$ implies an *operation* to be performed, viz. division, and the value is the *result* of that operation. When the form $\frac{a^2 - a^2}{a - a}$ is isolated, or detached from its interpretation, it is of course indeterminate; for the operation indicated requires that we determine such a quantity, P, that, when it is submitted to the reverse operation, $(a - a)$ P, the result may be $a^2 - a^2$; and it is easily seen that an infinite variety of suitable values for P exist; for it may be generally expressed by

$$P = a + a \pm p.$$

But the result of the operation $\frac{x^2-a^2}{x-a}$ is definite, and distinctly points out which of the above infinite variety is comprehended among the values of $\frac{x^2-a^2}{x-a}$.

It is one of the perfections of the symbolical language of analysis, that the form $\frac{a^2-a^2}{a-a}$ *is* indeterminate; for it may arise indifferently from $\frac{x^2-a^2}{x-a}$, $\frac{ax-a^2}{x-a}$, $\frac{ax-x^2}{x-a}$, $\frac{x^2-ax}{x-a}$, &c.: and thus it is very properly ambiguous for want of sufficient data.

Sometimes the form $\frac{0}{0}$ may arise in the result of an algebraic process from introducing *two* hypotheses among the disposable quantities; one causing the numerator to vanish, and the other causing the denominator to vanish. As the form in such a case is not due to a common vanishing factor in numerator and denominator, its discussion does not properly belong to the present chapter; for by the term *vanishing fraction* is to be understood a fraction which merges into the form $\frac{0}{0}$, in virtue of the simultaneous evanescence of a factor, common to numerator and denominator. We may, however, just observe, with respect to the other case of the symbol $\frac{0}{0}$, that it may be an indication that the hypotheses which have led to it are such as to cause one or more of the original analytical conditions to disappear; it will be proper, therefore, to ascertain, by an examination of those conditions, if such be the case; and if it be, we are then entitled to conclude that the solutions satisfying the conditions under the proposed hypotheses are innumerable.*

---

* Thus in the Analytical Geometry, page 202, we have the equation

$$2(x-x')\cos A - 2y' = 0.$$

which restricts $x - x'$ to the value

$$x - x' = \frac{y'}{\cos A}.$$

Now if we admit the hypotheses $y' = 0$, $\cos A = 0$, this result takes the

If, however, conditions do not disappear, and $\frac{0}{0}$ be not accompanied with any definite interpretation, then we must infer that the solution is not accomplishable in the case proposed, by the means we have adopted.*

We have just mentioned the circumstance of the symbol $\frac{0}{0}$ arising from suppositions which, at the same time, destroy original analytical conditions. It will readily occur to the student that in vanishing fractions, properly so called, the hypothesis, in virtue of which numerator and denominator vanish, may affect the original analytical conditions of the enquiry in the same way; that is, the same value which renders the resulting fraction $\frac{0}{0}$, may *also* render the problem indeterminate, by removing restrictions which had all along governed our several deductions, up to the last inclusively. It will be advisable, therefore, in analytical investigations, when any proposed hypothesis, introduced into the result, leads to a vanishing fraction, to observe the effect of that hypothesis upon the original conditions.

We shall now proceed to investigate methods for determining, in all cases, the value of a fraction which takes the form $\frac{0}{0}$, in consequence of giving a particular value to some quantity common to both numerator and denominator. Of these methods the most direct and general depend upon the differential calculus, and upon the algebraical development of functions, and require that we premise the following lemma, viz.

In the general development

$$\mathrm{F}(x+h) = \mathrm{F}(x) + \frac{\mathrm{d}\,\mathrm{F}(x)}{\mathrm{d}x}\,h + \frac{\mathrm{d}^2\mathrm{F}(x)}{\mathrm{d}x^2}\,\frac{h^2}{1\cdot 2} + \&c.$$

it is impossible that any particular value given to x can cause $\mathrm{F}(x)$, and at the same time *all* the differential coefficients, to vanish.

For if such could be the case, then, for that particular value, a, we should have

---

form $\frac{0}{0}$, and we see that the same hypotheses renders the conditional equation indeterminate. In this case, therefore, the values of $x - x'$ are innumerable, which is the deduction in the place referred to.

* See an instance in the Algebra, page 219, Ex. 2.

$$F(a+h)=0,$$

whatever be the value of h; but $F(a)=0$, consequently, whatever values put for a satisfy this equation, the same values put for $a+h$ must satisfy the preceding equation. In other words, $a=a+h$, which cannot be, unless $h=0$, whereas the hypothesis supposes the equations to exist independently of particular values of h; that hypothesis is therefore inadmissible.

Let now, in the proposed fraction, x be changed into $x+h$; then, by developing both numerator and denominator, it becomes

$$\frac{F(x)+F'(x)h+F''(x)h^2+F'''(x)h^3+\&c.}{f(x)+f'(x)h+f''(x)h^2+f'''(x)h^3+\&c.} \quad \ldots\ldots (1),$$

where $F'(x)$, $F''(x)$, &c. $f'(x)$, $f''(x)$, &c. are put for the successive differential coefficients divided by as many of the factors $1\cdot 2\cdot 3$, &c. as there are accents.

If in this we substitute a for x, then, since both $F(x)$ and $f(x)$ vanish, the fraction becomes, after dividing numerator and denominator by h,

$$\frac{F'(a)+F''(a)h+F'''(a)h^2+\&c.}{f'(a)+f''(a)h+f'''(a)h^2+\&c.} \quad \ldots\ldots (2),$$

and this fraction, when $h=0$, must obviously be equal to $\frac{F(a)}{f(a)}$; that is

$$\frac{F(a)}{f(a)}=\frac{F'(a)}{f'(a)}.$$

If, however, both $F'(a)$ and $f'(a)$ are also 0, then, expunging these terms from the fraction (2), and dividing numerator and denominator again by h, we have, when $h=0$,

$$\frac{F(a)}{f(a)}=\frac{F''(a)}{f''(a)},$$

and so on, till we at length obtain for $\frac{F(a)}{f(a)}$ a fraction of which the numerator and denominator do not both vanish, and such a fraction we eventually shall obtain in virtue of the preceding lemma.

Hence the following rule to determine the value of a fraction whose

numerator and denominator both vanish when $x = a$, viz. *For the numerator and denominator substitute their first differential coefficients, their second differential coefficients, and so on till we obtain a fraction in which numerator and denominator do not both vanish, for $x = a$; this will be the true value of the vanishing fraction.*

EXAMPLES.

1. Required the value of $\frac{a^x - b^x}{x}$ when $x = 0$.

$$\frac{F'(x)}{f'(x)} = \log a \,.\, a^x - \log b \,.\, b^x \quad \therefore \quad \frac{F'(a)}{f'(a)} = \log a - \log b = \log \frac{a}{b}.$$

2. Required the value of $\frac{x^3 - 3x + 2}{x^4 - 6x^2 + 8x - 3}$ when $x = 1$.

$$\frac{F'(x)}{f'(x)} = \frac{3x^2 - 3}{4x^3 - 12x + 8} \quad \therefore \quad \frac{F'(a)}{f'(a)} = \frac{0}{0} \quad \therefore$$

differentiating again,

$$\frac{F''(x)}{f''(x)} = \frac{6x}{12x^2 - 12} \quad \therefore \quad \frac{F''(a)}{f''(a)} = \frac{1}{0} = \infty.$$

3. Required the value of $\frac{1 - \sin x + \cos x}{\sin x + \cos x - 1}$ when $x = 90°$.

$$\frac{F'(x)}{f'(x)} = -\frac{\cos x + \sin x}{\cos x - \sin x} \quad \therefore \quad \frac{F'(a)}{f'(a)} = 1.$$

4. Required the value of

$$\frac{x + x^2 - (n+1)^2 x^{n+1} + (2n^2 + 2n - 1) x^{n+2} - n^2 x^{n+3}}{(1-x)^3}$$

when $x = 1$.

$$\frac{F'(x)}{f'(x)} =$$

$$\frac{1 + 2x - (n+1)^3 x^n + (n+2)(2n^2 + 2n - 1) x^{n+1} - n^2 (n+3) x^{n+2}}{-3(1-x)^2}$$

$$\therefore \quad \frac{F'(a)}{f'(a)} = \frac{0}{0}$$

$$\frac{F''(x)}{f''(x)} =$$

$$\frac{2-n(n+1)^3x^{n-1}+(n+1)(2n^3+6n^2+3n-2)x^n-(n+2)(n^3+3n^2)x^{n+1}}{6(1-x)}$$

$$\therefore \frac{F''(a)}{f''(a)} = \frac{0}{0}$$

$$\frac{F'''(x)}{f'''(x)} =$$

$$\frac{-(n^3-n)(n+1)^3x^{n-2}+(n^2+n)(2n^3+6n^2+3n-2)x^{n-1}-(n+1)(n+2)(n^3+3n^2)x^n}{-6}$$

$$\therefore \frac{F'''(a)}{f'''(a)} = \frac{n(n+1)(2n+1)}{6} = \frac{F(a)}{f(a)}.$$

5. Required the value of $\dfrac{a-\sqrt{a^2-x^2}}{x^2}$ when $x=0$.

$$\frac{F''(a)}{f''(a)} = \frac{1}{2a}.$$

6. Required the value of $\dfrac{x \sin x - 90^\circ}{\cos x}$ when $x = 90^\circ$.

$$\frac{F'(a)}{f'(a)} = -1.$$

7. Required the value of $\dfrac{1-x}{\cot x \dfrac{\pi}{2}}$ when $x=1$.

$$\frac{F'(a)}{f'(a)} = \frac{2}{\pi}.$$

8. Required the value of $\dfrac{a^n-x^n}{\log a - \log x}$ when $x=a$.

$$\frac{F'(a)}{f'(a)} = na^n.$$

(42.) If, in the application of the foregoing rule, we happen to arrive at a differential coefficient, which becomes infinite for the proposed value $x = a$, we must conclude that the development, in a series of ascending integral powers of h, is impossible for that particular value of the variable; and that, therefore, the rule, which is founded on the possibility of this development, becomes inapplicable. The process, however, to be adopted in such cases, is still analogous to that above; depending upon the development of the numerator and denominator of the proposed fraction; but here this development must be sought for by the common algebraical methods.

As before, let $\frac{F(x)}{f(x)}$ be a fraction which becomes $\frac{0}{0}$ when we change x into a. Substitute $a + h$ for x, and let the terms of the fraction be developed according to the increasing powers of h, either by involution, the extraction of roots, or some other algebraical process; then we shall have

$$\frac{F(a+h)}{f(a+h)} = \frac{Ah^{\alpha} + Bh^{\beta} + \&c.}{A'h^{\alpha'} + B'h^{\beta'} + \&c.},$$

in which the terms of the series are supposed to be so arranged that α and α' are the smallest exponents in each series; β and β' the next in magnitude, and so on. Now these three cases present themselves, viz.

$$1^{\circ}\ \alpha > \alpha';\ 2^{\circ}\ \alpha = \alpha';\ 3^{\circ}\ \alpha < \alpha'.$$

In the first case, by dividing the two terms of the fraction by $h^{\alpha'}$, and then supposing $h = 0$, there results

$$\frac{F(a)}{f(a)} = \frac{0}{A'} = 0.$$

In the second case the result of the same process is

$$\frac{F(a)}{f(a)} = \frac{A}{A'}$$

In the third case, by dividing the two terms of the fraction by h^{α}, and then supposing $h = 0$, the result is

$$\frac{F(a)}{f(a)} = \frac{A}{0} = \infty .$$

It appears from these results that the development of the numerator and denominator need not be carried beyond the first term, or that involving the lowest exponent of h;* and according as the exponent in the numerator is greater than, equal to, or less than, that in the denominator, will the true value of the fraction be 0, finite, or infinite. We have, therefore, the following rule:

Substitute $a + h$ for x, in the proposed fraction. Find the term containing the lowest exponent of h, in the development of the numerator; and that containing the lowest exponent of h, in the development of the denominator. If the former exponent be *greater* than this latter, the true value of the fraction will be 0; if *less*, it will be infinite. But if these exponents are *equal*, divide the coefficient of the term in the numerator, by the coefficient of that in the denominator; and the true result will be obtained.

This method, which is applicable in all cases, may frequently be employed advantageously, even where the preceding rule applies.

EXAMPLES.

9. Required the value of $\frac{(x^2 - 3ax + 2a^2)^{\frac{2}{3}}}{(x^3 - a^3)^{\frac{1}{2}}}$ when $x = a$.

Substituting $a + h$ for x, we have

$$\frac{F(a+h)}{f(a+h)} = \frac{h^{\frac{2}{3}}(h-a)^{\frac{2}{3}} \dagger}{h^{\frac{1}{2}}(3a^2 + 3ah + h^2)^{\frac{1}{2}}} = \frac{(-ah)^{\frac{2}{3}} + \&c.}{(3a^2h)^{\frac{1}{2}} + \&c.}.$$

---

* The first term which actually *appears* in the development is of course meant here: those which may vanish in consequence of the coefficients vanishing not being considered.

† To develop this according to the ascending powers of h, we must write it thus: $(-a+h)^{\frac{2}{3}}$, and apply the binomial theorem, when we shall have the series $(-a)^{\frac{2}{3}} + \frac{2}{3}a^{-\frac{1}{3}}h + \&c.$

Since the exponent of h in the numerator exceeds that in the denominator, we have

$$\frac{F(a)}{f(a)} = 0.$$

10. Required the value of $\dfrac{\sqrt{x} - \sqrt{a} + \sqrt{x-a}}{\sqrt{x^2-a^2}}$, when $x = a$,

(see p. 69.)

Substituting $a + h$ for x,

$$\frac{F(a+h)}{f(a+h)} = \frac{(a+h)^{\frac{1}{2}} - a^{\frac{1}{2}} + h^{\frac{1}{2}}}{h^{\frac{1}{2}}(2a+h)^{\frac{1}{2}}} = \frac{h^{\frac{1}{2}} + \&c.}{(2ah)^{\frac{1}{2}} + \&c.}$$

$$\therefore \frac{F(a)}{f(a)} = \frac{1}{(2a)^{\frac{1}{2}}}.$$

11. Required the value of $\dfrac{(x^2-a^2)^{\frac{3}{2}} + x - a}{(1+x-a)^3 - 1}$ when $x = a$.

Substituting $a + h$ for x,

$$\frac{F(a+h)}{f(a+h)} = \frac{h^{\frac{3}{2}}(2a+h)^{\frac{3}{2}} + h}{(1+h)^3 - 1} = \frac{h + \&c.}{3h + \&c.}$$

$$\therefore \frac{F(a)}{f(a)} = \tfrac{1}{3}.$$

12. Required the value of $\dfrac{a(4a^3+4x^3)^{\frac{1}{3}} - ax - a^2}{(2a^2+2x^2)^{\frac{1}{2}} - a - x}$ when $x = a$.

Substituting $a + h$ for x,

$$\frac{F(a+h)}{f(a+h)} = \frac{a(8a^3+12a^2h+12ah^2+4h^3)^{\frac{1}{3}} - 2a^2 - ah}{(4a^2+4ah+2h^2)^{\frac{1}{2}} - 2a - h},$$

which, by actually extracting the roots indicated,

$$= \frac{a\left(2a+h+\dfrac{h^2}{2a} + \&c. - 2a - h\right)}{2a+h+\dfrac{h^2}{4a} + \&c. - 2a - h}$$

$$= \frac{a\left(\frac{h^2}{2a} + \&c.\right)}{\frac{h^2}{4a} + \&c.} \quad \therefore \quad \frac{F(a)}{f(a)} \quad \frac{a}{\frac{1}{2}} = 2a.$$

This example is perhaps more easily performed by differentiation, according to the first rule: thus

$$\frac{F'(x)}{f'(x)} = \frac{a\,(4a^3 + 4x^3)^{-\frac{2}{3}}\,4x^2 - a}{(2a^2 + 2x^2)^{-\frac{1}{2}}\,2x - 1} \quad \therefore \quad \frac{F'(a)}{f'(a)} = \frac{0}{0}$$

$$\frac{F''(x)}{f''(x)} = \frac{-a\,(4a^3 + 4x^3)^{-\frac{5}{3}}\,32x^4 + a\,(4a^3 + 4x^3)^{-\frac{2}{3}}8x}{-(2a^2 + 2x^2)^{-\frac{3}{2}}\,4x^2 + 2\,(2a^2 + 2x^2)^{-\frac{1}{2}}}$$

$$\therefore \quad \frac{F''(a)}{f''(a)} = \frac{2}{\frac{1}{a}} = 2a.$$

(43.) Having thus seen how to determine the value of any fraction of which the numerator and denominator become each 0 for particular values of the variable, we readily perceive how the value may be found when particular substitutions make the numerator and denominator each infinite. For if $\frac{F(a)}{f(a)} = \frac{\infty}{\infty}$, then obviously

$$\frac{F(a)}{f(a)} = \frac{\frac{1}{f(a)}}{\frac{1}{F(a)}} = \frac{0}{0}.$$

So that if we find, by the preceding methods, the value of this last fraction, the value of the proposed fraction will be also obtained.

The following example will illustrate this.

13. Required the value of $\frac{\tan\left(\frac{1}{2}\,\pi\,.\,\frac{x}{a}\right)}{x^2\,a^{-1}\,(x^2 - a^2)^{-1}}$, when $x = a$.

In this case the fraction takes the form $\frac{\infty}{\infty}$, therefore,

$$\frac{\frac{1}{f(x)}}{\frac{1}{F(x)}} = \frac{a(x^2-a^2)\,x^{-2}}{\cot\left(\frac{1}{2}\pi \cdot \frac{x}{a}\right)} \therefore \frac{F'(x)}{f'(x)} = \frac{2a^3\,x^{-3}}{-\operatorname{cosec}^2\left(\frac{1}{2}\pi \cdot \frac{x}{a}\right)\frac{\pi}{2a}};$$

$$\therefore \frac{F'(a)}{f'(a)} = \frac{2}{-\frac{\pi}{2a}} = -\frac{4a}{\pi}.$$

(44.) By the same principles we may also find the true value of a *product* consisting of two factors, which for a particular value of the variable becomes, the one 0, and the other ∞. For if $F(a) = 0$ and $f(a) = \infty$, then,

$$F(a) \times f(a) = \frac{F(a)}{\frac{1}{f(a)}} = \frac{0}{0}.$$

We shall give an example of this.

11. Required the value of the product $(1 - x)$ tan. $(\frac{1}{2}\pi x)$ when $x = 1$.

In this case the first factor becomes 0 and the second ∞.

$$\frac{F(x)}{\frac{1}{f(x)}} = \frac{1-x}{\frac{1}{\tan\left(\frac{1}{2}\pi x\right)}} = \frac{1-x}{\cot\left(\frac{1}{2}\pi x\right)} \therefore F'(x) \times f'(x) = \frac{1}{\operatorname{cosec}^2\left(\frac{1}{2}\pi x\right)\frac{\pi}{2}}$$

$$\therefore F'(a) \times f'(a) = \frac{1}{\frac{1}{2}\pi} = \frac{2}{\pi}.$$

(45.) And finally, by the same principles, the true value of the difference of two functions may be ascertained, in the case where the substitution of a particular value for the variable causes each of them to become infinite.

For if $F(a) = \infty$ and $f(a) = \infty$, then

$$F(a) - f(a) = \frac{\frac{1}{f(a)} - \frac{1}{F(a)}}{\frac{1}{F(a) \times f(a)}} = \frac{0}{0}.$$

The following example belongs to this case:

15. Required the true value of the difference $x \tan x - \frac{1}{2}\pi \sec x$, when $x = 90°$.

$$\frac{\frac{1}{f(x)} - \frac{1}{F(x)}}{\frac{1}{F(x) + f(x)}} = \frac{\frac{1}{\frac{1}{2}\pi \sec x} - \frac{1}{x \tan x}}{\frac{1}{x \tan x \times \frac{1}{2}\pi \sec x}} = \frac{x \sin x - \frac{1}{2}\pi}{\cos x}$$

by substituting $\cos x$ for $\frac{1}{\sec x}$, and then multiplying numerator and denominator by $\frac{1}{2}\pi x \tan x$,

$$\therefore F'(x) - f'(x) = \frac{x \cos x + \sin x}{-\sin x} \therefore F(a) - f(a) = -1.$$

It should be remarked that in this, as well, indeed, as in the preceding cases, the transformation requisite to reduce the expression to the form $\frac{0}{0}$, in many instances at once presents itself to the mind; when of course it will be unnecessary to recur to the preceding formulas. The example just given is one of these instances, for since $\tan x = \frac{\sin x}{\cos x}$, and $\sec x = \frac{1}{\cos x}$, the proposed expression at once reduces to

$$\frac{x \sin x - \frac{1}{2}\pi}{\cos x},$$

which is the required form.

(46.) We shall terminate this chapter with a few miscellaneous examples for the exercise of the student.

16. Required the value of $\frac{x^n - 1}{x - 1}$, when $x = 1$.

Ans. n.

17. Required the value of $\dfrac{ax^2+ac^2-2acx}{bx^2-2bcx+bc^2}$ when $x=c$.

Ans. $\dfrac{a}{b}$.

18. Required the value of $\dfrac{x^3-ax^2-a^2x+a^3}{x^2-a^2}$ when $x=a$.

Ans. 0.

19. Required the value of $\dfrac{(2a^3x-x^4)^{\frac{1}{2}}-a\,(a^2x)^{\frac{1}{2}}}{a-(ax^3)^{\frac{1}{4}}}$, when $x=a$.

Ans. $\dfrac{16a}{9}$.

20. Required the value of $\dfrac{1-x+\log x}{1-(2x-x^2)^{\frac{1}{2}}}$, when $x=1$.

Ans. -1.

21. Required the value of $\dfrac{x^x-x}{1-x+\log x}$, when $x=1$.

Ans. -2.

22. Required the value of $\dfrac{\tan x-\sin x}{\sin x^3}$, when $x=0$.

Ans. $\frac{1}{2}$.

23. Required the value of $\dfrac{x\log x-(x-1)}{(x-1)\log x}$, when $x=1$.

Ans. $\frac{1}{2}$.

24. Required the value of $\dfrac{(x^2-a^2)^{\frac{3}{2}}}{(x-a)^{\frac{3}{2}}}$, when $x=a$.

Ans. $(2a)^{\frac{1}{2}}$.

25. Required the value of $\dfrac{x}{x-1}-\dfrac{1}{\log x}$, when $x=1$.

Ans. $\frac{1}{2}$.

26. Required the value of $\dfrac{ax-x^2}{a^4-2a^3x+2ax^3-x^4}$, when $x=a$.

Ans. ∞.

27. Required the value of $\frac{1}{\log x} - \frac{x}{\log x}$, when $x = 1$.

Ans. -1.

28. Required the value of $\frac{x^2 - a^2}{x^2} \tan \frac{\pi x}{2a}$, when $x = a$.

Ans. $-\frac{4}{\pi}$.

29. Required the value of $\frac{a(1-x)}{\cot \frac{1}{2}\pi x}$, when $x = 1$.

Ans. $\frac{2a}{\pi}$.

30. Required the value of $\frac{e^x - e^{\sin x}}{x - \sin x}$, when $x = 0$.

Ans. 1.

CHAPTER VI.

ON THE MAXIMA AND MINIMA VALUES OF FUNCTIONS OF A SINGLE VARIABLE.

(47.) In any function $y = F(x)$ let the independent variable take a particular value $x = a$, as also a preceding and succeeding value $x = a - h$, and $x = a + h$; then the corresponding values of the function, arranged according to those of the variable, will be

$$F(a-h),\ F(a),\ F(a+h);$$

and if a be such that for any finite value of h, however small, and for all intermediate values between this and 0, the middle value, $F(a)$ *exceeds* that on each side, the value $x = a$ is said to render the function a *maximum;* but if the middle value continue *less* than that on each side, between the same limits of h, the value $x = a$ is said to render the

function a *minimum;* so that we are not always to understand by the expression, *maximum* value of a function, the *greatest* value such function can possibly take, the term being of more comprehensive meaning, applying to *every* state of the function which exceeds its immediately preceding and succeeding state. In like manner, the *minimum* value of a function does not always imply the *least* possible value of such function, but equally characterizes every state of the function which is less than its immediately preceding and succeeding state.

Before proceeding to the general method of determining the values of x, necessary to render any proposed function a maximum or a minimum, we must premise this lemma:

If the function $F(a + h)$ *be developed according to the ascending powers of* h, *a value so small may be given to* h *that any proposed term in the series shall exceed the sum of all that follow.*

(48.) Let Ah^{α} be any proposed term in the development, and let Bh^{β}, Ch^{γ}, &c. be those which follow, each exponent being greater than that which precedes it. We are to prove that h may be taken so small that

$$Ah^{\alpha} > h^{\beta}\,(B + Ch^{\gamma-\beta} + Dh^{\delta-\beta} + \&c.)$$

Putting S for the sum of the series within the parentheses, it is obvious that h may be taken so small that $Sh^{\beta-\alpha}$ may be less than any proposed quantity, A; and that therefore, if h' be such a value, we must have

$$Ah'^{\alpha} > Sh'^{\beta}$$

which establishes the proposition. As $Sh^{\beta-\alpha}$ is less than A, for $h = h'$, the expression continues less than A for every value of h less than h'.

(49.) Let us now enquire by what means we may determine those values of x which render any proposed function, $y = F(x)$, a maximum or a minimum. In order to do this, let x be changed into $x \pm h$, then, by Taylor's theorem,

$$F(x \pm h) = F(x) \pm \frac{dy}{dx} h + \frac{d^2y}{dx^2} \frac{h^2}{1 \cdot 2} \pm \frac{d^3y}{dx^3} \frac{h^3}{1 \cdot 2 \cdot 3} +$$

$$\frac{d^4y}{dx^4} \frac{h^4}{1 \cdot 2 \cdot 3 \cdot 4} \pm \&c.$$

Now if $x = a$ render the proposed function a *maximum*, then there exists for h some finite value h', such that for all the intermediate values between this and 0 we must have

$$F(a) > F(a \pm h),$$

and, consequently, for every such value that can occur among those for which Taylor's theorem does not fail, we must have the condition

$$\pm \left[\frac{dy}{dx}\right] h + \left[\frac{d^2y}{dx^2}\right] \frac{h^2}{1 \cdot 2} \pm \left[\frac{d^3y}{dx^3}\right] \frac{h^3}{1 \cdot 2 \cdot 3} + \&c. < 0 \ldots (1).$$

But if the supposed value render the function a *minimum*, then, for all the intermediate values of h between $h = h'$ and $h = 0$, we must have

$$F(a) < F(a \pm h),$$

and, consequently,

$$\pm \left[\frac{dy}{dx}\right] h + \left[\frac{d^2y}{dx^2}\right] \frac{h^2}{1 \cdot 2} \pm \left[\frac{d^3y}{dx^3}\right] \frac{h^3}{1 \cdot 2 \cdot 3} + \&c. > 0 \ldots (2).$$

It has, however, been proved above, that a value may be given to h small enough to render the first term in each of the series (1) and (2) greater than the sum of all the other terms; and that this first term will continue greater for all other values of h between this small value and 0; so that, for each of these values of h, the sign belonging to the sum of the whole series is the same as that of the first term; it is impossible, therefore, that either of the conditions (1) or (2) can exist for *both* $+\left[\frac{dy}{dx}\right] h$ and $-\left[\frac{dy}{dx}\right] h$, unless $\left[\frac{dy}{dx}\right] = 0$; we conclude, therefore, that of the values of x, for which Taylor's theorem holds, those only can render the function a maximum or minimum which fulfil the condition,

$$\frac{dy}{dx} = 0;$$

and even these must be tested as follows: Expunge the first term from each of the series, (1), (2), then in virtue of the condition just stated, we must have, in the case of a maximum,

$$[\frac{d^2y}{dx^2}]\ \frac{h^2}{1\cdot 2} \pm [\frac{d^3y}{dx^3}]\ \frac{h^3}{1\cdot 2\cdot 3} + \&c. < 0 \quad . \quad . \quad . \quad (3),$$

and in the case of a minimum,

$$[\frac{d^2y}{dx^2}]\ \frac{h^2}{1\cdot 2} \pm [\frac{d^3y}{dx^3}]\ \frac{h^3}{1\cdot 2\cdot 3} + \&c. > 0 \quad . \quad . \quad . \quad (4).$$

But the former of these conditions cannot exist for any of the values of h between $h = h'$ and $h = 0$, by virtue of the foregoing principle, unless $[\frac{d^2y}{dx^2}]$ is negative; nor can the latter condition exist unless $[\frac{d^2y}{dx^2}]$ is positive; that is, supposing that these coefficients do not vanish from the series (3) and (4).

We may infer, therefore, that of the values of x which satisfy the condition $\frac{dy}{dx} = 0$, those only among them that also satisfy the condition $\frac{d^2y}{dx^2} < 0$ belong to maximum values of the function, while those fulfilling the condition $\frac{d^2y}{dx^2} > 0$ belong to minimum values of the function.

It is possible, however, that some of the values derived from the equation $\frac{dy}{dx} = 0$ may, when substituted for x in $\frac{d^2y}{dx^2}$, cause this coefficient to vanish, in which case the conditions (1), (2), become

$$\pm [\frac{d^3y}{dx^3}]\ \frac{h^3}{1\cdot 2\cdot 3} + [\frac{d^4y}{dx^4}]\ \frac{h^4}{1\cdot 2\cdot 3\cdot 4} \pm \&c. < 0 \quad . \quad . \quad . \quad (5),$$

and

$$\pm [\frac{d^3y}{dx^3}]\ \frac{h^3}{1\cdot 2\cdot 3} + [\frac{d^4y}{dx^4}]\ \frac{h^4}{1\cdot 2\cdot 3\cdot 4} \pm \&c. > 0 \quad . \quad . \quad . \quad (6),$$

which are both impossible unless $[\frac{d^3y}{dx^3}] = 0$, for reasons similar to those assigned above, and, unless, also $[\frac{d^4y}{dx^4}] < 0$ in the case of a maximum, and $[\frac{d^4y}{dx^4}] > 0$ in the case of a minimum; that is, on the supposition that this coefficient does not vanish from the series (5) and (6.) If, however, this coefficient does vanish, then, for reasons similar to those assigned in the preceding cases, the following coefficient $\frac{d^5y}{dx^5}$ must also vanish; the condition of maximum will then be $[\frac{d^6y}{dx^6}] < 0$, and the condition of minimum $[\frac{d^6y}{dx^6}] > 0$, and so on.

It hence appears, that to determine what values of x correspond to the maxima and minima values of the function $y = F(x)$, we must proceed as follows:

Determine the real roots of the equation $\frac{dy}{dx} = 0$, and substitute them, one by one, in the following coefficients $\frac{d^2y}{dx^2}$, $\frac{d^3y}{dx^3}$, &c. stopping at the first, which does not vanish. If this is of an *odd* order, the root that we have employed is *not* one of those values of x that renders the function either a maximum or a minimum; but if it is of an *even* order, then, according as it is *negative* or *positive*, will the root employed correspond to a *maximum* or to a *minimum* value of the function.

(50.) It must, however, be remarked, that should any of the roots of the equation $\frac{dy}{dx} = 0$ cause the first of the following coefficients, which does not vanish, to become infinite, we cannot apply to such roots the foregoing tests for detecting the admissible values, and for distinguishing the maxima from the minima, because the true development of the

function for any such value of x begins to differ in form from Taylor's development, at that term which is thus rendered infinite (4); so that we cannot infer, from Taylor's series, whether the power of h, which ought to enter this term, is odd or even.

In a case of this kind, therefore, we must find, by actual involution, extraction, &c. the true term that ought to supply the place of that rendered infinite in Taylor's series for $x=a$. If this term take an odd power of h, or, rather, if its sign change with the sign of h, then $x=a$ does not render the function either a maximum or a minimum; but if the sign does not change with that of h, then the value of x renders the function a maximum or a minimum, according as the sign of this term is negative or positive.

To illustrate this case, suppose the function were

$$y=b+(x-a)^{\frac{5}{3}}$$

$$\therefore \frac{dy}{dx}=\frac{5}{3}(x-a)^{\frac{2}{3}}$$

$$\frac{d^2y}{dx^2}=\frac{10}{9}(x-a)^{-\frac{1}{3}}.$$

Now the equation $\frac{dy}{dx}=0$ gives $x=a$, so that if any value of x could render the proposed function a maximum or a minimum, this probably would be it. We cannot, however, without examination, be confident on this point, because, although suitable values of x may exist, yet they may be such as to cause Taylor's theorem to fail even at its commencement, and consequently cannot be among the roots of the equation $\frac{dy}{dx}=0$. In order however to test the root just found, substitute its value in $\frac{d^2y}{dx^2}$; the result we see is infinite, and we cannot therefore infer the state of the function from this coefficient; substituting, then, $a\pm h$ for x in the proposed, we have

$$F(a\pm h)=b\pm h^{\frac{5}{3}};$$

and, as $h^{\frac{5}{3}}$ obviously changes its sign when h does, we conclude that the function proposed becomes neither a maximum nor a minimum for the proposed value of x.

Again, let

$$y = b + (x - a)^{\frac{5}{3}}$$

$$\therefore \frac{dy}{dx} = \frac{4}{3}(x - a)^{\frac{1}{3}}$$

$$\frac{d^2y}{dx^2} = \frac{4}{9}(x - a)^{-\frac{2}{3}}.$$

The equation $\frac{dy}{dx} = 0$ gives $x = a$, a value which causes $\frac{d^2y}{dx^2}$ to become infinite: therefore, substituting $a \pm h$ for x in the proposed, we have

$$F(a \pm h) = b + h^{\frac{5}{3}};$$

and, as the sign of $h^{\frac{5}{3}}$ is positive whatever be the sign of h, we conclude that the value $x = a$ renders the function a minimum.

(51.) There remains to be considered one more case to which the general rule is not applicable, and which, like the preceding, arises from the failure of Taylor's theorem. We have hitherto examined only those values of x for which Taylor's development is possible, as far at least as the first power of h, but, as before remarked, we cannot say that among those values of x, which would render the coefficient of this first power *infinite*, there may not be some which cause the function to fulfil the conditions of maxima or minima; therefore, before we can conclude in any case that the values of x, deduced from the condition $\frac{dy}{dx} = 0$, comprise among them *all* those which can render the function a maximum or minimum, we must examine those values arising from the condition $\frac{dy}{dx} = \infty$, by substituting each of these $\pm h$ for x in the proposed equation, and observing which of the results agree with the conditions of maxima and minima in (47.) It will be frequently

found, however, that this new condition can only be satisfied by $x = \infty$, which is of course not admissible inasmuch as it does not admit of preceding and *succeeding* values. In the two examples above, for instance, this new condition furnishes no admissible value for x, so that, in the first example, no suitable value exists, and in the second only one, and that renders the function a minimum.

(52.) If the function that y is of x be *implicitly* given, that is, if

$$u = F(x, y) = 0;$$

then, by (39), we have, for the differential coefficient,

$$\frac{dy}{dx} = -\frac{du}{dx} \div \frac{du}{dy} \quad \ldots . \quad (1),$$

and therefore, when $\frac{dy}{dx} = 0$, we must have $\frac{du}{dx} = 0$; hence, the values corresponding to maxima and minima are determinable from the two equations*

$$\left.\begin{aligned} u &= 0 \\ \frac{du}{dx} &= 0 \end{aligned}\right\} \quad \ldots . \quad (2).$$

Having found from these the values of x that *may* render y a maximum or a minimum,† as also the corresponding values of y itself, we must substitute them for x and y in $\frac{d^2y}{dx^2}$, when those values of y will be

---

* Other values may be implied in the condition $\frac{dy}{dx} = \infty$, which leads to $\frac{du}{dy} = 0$, but to ascertain which of these are applicable would require us to solve the equation for y.

† Garnier, at p. 271 of his *Calcul Differentiel*, says, that, by means of the equations (2), "on obtient les valeurs de x et y par lesquelles $F(x, y)$ devient ou peut devenir maximum ou minimum;" but this is evidently a mistake, since, by hypothesis, $F(x, y)$ is always $= 0$.

maxima that render this coefficient negative, and those will be minima that render it positive. But those values that cause it to vanish, belong neither to maxima nor to minima, unless the same values cause also $\frac{d^3y}{dx^3}$ to vanish, and so on.

The second differential coefficient may be readily derived from (1); for, putting for brevity

$$\frac{dy}{dx} = -\frac{M}{N},$$

we have

$$\frac{d^2y}{dx^2} = -\frac{N\left(\frac{dM}{dx} + \frac{dM}{dy}\frac{dy}{dx}\right) - M\left(\frac{dN}{dx} + \frac{dN}{dy}\frac{dy}{dx}\right)}{N^2},$$

which, because $\frac{M}{N}$, or $\frac{dy}{dx}$, $= 0$, becomes for the particular values of x resulting from this condition,

$$\left[\frac{d^2y}{dx^2}\right] = -\left[\frac{d^2u}{dx^2}\right] \div \left[\frac{du}{dy}\right] \quad \ldots \quad (3).$$

By differentiating the above expression for $\frac{d^2y}{dx^2}$ we should find

$$\left[\frac{d^3y}{dx^3}\right] = -\left[\frac{d^3u}{dx^3}\right] \div \left[\frac{du}{dy}\right] \quad \ldots \quad (4),$$

and so on.

(53.) Before we proceed to apply the foregoing theory to examples, we shall state a few particulars that may, in many instances, be serviceable in abridging the process of finding maxima and minima.

1. If the proposed function appears with a constant factor, such factor may be omitted. Thus, calling the function Ay, the first differential coefficient will be $A\frac{dy}{dx}$, and $A\frac{dy}{dx} = 0$ leads to $\frac{dy}{dx} = 0$, also $\frac{1}{A\frac{dy}{dx}} = 0$ leads to $\frac{1}{\frac{dy}{dx}} = 0$, so that A may be expunged from the function.

2. Whatever value of x renders a function a maximum or minimum, the same value must obviously render its square, cube, and every other power, a maximum or minimum; so that when a proposed function is under a radical, this may be removed. The rational function may, however, become a maximum or a minimum for more values of x than the original root; indeed, all values of x which render the rational function *negative* will render every even root of it imaginary; such values, therefore, do not belong to that root; moreover, if the rational function be $= 0$, when a maximum, the corresponding value of the variable will be inadmissible in any even root, because the contiguous values of the function must be negative.

3. Whatever value of x renders a function a maximum, the same value will obviously render its reciprocal a minimum, and vice versâ. Hence if it be found more convenient to take the reciprocal of any proposed function, instead of the function itself, we are at liberty to do so, recollecting, however, that the maxima and minima values of the changed function correspond to the minima and maxima of the original one.

EXAMPLES.

(54.) 1. To determine for what values of x the function

$$y = a^4 + b^3x - c^2x^2$$

becomes a maximum or minimum,

$$\frac{dy}{dx} = b^3 - 2c^2x, \quad \frac{d^2y}{dx^2} = -2c^2.$$

From the second equation it appears that, whatever be the values of x, given by the condition $\frac{dy}{dx} = 0$, they must all belong to maxima.

From $b^3 - 2c^2x = 0$ we get $x = \frac{b^3}{2c^2}$; hence

$$\text{when } x = \frac{b^3}{2c^2} \therefore y = a^4 + \frac{b^6}{4c^2}, \text{ a maximum.}$$

The equation $\frac{dy}{dx} = \infty$ would give, in the present case, $x = \infty$, a value which is inadmissible (51).

2. To determine the maxima and minima values of the function

$$y = 3a^2x^3 - b^4x + c^5$$

$$\frac{dy}{dx} = 9a^2x^2 - b^4, \quad \frac{d^2y}{dx^2} = 18a^2x$$

putting

$$9a^2x^2 - b^4 = 0 \quad \therefore \; x = \pm \frac{b^2}{3a}.$$

Substituting each of these values in $\frac{d^2y}{dx^2}$, we infer from the results that

$$\text{when } x = \frac{b^2}{3a} \; \ldots \; y = c^5 - \frac{2b^6}{9a}, \text{ a minimum,}$$

$$x = -\frac{b^2}{3a} \; \ldots \; y = c^5 + \frac{2b^6}{9a}, \text{ a maximum.}$$

3. To determine the maxima and minima values of the function

$$y = \sqrt{2ax}.$$

Omitting the radical

$$u = 2ax \; \therefore \; \frac{du}{dx} = 2a,$$

as this can never become 0 or ∞, we infer that the function has no maximum or minimum value.

4. To determine the maximum and minimum values of the function

$$y = \sqrt{4a^2x^2 - 2ax^3}.$$

Omitting the radical and the constant factor $2a$ (53),

$$u = 2ax^2 - x^3,$$

$$\therefore \; \frac{du}{dx} = 4ax - 3x^2, \; \frac{d^2u}{dx^2} = 4a - 6x,$$

$$\therefore x(4a - 3x) = 0 \ \therefore x = 0, \text{ or } x = \frac{4a}{3}.$$

Substituting each of these values in $\frac{d^2u}{dx^2}$, the results are $4a$ and $-4a$; hence

when $x = 0$ . . . $y = 0$, a minimum,

$$x = \frac{4a}{3} \ . \ . \ y = \frac{8\sqrt{3}}{9}a^2, \text{ a maximum.}$$

If, instead of the preceding, the example had been

$$y = \sqrt{2ax^3 - 4a^2x^2},$$

we should have had

$$\frac{du}{dx} = 3x^2 - 4ax, \ \frac{d^2u}{dx^2} = 6x - 4a;$$

$$\therefore x(3x - 4a) = 0, \ \therefore x = 0, \text{ or } x = \frac{4a}{3},$$

the same values as before; but the first corresponds here to a *maximum*, since it makes $\frac{d^2u}{dx^2}$ negative; this value, therefore, must, by (53), be rejected. If, indeed, we substitute $0 \pm h$ for x, in the proposed function, it becomes

$$y = \sqrt{-4a^2h^2 \mp 2ah^3},$$

where h may be taken so small as to cause the expression under the radical to be negative for all values of h between this and 0.

5. To determine the maxima and minima values of the function

$$y = a + \sqrt[3]{a^3 - 2a^2x + ax^2}.$$

If y is a maximum or minimum, $y - a$ will be so; therefore, transposing the a, and omitting the radical (53),

$$u = a^3 - 2a^2x + ax^2$$

$$\frac{du}{dx} = -2a^2 + 2ax, \ \frac{d^2u}{dx^2} = 2a,$$

$$\therefore -2a^2 + 2ax = 0 \ \therefore x = a,$$

$$\therefore \text{when } x = a \ . \ . \ . \ y = a, \text{ a minimum.}$$

6. To determine the maxima and minima values of the function

$$y = \frac{a^2 x}{(a - x)^2}.$$

In solving this example we shall employ the third principle at p. 68, which is often found useful, when the proposed function is a fraction with a denominator more complex than the numerator. Instead of the function itself, we shall, therefore, take its reciprocal, which will give us a more simple form, and it is plain that the maxima and minima values of the reciprocal of a function correspond respectively to the minima and maxima values of the function itself. Omitting, then, the constant a^2, and, taking the reciprocal, we have

$$u = \frac{a^2 - 2ax + x^2}{x} = \frac{a^2}{x} - 2a + x.$$

$$\therefore \frac{du}{dx} = -\frac{a^2}{x^2} + 1, \ \frac{d^2u}{dx^2} = \frac{2a^2}{x^3}.$$

$$\therefore -\frac{a^2}{x^2} + 1 = 0 \ \therefore x = \pm a \ \therefore \left[\frac{d^2u}{dx^2}\right] = \pm \frac{2}{a};$$

hence $x = a$ makes u a minimum, and $x = -a$ makes it a maximum; therefore,

$$\text{when } x = a \ . \ . \ . \ y = \infty, \text{ a maximum,}$$

$$x = -a \ . \ . \ . \ y = -\tfrac{1}{4}a, \text{ a minimum.}$$

7. To determine the maxima and minima values of the function

$$y = b + \sqrt[3]{(x - a)^5}.$$

Omitting b and the radical, we have

$$u = (x - a)^5;$$

$$\therefore \frac{du}{dx} = 5\,(x - a)^4, \ \frac{d^2u}{dx^2} = 4 \ . \ 5\,(x - a)^3;$$

$$\therefore\ 5\,(x-a)^4=0,\ \therefore\ x=a,\ \therefore\ [\frac{d^2u}{dx^2}]=0.$$

As this coefficient vanishes, we must proceed to the following, which however all contain $x-a$, and therefore vanish, till we come to $\frac{d^5u}{dx^5}$ $=2\cdot3\cdot4\cdot5$;* as therefore the first coefficient which does not vanish is of an odd order, the function does not admit of a maximum or a minimum.

8. To determine the maxima and minima values of the function

$$y=x^x$$

$$\frac{dy}{dx}=x^x\,(1+\log x),\ \frac{d^2y}{dx^2}=x^x\,\{\frac{1}{x}+(1+\log x)^2\}.$$

The factor x^x can never become 0, therefore,

$$(1+\log x)=0,\ \therefore\ \log x=-1;$$

$$\therefore\ x=e^{-1}=\frac{1}{e};$$

$$\therefore\ [\frac{d^2y}{dx^2}]=(\frac{1}{e})^{\frac{1}{e}}\cdot e;$$

$$\therefore\ \text{when } x=\frac{1}{e},\ x^x=(\frac{1}{e})^{\frac{1}{e}},\ \text{a minimum.}$$

9. To determine the maxima and minima values of y in the function

$$u=x^3-3axy+y^3=0$$

$$\frac{du}{dx}=3x^2-3ay,\ \therefore\ (52)$$

---

* All the coefficients preceding the nth will always obviously vanish from the function $P\,(x\pm a)^n$; so that if n be odd, $x=\mp a$ renders the function neither a maximum nor a minimum. But when n is even, then a maximum or minimum obviously exists.

$$\left.\begin{aligned} x^3 - 3axy + y^3 = 0 \\ 3x^2 - 3ay = 0 \end{aligned}\right\} \therefore y = \frac{x^2}{a} \therefore x^6 - 2a^3x^3 = 0,$$

$$\therefore x = 0, \text{ or } x = a\sqrt[3]{2} \therefore (52)$$

$$\left[\frac{d^2y}{dx^2}\right] = -\left[\frac{d^2u}{dx^2}\right] \div \left[\frac{du}{dy}\right] = -[6x] \div \left[3\,\frac{x^4}{a^2} - 3ax\right] = -\left[\frac{2a^2}{x^3 - a^3}\right]$$

$$= \frac{2}{a} \text{ or } -\frac{2}{a},$$

$\therefore$ when $x = 0$ . . . . $y = 0$, a minimum,

$x = a\sqrt[3]{2}$ . . . . $y = a\sqrt[3]{4}$, a maximum.

10. To divide a given number a into two parts, such that the product of the mth power of the one and the nth power of the other shall be the greatest possible.

Let x be one part, then $a - x$ is the other, and

$$y = x^m (a - x)^n = \text{maximum},$$

$$\therefore \frac{dy}{dx} = mx^{m-1} (a - x)^n - nx^m (a - x)^{n-1}$$

$$= x^{m-1} (a - x)^{n-1} \{ma - (m + n) x\} = 0,$$

$$\therefore x = 0, \text{ or } a - x = 0, \text{ or}$$

$$ma - (m + n) x = 0,$$

which give

$$x = 0,\ x = a,\ x = \frac{ma}{m + n}.$$

The first and second of these values are inadmissible, because the number is not *divided* when $x = 0$, or when $x = a$.

Substituting the third value in

$$\frac{d^2y}{dx^2} = x^{m-2} (a - x)^{n-2} \{(ma - (m + n) x)^2 - m (a - x)^2 - nx^2\},$$

we have

$$[\frac{d^2y}{dx^2}] = -[x]^{m-2}[a-x]^{n-2}\{m[a-x]^2 + n[x]^2\}$$

which is negative because each factor is positive; hence the two required parts are

$$\frac{ma}{m+n} \text{ and } \frac{na}{m+n}, \text{ being to each other as } m \text{ to } n.$$

Cor. If $m = n$ the parts must be equal.

An easier solution to this problem may be obtained as follows:

Put $\frac{m}{n} = p$ and determine x, so that we may have

$$u = x^p(a-x) = \text{a maximum},$$

$$\therefore \frac{du}{dx} = px^{p-1}(a-x) - x^p$$

$$= x^{p-1}\{pa - (p+1)x\} = 0,$$

$$\therefore x = 0, \text{ or } pa - (p+1)x = 0 \ \therefore x = \frac{pa}{p+1}.$$

This last value substituted in

$$\frac{d^2u}{dx^2} = x^{p-2}\{pa - (p+1)x\} - (p+1)x^{p-1}$$

causes the first term to vanish; the result is therefore negative, so that $x = \frac{pa}{p+1} = \frac{ma}{m+n}$ corresponds to a maximum value of u, and therefore (53) to a maximum value of $u^n = x^m(a-x)^n$.

Another easy mode of solution is had by using logarithms; for it is obvious that since the logarithm of any number increases with the number, when this number is the greatest possible, its logarithm will be so too.

$$\therefore m\log x + n\log(a-x) = \text{maximum},$$

$$\therefore \frac{m}{x} - \frac{n}{a-x} = 0 \ \therefore ma - (m+n)x = 0,$$

$$\therefore x = \frac{ma}{m+n},$$

as before.

The expression for the second differential coefficient is $-(m+n)$ showing that the foregoing value of x renders the logarithmic expression a maximum.

11. To divide a number a into so many parts that their continued product may be the greatest possible.

It is obvious, from the corollary to the last example, that the parts must be equal; for the product of any two unequal parts of a number is less than that of equal parts.

Let x be the number of factors, then,

$$\left(\frac{a}{x}\right)^x = \text{a maximum},$$

$$\therefore \log\left(\frac{a}{x}\right)^x = x \log\left(\frac{a}{x}\right) = \text{a maximum},$$

$$\therefore \log\frac{a}{x} - 1 = 0,$$

$$\therefore \frac{a}{x} = \log^{-1} 1 = e \therefore x = \frac{a}{e}.^*$$

hence the proposed number must be divided by the number $e = 2{\cdot}718281828$.

12. To determine those conjugate diameters of an ellipse which include the greatest angle.

Call the principal semi-diameters of the ellipse a, b, the sought semi-conjugates x, x', and the sine of the angle they include y. Then (*Anal. Geom.* p. 78,)

$$x^2 + x'^2 = a^2 + b^2 \therefore x' = \sqrt{a^2 + b^2 - x^2}$$

$$xx'y = ab \therefore y = \frac{ab}{xx'}$$

---

* There is obviously no necessity to recur to the second differential coefficient to ascertain whether this value render the function a maximum or a minimum, since it is plain that there is no minimum unless one of the parts may be 0.

$$\therefore y = \frac{ab}{x\sqrt{a^2 + b^2 - x^2}}.$$

Omitting the constant ab, inverting the function (Ex. 6,) and squaring, we have

$$u = a^2x^2 + b^2x^2 - x^4,$$

$$\therefore \frac{du}{dx} = 2a^2x + 2b^2x - 4x^3 = 0,$$

$$\therefore x = 0,\ a^2 + b^2 - 2x^2 = 0\ \therefore x^2 = \frac{a^2 + b^2}{2} = \frac{x^2 + x'^2}{2}.$$

The first of these values is inadmissible; from the second we find that

$$x^2 = x'^2$$

hence the conjugates are equal. For the second differential coefficient we have

$$\frac{d^2u}{dx^2} = 2a^2 + 2b^2 - 12x^2,$$

$$\therefore \left[\frac{d^2u}{dx^2}\right] = -4(a^2 + b^2).$$

This being negative, shows that $x = \sqrt{\frac{a^2 + b^2}{2}}$ corresponds to a maximum value of u, or to a minimum value of y; so that the conjugates here determined, include an angle whose *sine* is the least possible; and this happens when the angle itself is the greatest possible (being obtuse), as well as when it is the least possible.

13. To divide an angle θ into two parts, such that the product of the nth power of the sine of one part, by the mth power of the sine of the other part, may be the greatest possible.

Let x be one part, then $\theta - x$ is the other, and

$$\sin^n x \cdot \sin^m(\theta - x) = \text{maximum},$$

$$\therefore n \log \sin x + m \log \sin(\theta - x) = \text{maximum},$$

$$\therefore \frac{n \cos x}{\sin x} - \frac{m \cos(\theta - x)}{\sin(\theta - x)} = 0,$$

$$\therefore n \tan(\theta - x) = m \tan x,$$

$$\therefore n : m :: \tan x : \tan(\theta - x),$$

$$\therefore n + m : n - m :: \tan x + \tan(\theta - x) : \tan x - \tan(\theta - x),$$

$$:: \sin\theta : \sin(2x - \theta),^*$$

$$\therefore \sin(2x - \theta) = \frac{n - m}{n + m} \sin\theta$$

$$\therefore 2x - \theta = \sin^{-1}\left\{\frac{n - m}{n + m} \sin\theta\right\},$$

$$\therefore x = \frac{\theta}{2} + \frac{1}{2} \sin^{-1}\left\{\frac{n - m}{n + m} \sin\theta\right\}.$$

14. Given the hypothenuse of a right-angled triangle to determine the other sides, when the surface is the greatest possible.

Call the hypothenuse a, and one of the sides x, then the other will be $\sqrt{a^2 - x^2}$ and the area of the triangle will be

$$\frac{x}{2}\sqrt{a^2 - x^2} = \text{maximum},$$

$$\therefore u = a^2x^2 - x^4 = \text{maximum},$$

$$\therefore \frac{du}{dx} = 2a^2x - 4x^3 = 0 \quad \therefore x = 0 \text{ or } x = \frac{a}{\sqrt{2}}.$$

Substituting the second value in

$$\frac{d^2u}{dx^2} = 2a^2 - 12x^2,$$

the result being negative, shows that the above value of x corresponds to a maximum. Therefore the required sides are each $\frac{a}{\sqrt{2}}$.

15. To determine the maxima and minima values of the function

$$y = x^3 - 18x^2 + 96x - 20,$$

---

* Trigonometry, p. 48, Form 6.

when $x = 4 \ldots . y = 140$ a maximum,

$x = 8 \ldots . y = 108$ a minimum.

16. To determine a number x, such that the xth root may be the greatest possible.

Ans. $x = e = 2{\cdot}71828 \ldots .$

17. What fraction is that which exceeds its mth power by the greatest possible number?

Ans. $\sqrt[m-1]{\frac{1}{m}}$.

18. Given the equation

$$y^2 - 2mxy + x^2 = a^2$$

to determine the maxima and minima values of y.

When $x = \frac{ma}{\sqrt{1-m^2}} \ldots . y = \frac{a}{\sqrt{1-m^2}}$, a maximum,

$x = \frac{-ma}{\sqrt{1-m^2}} \ldots . y = \frac{-a}{\sqrt{1-m^2}}$, a minimum.

19. Given the position of a point between the sides of a given angle to draw through it a line so that the triangle formed may be the least possible.

Ans. The line must be bisected by the point.

20. The equation of a certain curve is $a^2y = ax^2 - x^3$, required its greatest and least ordinates.

When $x = \frac{2}{3}a \ldots . y =$ maximum,

$x = 0 \ldots . y =$ minimum.

21. To divide a given angle θ, less than 90°, into two parts, x and $\theta - x$, such that $\tan^n x \,.\, \tan^m (\theta - x)$ may be the greatest possible.

$$\tan (2x - \theta) = \frac{n-m}{n+m} \tan \theta,$$

$$\therefore 2x - \theta = \sin^{-1} \left\{ \frac{n-m}{n+m} \tan \theta \right\},$$

$$\therefore x = \frac{\theta}{2} + \frac{1}{2}\sin^{-1}\{\frac{n-m}{n+m}\tan\theta\}.$$

22. To determine the greatest parabola that can be formed by cutting a given right cone.*

SCHOLIUM.

(55.) It will be proper, before terminating the present chapter, to apprize the student that in the application of the theory of maxima and minima to geometrical enquiries, care must be taken that we do not adopt results inconsistent with the geometrical restrictions of the problem. We know, indeed, from the first principles of Analytical Geometry, that when the geometrical conditions of a problem are translated into an algebraical formula, that formula is not necessarily restricted to those conditions, but, in addition to all the possible solutions of the problem, may also furnish others that belong merely to the analytical expression, and have no geometrical signification.† If, therefore, among these latter solutions there be any belonging to maxima or minima, they are inadmissible in the application of this theory to Geometry. The following example is given by *Simpson*, at Art. 47 of his *Treatise on Fluxions*, to illustrate this.

From the extremity C of the minor axis of an ellipse to draw the longest line to the curve. Suppose F to be the point to which the line must be drawn, and call the abscissa CE, x, then the geometrical restrictions of this variable are such that its values must always lie between the limits $x = 0$, and $x = 2a$, a and b denoting the semi-axes.



By the equation of the curve

$$EF^2 = y^2 = \frac{a^2}{b^2}(2bx - x^2),$$

* A parabola is equal in surface to $\frac{2}{3}$ of a rectangle of the same base and altitude.

† See the *Analytical Geometry*, pp. 3—4.

$$\therefore \mathrm{CF}^2 = u = x^2 + \frac{a^2}{b^2}(2bx - x^2) = \text{maximum},$$

$$\therefore \frac{\mathrm{d}u}{\mathrm{d}x} = 2\left(x + \frac{a^2}{b} - \frac{a^2}{b^2}x\right) = 0,$$

$$\therefore x = \frac{a^2 b}{a^2 - b^2},$$

and since $\frac{\mathrm{d}^2 u}{\mathrm{d}x^2} = 2\left(1 - \frac{a^2}{b^2}\right)$ it follows that the foregoing expression for x renders u a maximum for all values of b less than a, and a minimum for all values of b greater than a. Hence if the relation between a and b be such that $\frac{a^2 b}{a^2 - b^2}$ may exceed $2b$, the analytical expression for CF will admit of a maximum value, although such value, not coming within the geometrical restrictions of the problem, is inadmissible. If the relation between a and b be such that $\frac{a^2 b}{a^2 - b^2} = 2b$, that is, if $a^2 = 2b^2$, the solution will be valid; and in the ellipse whose axes are thus related, CD will be the longest line that can be drawn from C, agreeably to the analytical determination, and the solution will always be valid if the axes of the ellipse are related so that $\frac{a^2 b}{a^2 - b^2}$ is not greater than $2b$, which leads to the condition $2b^2$ not greater than a^2.

CHAPTER VII.

ON THE DIFFERENTIATION AND DEVELOPMENT OF FUNCTIONS OF TWO INDEPENDENT VARIABLES.

Differentiation of functions of two independent variables.

(56.) Let $z = \mathrm{F}(x, y)$ be a function of two independent variables; then, since in consequence of this independence, however either be supposed to vary, the other will remain unchanged: the function ought to furnish two differential coefficients; the one arising from ascribing a variation to x, and the other from ascribing a variation to y, y entering the first coefficient as if it were a constant, and x entering the second as if it were a constant. The differential coefficient arising from the variation of x is expressed thus, $\frac{\mathrm{d}z}{\mathrm{d}x}$; and that arising from the variation of y thus, $\frac{\mathrm{d}z}{\mathrm{d}y}$: and these are called the *partial differential coefficients*, being analogous to those bearing the same name considered in chapter IV. We have seen, in functions of a single variable, that if that variable take an increment, and the function be developed, what we have called the differential coefficient will be the coefficient of the first power of the increment in that development; so here, as will be shortly shown, the partial differential coefficients are no other than the coefficients of the first power of the increments in the development of the function from which they are derived. As to the *partial differentials* they are obviously $\frac{\mathrm{d}z}{\mathrm{d}x}\mathrm{d}x$ and $\frac{\mathrm{d}z}{\mathrm{d}y}\mathrm{d}y$, and hence we call $\frac{\mathrm{d}z}{\mathrm{d}x}\mathrm{d}x + \frac{\mathrm{d}z}{\mathrm{d}y}\mathrm{d}y$ the *total differential* of the function, that is,

$$\mathrm{d}z = \frac{\mathrm{d}z}{\mathrm{d}x}\mathrm{d}x + \frac{\mathrm{d}z}{\mathrm{d}y}\mathrm{d}y,$$

and we immediately see that this form becomes the same as that given

in chapter IV. for the differential of $F(x, y)$ as soon as we suppose y to be a function of x, for we then have

$$\left\{\frac{dz}{dx}\right\} = \frac{dz}{dx} + \frac{dz}{dy}\frac{dy}{dx},$$

as indeed we ought.

In a similar manner, if the function consist of a greater number of independent variables as $u = F(x, y, z, \&c.)$ we should necessarily have as many independent differentials, of which the aggregate would be the *total* differential of the function; that is,

$$du = \frac{du}{dx}dx + \frac{du}{dy}dy + \frac{du}{dz}dz + \&c.$$

Hence, whether the variables are dependent or independent, we infer, generally, that

The total differential of any function is the sum of the several partial differentials arising from differentiating the function relatively to each variable in succession, as if all the others were constants.

We shall add but few examples in functions of independent variables, seeing that the process is exactly the same as for functions of dependent variables.

1. $d(x + y) = dx + dy.$

2. $d\,xy = ydx + xdy.$

3. $$d\frac{x}{y} = \frac{ydx - xdy}{y^2}.$$

4. $$d\frac{ay}{\sqrt{x^2 + y^2}} = \frac{ax^2dy - ayxdx}{(x^2 + y^2)^{\frac{3}{2}}}.$$

5. $$d\tan^{-1}\frac{x}{y} = \frac{ydx - xdy}{y^2 + x^2}.$$

6. $$d\frac{y}{3y^2 - x} = \frac{ydx - 3y^2dy - xdy}{(3y^2 - x)^2}.$$

7. $$d\,a^x b^y c^z = a^x b^y c^z(dx \log a + dy \log b + dz \log c).$$

8. $$\mathrm{d}\log\tan\frac{x}{y}=\frac{y\mathrm{d}x-x\mathrm{d}y}{y^2\sin\frac{x}{y}\cos\frac{x}{y}}=\frac{2\,(y\mathrm{d}x-x\mathrm{d}y)}{y^2\sin\frac{2x}{y}}.$$

9. $$\mathrm{d}y^x=y^x\log y\mathrm{d}x+y^{x-1}x\mathrm{d}y.$$

(57.) If the function that z is of x, y is given implicitly, that is by the equation

$$u=\mathrm{F}(x,y,z)=0,$$

then

$$\{\frac{\mathrm{d}u}{\mathrm{d}x}\}^{\bullet}=0, \text{ and } \{\frac{\mathrm{d}u}{\mathrm{d}y}\}=0,$$

but (39),

$$\{\frac{\mathrm{d}u}{\mathrm{d}x}\}=\frac{\mathrm{d}u}{\mathrm{d}x}+\frac{\mathrm{d}u}{\mathrm{d}z}\frac{\mathrm{d}z}{\mathrm{d}x}=0,$$

$$\{\frac{\mathrm{d}u}{\mathrm{d}y}\}=\frac{\mathrm{d}u}{\mathrm{d}y}+\frac{\mathrm{d}u}{\mathrm{d}z}\frac{\mathrm{d}z}{\mathrm{d}y}=0,$$

$$\therefore \mathrm{d}u=(\frac{\mathrm{d}u}{\mathrm{d}x}+\frac{\mathrm{d}u}{\mathrm{d}z}\frac{\mathrm{d}z}{\mathrm{d}x})\mathrm{d}x+(\frac{\mathrm{d}u}{\mathrm{d}y}+\frac{\mathrm{d}u}{\mathrm{d}z}\frac{\mathrm{d}z}{\mathrm{d}y})\mathrm{d}y=0,$$

$$=\{\frac{\mathrm{d}u}{\mathrm{d}x}\}\mathrm{d}x+\{\frac{\mathrm{d}u}{\mathrm{d}y}\}\mathrm{d}y.$$

• The brackets are employed here for the same purpose as at (37), viz. to imply the *total differential coefficient derived from u, considered as a function of a single variable.* This form it will be necessary to adopt whenever u contains, besides x, other variables that are functions of x, provided we wish to express the *total* coefficient with respect to x. No ambiguity can arise from our calling these same coefficients *partial* in one sense, and *total* in another. They are *partial* coefficients in relation to the *whole* variation of u, but they are *total* coefficients as far as that variable is concerned whose differential forms the denominator; and it may be remarked here, once for all, that when we enclose a differential coefficient in brackets, we mean the *total differential coefficient* to be understood, arising from considering the function, whose differential is the numerator, as simply a function of the variables whose differentials form the denominator.

Thus: let $Ax^2 + By^2 + Cz^2 - 1 = 0$,

$$\therefore \{\frac{du}{dx}\} = 2Ax + 2Cz\frac{dz}{dx} = 0,$$

$$\{\frac{du}{dy}\} = 2By + 2Cz\frac{dz}{dy} = 0;$$

$$\therefore du = (Ax + Cz\frac{dz}{dx})\,dx + (By + Cz\frac{dz}{dy})\,dy = 0.$$

(58.) If $u = F(z)$, z being a function of x and y, the two differential coefficients are (33)

$$\frac{du}{dx} = \frac{du}{dz}\frac{dz}{dx}, \quad \frac{du}{dy} = \frac{du}{dz}\frac{dz}{dy},$$

and the total differential is therefore

$$du = \frac{du}{dz}\frac{dz}{dx}dx + \frac{du}{dz}\frac{dz}{dy}dy.$$

Now it is worthy of notice, that *the ratio of the two partial differential coefficients, is independent of* F, so that this may be any function whatever. Thus

$$\frac{du}{dx} \div \frac{du}{dy} = \frac{du}{dz}\frac{dz}{dx} \div \frac{du}{dz}\frac{dz}{dy} = \frac{dz}{dx} \div \frac{dz}{dy},$$

which is an important property, since it enables us to eliminate any arbitrary function F of a determinate function $f(x, y)$ of two variables. We shall often have occasion to employ it in discussing the theory of curve surfaces. By means of this property too we may readily ascertain whether an expression containing two variables is a function of any proposed combination of those variables. For, calling this combination z and the function u, we shall merely have to ascertain whether or not the above condition exists; or, which is the same thing, whether or not the condition

$$\frac{du}{dx}\frac{dz}{dy} - \frac{du}{dy}\frac{dz}{dx} = 0$$

exists. For instance, suppose we wished to know whether $u = x^4 + 2x^2y^2 + y^4$ is a function of $z = x^2 + y^2$.

Here

$$\frac{du}{dx} = 4x^3 + 4xy^2, \quad \frac{du}{dy} = 4x^2y + 4y^3; \quad \frac{dz}{dy} = 2y, \quad \frac{dz}{dx} = 2x;$$

$$\therefore \frac{du}{dx}\frac{dz}{dy} - \frac{du}{dy}\frac{dz}{dx} = (4x^3 + 4xy^2)\,2y - (4x^2y + 4y^3)\,2x = 0;$$

consequently, since the proposed condition exists, we infer that u *is* a function of z.

We shall now proceed to apply Taylor's theorem to functions of two independent variables.

Development of Functions of two Independent Variables.

(59.) In the function $z = \mathrm{F}(x, y)$ suppose x takes the increment h. the function will become $\mathrm{F}(x + h, y)$, y remaining unchanged, since it is independent of x; then, by Taylor's theorem,

$$\mathrm{F}(x + h, y) = z + \frac{dz}{dx}h + \frac{d^2z}{dx^2}\frac{h^2}{1 \cdot 2} + \frac{d^3z}{dx^3}\frac{h^3}{1 \cdot 2 \cdot 3} + \&c. \ldots (1).$$

But if y also take an increment k, then z will become

$$z + \frac{dz}{dy}k + \frac{d^2z}{dy^2}\frac{k^2}{1 \cdot 2} + \frac{d^3z}{dy^3}\frac{k^3}{1 \cdot 2 \cdot 3} + \&c.$$

so that in the expression (1) we must for $\dfrac{dz}{dx}$ substitute

$$\frac{dz}{dx} + \frac{d\dfrac{dz}{dy}}{dx}k + \frac{d\dfrac{d^2z}{dy^2}}{dx}\frac{k^2}{1 \cdot 2} + \frac{d\dfrac{d^3z}{dy^3}}{dx}\frac{k^3}{1 \cdot 2 \cdot 3} + \&c.$$

for $\dfrac{d^2z}{dx^2}$;

$$\frac{d^2z}{dx^2} + \frac{d^2\frac{dz}{dy}}{dx^2}k + \frac{d^2\frac{d^2z}{dy^2}}{dx^2}\frac{k^2}{1\cdot 2} + \frac{d^2\frac{d^3z}{dy^3}}{dx^2}\frac{k^3}{1\cdot 2\cdot 3} + \&c.$$

for $\frac{d^3z}{dx^3}$,

$$\frac{d^3z}{dx^3} + \frac{d^3\frac{dz}{dy}}{dx^3}k + \frac{d^3\frac{d^2z}{dy^2}}{dx^3}\frac{k^2}{1\cdot 2} + \frac{d^3\frac{d^3z}{dy^3}}{dx^3}\frac{k^3}{1\cdot 2\cdot 3} + \&c.$$

and so on. Before, however, we actually make these substitutions, we shall, for abridgment, write

$$\frac{d^2z}{dy\,dx} \text{ for } \frac{d\frac{dz}{dy}}{dx}, \quad \frac{d^3z}{dy^2\,dx} \text{ for } \frac{d\frac{d^2z}{dy^2}}{dx},$$

$$\text{and generally } \frac{d^{q+p}z}{dy^q\,dx^p} \text{ for } \frac{d^p\frac{d^qz}{dy^q}}{dx^p},$$

this last expression implying that after having determined the qth differential coefficient of the function z relatively to the variable y, the pth differential coefficient of this is taken relatively to the other variable x. Hence, the result of the proposed substitutions in (1) will be

$$F(x+h,\ y+k) =$$

$$z + \frac{dz}{dx}h + \frac{dz}{dy}k \quad \Big| \quad + \frac{d^2z}{dx^2}\frac{h^2}{1\cdot 2} + \frac{d^2z}{dy\,dx}kh + \frac{d^2z}{dy^2}\frac{k^2}{1\cdot 2} \quad \Big| \quad + \frac{d^3z}{dx^3}\frac{h^3}{1\cdot 2\cdot 3} + \frac{d^3z}{dy\,dx^2}\frac{kh^2}{1\cdot 2} + \frac{d^3z}{dy^2\,dx}\frac{k^2h}{1\cdot 2} + \frac{d^3z}{dy^3}\frac{k^3}{1\cdot 2\cdot 3} \quad \Big| \quad + \&c.$$

The general term of the development being

$$\frac{\mathrm{d}^{q+p} z}{\mathrm{d}y^q \, \mathrm{d}x^p} \, \frac{k^q h^p}{(1 \cdot 2 \ldots q)(1 \cdot 2 \ldots p)},$$

if in the proposed function, $z = \mathrm{F}(x, y)$, we had supposed y to vary first, then, instead of (1), we should have had

$$\mathrm{F}(x, y + k) = z + \frac{\mathrm{d}z}{\mathrm{d}y} k + \frac{\mathrm{d}^2 z}{\mathrm{d}y^2} \, \frac{k^2}{1 \cdot 2} + \frac{\mathrm{d}^3 z}{\mathrm{d}y^3} \, \frac{k^3}{1 \cdot 2 \cdot 3} + \&\mathrm{c.} \ldots (2).$$

But, if x take the increment h, z will become

$$z + \frac{\mathrm{d}z}{\mathrm{d}x} h + \frac{\mathrm{d}^2 z}{\mathrm{d}x^2} \, \frac{h^2}{1 \cdot 2} + \frac{\mathrm{d}^3 z}{\mathrm{d}x^3} \, \frac{h^3}{1 \cdot 2 \cdot 3} + \&\mathrm{c.}$$

and, therefore, we must substitute, agreeably to the foregoing notation,

for $\dfrac{\mathrm{d}z}{\mathrm{d}y}$

$$\frac{\mathrm{d}z}{\mathrm{d}y} + \frac{\mathrm{d}^2 z}{\mathrm{d}x \, \mathrm{d}y} h + \frac{\mathrm{d}^3 z}{\mathrm{d}x^2 \, \mathrm{d}y} \, \frac{h^2}{1 \cdot 2} + \frac{\mathrm{d}^4 z}{\mathrm{d}x^3 \, \mathrm{d}y} \, \frac{h^3}{1 \cdot 2 \cdot 3} + \&\mathrm{c.}$$

for $\dfrac{\mathrm{d}^2 z}{\mathrm{d}y^2}$

$$\frac{\mathrm{d}^2 z}{\mathrm{d}y^2} + \frac{\mathrm{d}^3 z}{\mathrm{d}x \, \mathrm{d}y^2} h + \frac{\mathrm{d}^4 z}{\mathrm{d}x^2 \, \mathrm{d}y^2} \, \frac{h^2}{1 \cdot 2} + \frac{\mathrm{d}^5 z}{\mathrm{d}x^3 \, \mathrm{d}y^2} \, \frac{h^3}{1 \cdot 2 \cdot 3} + \&\mathrm{c.}$$

for $\dfrac{\mathrm{d}^3 z}{\mathrm{d}y^3}$

$$\frac{\mathrm{d}^3 z}{\mathrm{d}y^3} + \frac{\mathrm{d}^4 z}{\mathrm{d}x \, \mathrm{d}y^3} h + \frac{\mathrm{d}^5 z}{\mathrm{d}x^2 \, \mathrm{d}y^3} \, \frac{h^2}{1 \cdot 2} + \frac{\mathrm{d}^6 z}{\mathrm{d}x^3 \, \mathrm{d}y^3} \, \frac{h^3}{1 \cdot 2 \cdot 3} + \&\mathrm{c.}$$

and so on; so that the development would be

$$\mathrm{F}(x + h, y + k) =$$

$$z + \frac{dz}{dx}h \quad \frac{dz}{dy}k \qquad + \frac{d^2z}{dx^2}\frac{h^2}{1\cdot 2} \quad \frac{d^2z}{dx\,dy}hk \quad \frac{d^2z}{dy^2}\frac{k^2}{1\cdot 2} \qquad + \frac{d^3z}{dx^3}\frac{h^3}{1\cdot 2\cdot 3} \quad \frac{d^3z}{dx^2\,dy}\frac{h^2k}{1\cdot 2} \quad \frac{d^3z}{dx\,dy^2}\frac{hk^2}{1\cdot 2} \quad \frac{d^3z}{dy^3}\frac{k^3}{1\cdot 2\cdot 3} \qquad + \&c.$$

the general term being

$$\frac{d^{p+q}z}{dx^p\,dy^q}\,\frac{h^p\,k^q}{(1\cdot 2\ldots p)\,(1\cdot 2\ldots q)}.$$

As this development must be identical with that exhibited above, we have, by equating the like powers of h and k,

$$\frac{d^2z}{dy\,dx} = \frac{d^2z}{dx\,dy}, \quad \frac{d^3z}{dy\,dx^2} = \frac{d^3z}{dx^2\,dy},$$

and generally

$$\frac{d^{q+p}z}{dy^q\,dx^p} = \frac{d^{p+q}z}{dx^p\,dy^q};$$

we conclude, therefore, that if we first determine the qth differential coefficient relatively to the variable y, and then the pth differential coefficient of this relatively to the variable x, the final result will be the same as if we first determine the pth differential coefficient relatively to x, and then the qth differential coefficient of this relatively to y; so that *the result is the same in whichever order the differentiations are performed.*

(60.) We see from the foregoing development, that the partial differential coefficients of the first order are the coefficients of h and k, the first power of the increments; so that the term containing these first powers is, in this respect, analogous to that containing the first power of the increment in the development of functions of a single variable, and,

by a very slight transformation, it will be seen that the same analogy extends throughout all the terms of the two developments. For the development just given may be put under the form

$$F(x+h,\ y+k) =$$

$$z$$

$$+ \left(\frac{dz}{dx}h + \frac{dz}{dy}k\right)$$

$$+ \frac{1}{2}\left(\frac{d^2z}{dx^2}h^2 + 2\frac{d^2z}{dx\,dy}hk + \frac{d^2z}{dy^2}k^2\right)$$

$$+ \frac{1}{2\cdot 3}\left(\frac{d^3z}{dx^3}h^3 + 3\frac{d^3z}{dx^2\,dy}h^2k + 3\frac{d^3z}{dx\,dy^2}hk^2 + \frac{d^3z}{dy^3}k^3\right)$$

$$+ \quad \&c.$$

where the partial differential coefficients in each term are identical with those which appear in the differential of the preceding term, as the actual differentiation shows, thus:

$$dz = \frac{dz}{dx}dx + \frac{dz}{dy}dy \quad \ldots\ (1),$$

the coefficients $\frac{dz}{dx}$, $\frac{dz}{dy}$, being functions of x and y, we have

$$d\frac{dz}{dx} = \frac{d^2z}{dx^2}dx + \frac{d^2z}{dx\,dy}dy$$

$$d\frac{dz}{dy} = \frac{d^2z}{dy\,dx}dx + \frac{d^2z}{dy^2}dy,$$

and, consequently,

$$d^2z = \frac{d^2z}{dx^2}dx^2 + 2\frac{d^2z}{dx\,dy}dx\,dy + \frac{d^2z}{dy^2}dy^2 \quad \ldots\ (2).$$

---

* This can be regarded as the second differential of the function z only on the supposition that the variations of x and y are both constant and independent; so that we cannot deduce from this expression, that for the second differential of z, on the supposition that y is a function of x. For

In like manner, these coefficients being functions of x and y, we have

$$d\frac{d^2z}{dx^2} = \frac{d^3z}{dx^3}dx + \frac{d^3z}{dx^2\,dy}dy$$

$$d\frac{d^2z}{dx\,dy} = \frac{d^3z}{dx^2\,dy}dx + \frac{d^3z}{dx\,dy^2}dy$$

$$d\frac{d^2z}{dy^2} = \frac{d^3z}{dy^2\,dx}dx + \frac{d^3z}{dy^3}dy;$$

so that

$$d^3z = \frac{d^3z}{dx^3}dx^3 + 3\,\frac{d^3z}{dx^2\,dy}dx^2\,dy + 3\,\frac{d^3z}{dx\,dy^2}dx\,dy^2 +$$

$$\frac{d^3z}{dy^3}dy^3 \ldots . \quad (3),$$

and so on: the numerical coefficients agreeing with those in the corresponding powers of the expanded binomial.

(61.) Having now applied Taylor's theorem to functions of two variables, we may equally extend Maclaurin's theorem. For, if in the foregoing development, we suppose x and y each $= 0$, the development will become that of the function F (h, k) according to the powers of h and k; or, substituting x and y for the symbols h and k, since these are indeterminate, we have

$$\cdot\cdot z = F(x, y) = [z] + ([\frac{dz}{dx}]\,x + [\frac{dz}{dy}]\,y) + \tfrac{1}{2}\,([\frac{d^2z}{dx^2}]\,x^2 +$$

$$2\,[\frac{d^2z}{dx\,dy}]\,xy + [\frac{d^2z}{dy^2}]\,y^2) + \&c.$$

The principles by which we have thus far extended the theorems of Taylor and Maclaurin are sufficient to enable us to extend these

---

although when y is a function of x, the original integral equation is more restricted than when x and y are independent, yet, if in the latter case we regard the independent variations of x and y to be constant, the successive differentials of any function of them will obviously be more restricted than in the former case, where the variation of x alone is regarded as constant.

theorems still further, even to the development of functions of any number of variables whatever, but this is unnecessary. It may be remarked, however, that if we wish to develop a function of several variables according to the powers of one of them, it may be done independently of anything taught in this chapter; for, if all the variables but this one were constants, the development would agree with that already established for functions of a single variable, and, as these constants may take any value whatever, they may obviously be replaced by so many independent variables. We shall give one instance of this extension of Maclaurin's theorem to a function of two independent variables, choosing a form of extensive application, and of which the development is known by the name of

Lagrange's Theorem.

(62.) The function which we here propose to develop according to the powers of x, is

$$u = \mathrm{F}(z) \text{ in which } z = y + xf(z),$$

z being obviously a function of the independent variables x and y. We shall first develop $z = y + xf(z)$ according to the powers of x; this development is by Maclaurin's theorem

$$z = [z] + [\frac{\mathrm{d}z}{\mathrm{d}x}]\, x + [\frac{\mathrm{d}^2 z}{\mathrm{d}x^2}]\, \frac{x^2}{1 \cdot 2} + [\frac{\mathrm{d}^3 z}{\mathrm{d}x^3}]\, \frac{x^3}{1 \cdot 2 \cdot 3} + \&c.$$

and if we express, according to the notation of Lagrange, the successive partial differential coefficients of $f(z)$, relatively to x, by $f'(z)$, $f''(z)$, $f'''(z)$, &c. we shall have

$$z = y + xf(z)$$

$$\frac{\mathrm{d}z}{\mathrm{d}x} = f(z) + xf'(z)$$

$$\frac{\mathrm{d}^2 z}{\mathrm{d}x^2} = 2f'(z) + xf''(z) = 2\,\frac{\mathrm{d}f(z)}{\mathrm{d}z}\,\frac{\mathrm{d}z}{\mathrm{d}x} + xf''(z)$$

$$\frac{d^3z}{dx^3} = 3f''(z) + xf'''(z) = 3\left(\frac{d^2f(z)}{dz^2}\,\frac{dz^2}{dx^2} + \frac{df(z)}{dz}\,\frac{d^2z}{dx^2}\right) + xf'''(z)$$

&c. &c.

Consequently, when $x = 0$,

$$[z] = y$$

$$\left[\frac{dz}{dx}\right] = f(y)$$

$$\left[\frac{d^2z}{dx^2}\right] = 2\,\frac{df(y)}{dy}\,f(y) = \frac{d(f(y))^2}{dy}$$

$$\left[\frac{d^3z}{dx^3}\right] = 3\,\frac{d^2f(y)}{dy^2}\,(f(y))^2 + 3\,\frac{df(y)}{dy}\,\frac{d(f(y))^2}{dy} = \frac{d^2(f(y))^3}{dy^2}$$

&c. &c.

Hence

$$z = y + xf(z) = y + f(y)\,\frac{x}{1} + \frac{d(f(y))^2}{dy}\,\frac{x^2}{1\cdot 2} + \frac{d^2(f(y))^3}{dy^2}\,\frac{x^3}{1\cdot 2\cdot 3} +$$

&c. . . . . (1).

Now, instead of this development, we should obviously have obtained that of $F(z) = F(y + xf(z))$, if in place of z and its differential coefficients we had employed $F(z)$ and its differential coefficients. We should then have had

$$u = F(y + xf(z)) \quad . \quad . \quad \text{therefore} \quad . \quad . \quad [u] = F(y)$$

$$\frac{du}{dx} = \frac{du}{dz}\,\frac{dz}{dx} \quad . \quad . \quad . \quad . \quad . \quad . \quad . \quad . \quad . \quad . \quad \left[\frac{du}{dx}\right] = \frac{dF(y)}{dy}\,f(y)$$

$$\frac{d^2u}{dx^2} = \frac{d^2u}{dz^2}\left(\frac{dz}{dx}\right)^2 + \frac{du}{dz}\,\frac{d^2z}{dx^2} \quad . \quad . \quad . \quad \left[\frac{d^2u}{dx^2}\right] = \frac{d^2F(y)}{dy^2}\,(f(y))^2 +$$

$$\frac{dF(y)}{dy}\,\frac{d(f(y))^2}{dy} = \frac{d\,\dfrac{dF(y)}{dy}\,(f(y))^2}{dy}$$

&c. &c.

Hence

$$F(z)=F(y+xf(z))=F(y)+\frac{dF(y)}{dy}f(y)\,\frac{x}{1}+\frac{d\,\frac{dF(y)}{dy}(f(y))^2}{dy}$$

$$\frac{x^2}{1\cdot 2}+\frac{d^2\,\frac{dF(y)}{dy}(f(y))^3}{dy^2}\,\frac{x^3}{1\cdot 2\cdot 3}+\&c.\ldots.\ (2),$$

and this is *Lagrange's Theorem.*•

From this remarkable expression, which includes that marked (1), other forms may be readily deduced as particular cases. Of these the two following are the most important.

Put $x=1$, then the formula (1) becomes

$$z=y+f(z)=y+f(y)+\frac{d(f(y))^2}{dy}\,\frac{1}{1\cdot 2}+\frac{d^2(f(y))^3}{dy^2}\,\frac{1}{1\cdot 2\cdot 3}+$$

$$\&c.\ldots.\ (3),$$

and the formula (2),

$$u=F(y+f(z))=F(y)+\frac{dF(y)}{dy}f(y)+\frac{d\,\frac{dF(y)}{dy}(f(y))^2}{dy}\,\frac{1}{1\cdot 2}+$$

$$\frac{d^2\,\frac{dF(y)}{dy}(f(y))^3}{dy^2}\,\frac{1}{1\cdot 2\cdot 3}+\&c.\ldots.\ (4).$$

(63.) We shall terminate the present section with one or two examples of the application of these formulas, referring the student for more ample details on this subject to *Lagrange's Resolution des Equations Numeriques*, note xi.; and *Jephson's Fluxional Calculus*, vol. i.

• For another and very complete demonstration of this theorem see note (B) at the end.

EXAMPLES.

1. Given $z^3 - qz + r = 0$, to develop z according to the powers of r.

Since here $z = \frac{r}{q} + \frac{z^3}{q}$, we have $y = \frac{r}{q}, f(z) = \frac{1}{q} z^3 \therefore f(y) = \frac{1}{q} (\frac{r}{q})^3$

$$\therefore \frac{d(f(y))^2}{dy} = \frac{1}{q^2} \frac{dy^6}{dy} = \frac{6}{q^2} y^5, \frac{d^2(f(y))^3}{dy^2} = \frac{1}{q^3} \frac{d^2y^9}{dy^2} = \frac{9}{q^3} \frac{dy^8}{dy} =$$

$$\frac{8 \cdot 9}{q^3} y^7, \text{ \&c.}$$

Hence, by the formula (3), we have, by putting for y its value $\frac{r}{q}$,

$$z = \frac{r}{q} + \frac{1}{q} \frac{r^3}{q^3} + \frac{6}{1 \cdot 2q^2} \frac{r^5}{q^5} + \frac{8 \cdot 9}{1 \cdot 2 \cdot 3q^3} \frac{r^7}{q^7} + \text{\&c.}$$

$$= \frac{r}{q} (1 + \frac{r^2}{q^3} + \frac{6r^4}{1 \cdot 2q^6} + \frac{8 \cdot 9r^6}{1 \cdot 2 \cdot q^9} + \text{\&c.})$$

2. Given the radius vector of an ellipse, viz. (*Anal. Geom.* p. 133,)

$$r = a \frac{1 - e^2}{1 + e \cos \omega}$$

to develop r^n, according to the powers of $\cos \omega$.

Since $r = a(1 - e^2) - e^2 \cos \omega \cdot r$, we have, by putting y for $a(1 - e^2)$ and x for $-e \cos \omega$,

$$F(r) = F(y + xf(r)) = F(y + x \cdot r) = (y + x \cdot r)^n.$$

Hence, by the formula (2),

$$r^n = y^n + \frac{dy^n}{dy} y \frac{x}{1} + \frac{d \frac{dy^n}{dy} y^2}{dy} \frac{x^2}{1 \cdot 2} +$$

$$\frac{d^2 \frac{dy^n}{dy} y^3}{dy^2} \frac{x^3}{1 \cdot 2 \cdot 3} + \text{\&c.}$$

$$= y^n + ny^n \frac{x}{1} + n(n+1)\, y^n \frac{x^2}{1\cdot 2} +$$

$$n(n+1)(n+2)\, y^n \frac{x^3}{1\cdot 2\cdot 3} + \&c.$$

$$= a^n (1-e^2)^n \left(1 - \frac{ne\cos\omega}{1} + \frac{n(n+1)}{1\cdot 2} e^2 \cos^2\omega - \right.$$

$$\left. \frac{n(n+1)(n+2)}{1\cdot 2\cdot 3} e^3 \cos^3\omega + \&c.\right)$$

3. It is required to revert the series

$$\alpha + \beta z + \gamma z^2 + \delta z^3 + \&c. = 0,$$

that is, to express the value of z in terms of the coefficients. Here

$$z = -\frac{\alpha}{\beta} - \frac{z^2}{\beta}(\gamma + \delta z + \&c.) = y + f(z);$$

therefore, by the formula (3),

$$z = y - \frac{y^2}{\beta}(\gamma + \delta y + \&c.) + \frac{\mathrm{d}\frac{y^4}{\beta^2}(\gamma + \delta y + \&c.)^2}{\mathrm{d}y} \frac{1}{1\cdot 2}$$

$$- \frac{\mathrm{d}^2\frac{y^6}{\beta^3}(\gamma + \delta y + \&c.)^3}{\mathrm{d}y^2} \frac{1}{1\cdot 2\cdot 3} + \&c.$$

$$= y - \frac{y^2}{\beta}(\gamma + \delta y + \&c.) + 2\frac{y^3}{\beta^2}(\gamma + \delta y + \&c.)^2 +$$

$$\frac{y^4}{\beta^2}(\gamma + \delta y + \&c.)(\delta + 2\varepsilon y + \&c.)$$

$$- 5\frac{y^4}{\beta^3}(\gamma + \delta y + \&c.)^3 + \&c.$$

$$= y - \frac{\gamma}{\beta} y^2 - \frac{\delta}{\beta} y^3 - \frac{\varepsilon}{\beta} y^4 - \&c.$$

$$+\frac{2\gamma^2}{\beta^2}y^3+\frac{5\gamma\delta}{\beta^2}y^4+\&c.$$

$$-\frac{5\gamma^3}{\beta^3}y^4+\&c.$$

$$+\&c.$$

where $y=-\frac{\alpha}{\beta}$, consequently

$$z=-\frac{1}{\beta}\alpha-\frac{\gamma}{\beta^3}\alpha^2+\frac{\delta\beta-2\gamma^2}{\beta^5}\alpha^3-\frac{\beta^2\epsilon-5\gamma\delta\beta+5\gamma^3}{\beta^7}\alpha^4+\&c.$$

4. Given $1-z+az=0$ to develop $\log z$, according to the powers of a.

$$\log z=a+\tfrac{1}{2}a^2+\tfrac{1}{3}a^3+\tfrac{1}{4}a^4+\&c.^*$$

CHAPTER VIII.

ON THE MAXIMA AND MINIMA VALUES OF FUNCTIONS OF TWO VARIABLES, AND ON CHANGING THE INDEPENDENT VARIABLE.

(64.) It remains to complete the theory of maxima and minima by applying the principles established in chapter vi. to functions of two independent variables.

The same character belongs to a maximum or minimum function of two variables that belongs to a maximum or minimum function of one

---

* This we know from other principles; for, since the proposed expression reduces to $z=\frac{1}{1-a}$ $\therefore$ $\log z=-\log(1-a)$ and this, in the hyperbolic system, is equal to the above series. (*See the Algebra*, p. 186.)

variable, that is, the maximum value exceeds the contiguous values of the function, and the minimum value falls short of them.

Hence, if

$$z = \mathrm{F}(x, y)$$

be any function of two variables, which becomes a maximum for certain particular values of them, then, h and k being finite increments, however small, the condition is that, between such finite values and 0, we must always have

$$\mathrm{F}[x, y] > \mathrm{F}[x \pm h, y \pm k],$$

and, consequently, (60),

$$[\pm \frac{\mathrm{d}z}{\mathrm{d}x} h \pm \frac{\mathrm{d}z}{\mathrm{d}y} k] + \tfrac{1}{2} [\frac{\mathrm{d}^2 z}{\mathrm{d}x^2} h^2 \pm 2 \frac{\mathrm{d}^2 z}{\mathrm{d}x\,\mathrm{d}y} hk + \frac{\mathrm{d}^2 z}{\mathrm{d}y^2} k^2] + \&c. < 0.$$

If, therefore, of the small values which we suppose h and k to take, h be the smaller, a part of k may be taken so small as to be less than h; or, which is the same thing, equal to one of the values of h between the proposed value and 0, so that we have $h' = k'$; therefore, the above condition is

$$[\pm \frac{\mathrm{d}z}{\mathrm{d}x} \pm \frac{\mathrm{d}z}{\mathrm{d}y}] h' + \tfrac{1}{2} [\frac{\mathrm{d}^2 z}{\mathrm{d}x^2} \pm 2 \frac{\mathrm{d}^2 z}{\mathrm{d}x\,\mathrm{d}y} + \frac{\mathrm{d}^2 z}{\mathrm{d}y^2}] h'^2 + \&c. < 0.$$

This condition being similar to (1) art. (49), we infer, by the same reasoning, that

$$\pm \frac{\mathrm{d}z}{\mathrm{d}x} \pm \frac{\mathrm{d}z}{\mathrm{d}y} = 0,$$

which cannot be, for both the signs $\pm$, unless

$$\frac{\mathrm{d}z}{\mathrm{d}x} = 0, \frac{\mathrm{d}z}{\mathrm{d}y} = 0 \ldots . (1).$$

By continuing to imitate the reasoning in (49), we find that these same conditions must exist for all the values of the variables that render the function a *minimum*.

Hence (49), we have, in the case of a maximum, the condition

$$\tfrac{1}{2} [\frac{\mathrm{d}^2 z}{\mathrm{d}x^2} \pm 2 \frac{\mathrm{d}^2 z}{\mathrm{d}x\,\mathrm{d}y} + \frac{\mathrm{d}^2 z}{\mathrm{d}y^2}] h'^2 + \&c. < 0,$$

and in the case of a minimum,

$$\tfrac{1}{2}\left[\frac{d^2z}{dx^2} \pm 2\,\frac{d^2z}{dx\,dy} + \frac{d^2z}{dy^2}\right]h'^2 + \&c. > 0,$$

so that, supposing these first terms do not vanish for the values of x and y given by (1), the condition of maximum is

$$\left[\frac{d^2z}{dx^2} \pm 2\,\frac{d^2z}{dx\,dy} + \frac{d^2z}{dy^2}\right] < 0,$$

and the condition of minimum,

$$\left[\frac{d^2z}{dx^2} \pm 2\,\frac{d^2z}{dx\,dy} + \frac{d^2z}{dy^2}\right] > 0,$$

In either case, therefore, the expression within the brackets must have the same sign independently of the sign of the middle term. To determine upon what other condition this depends, let us represent the expression by

$$A \pm 2B + C \text{ or } A\left(1 \pm 2\,\frac{B}{A} + \frac{C}{A}\right).$$

Adding $\frac{B^2}{A^2} - \frac{B^2}{A^2} = 0$ to the quantity within the parentheses, its form is

$$A\left\{\left(1 \pm \frac{B}{A}\right)^2 + \frac{C}{A} - \frac{B^2}{A^2}\right\}.$$

Now this expression will always have the same sign as A, provided C has, and that $\frac{C}{A} > \frac{B^2}{A^2}$, that is, $AC > B^2$ or $AC - B^2 > 0$, because then the factor of A will be necessarily positive. Hence, besides (1), the condition that a maximum or a minimum may exist is

$$\left[\frac{d^2z}{dx^2}\,\frac{d^2z}{dy^2} - \left(\frac{d^2z}{dx\,dy}\right)^2\right] > 0 \quad \ldots \quad (2),$$

and we are to distinguish the *maximum* from the *minimum* by ascertaining whether the proposed values of x and y render

$$\frac{d^2z}{dx^2} < 0 \text{ or } > 0,$$

or, which amounts to the same, whether

$$\frac{d^2z}{dy^2} < 0 \text{ or } > 0,$$

since $\frac{d^2z}{dx^2}$ and $\frac{d^2z}{dy^2}$ have the same sign.

Should any of the values determined from (1) cause the coefficient of h'^2 to vanish, there will be no maximum or minimum for those values, unless the coefficient of the following term vanishes also.

EXAMPLES.

(65.) 1. To determine the shortest distance between two straight lines situated in space.

Let the equations of the two lines be

$$\left.\begin{aligned} x &= az + \alpha \\ y &= bz + \beta \end{aligned}\right\} \text{ and } \left\{\begin{aligned} x' &= a'z' + \alpha' \\ y' &= b'z' + \beta' \end{aligned}\right. \quad \ldots\ldots (1)$$

then the expression for the distance between any two points (z, y, z), $(x' y' z')$ is *(Anal. Geom.* p. 226),

$$D^2 = u = (x - x')^2 + (y - y')^2 + (z - z')^2$$

$$= (\alpha - \alpha' + az - a'z')^2 + (\beta - \beta' + bz - b'z')^2 + (z - z')^2$$

and this expression, containing the two independent variables z, z' is to be a minimum. Hence by the conditions (1), p. 121,

$$\left.\begin{aligned} \frac{du}{dz} &= 2(z - z') + 2a(\alpha - \alpha' + az - a'z') + 2b(\beta - \beta' + bz - b'z') = 0 \\ \frac{du}{dz'} &= -2(z - z') + 2a'(\alpha - \alpha' + az - a'z') + 2b'(\beta - \beta' + bz - b'z') = 0 \end{aligned}\right\} \ldots (2),$$

and from these equations the proper values of z, z' may be readily determined, which substituted in the expression for u, render it the least possible. That these values really belong to a minimum is evident, because,

$\frac{d^2u}{dz^2}=2(1+a^2+b^2)$, $\frac{d^2u}{dz'^2}=2(1+a'^2+b'^2)$, $\frac{d^2u}{dzdz'}=-2(1+aa'+bb')$

and this proves that $\frac{d^2u}{dz^2}$, $\frac{d^2u}{dz'^2}$ are both positive, and that

$$\frac{d^2u}{dz^2}\,\frac{d^2u}{dz'^2}-\left(\frac{d^2u}{dz\,dz'}\right)^2>0.$$

Since the equations of a straight line passing through two points (x, y, z), (x', y', z') are

$$\left.\begin{aligned} x-x'&=a'(z-z')\\ y-y'&=b'(z-z')\end{aligned}\right\}$$

we have, by substitution, when these points are on the lines (1),

$$\left.\begin{aligned} \alpha-\alpha'+az-a'z'&=a''(z-z')\\ \beta-\beta'+bz-b'z'&=b''(z-z')\end{aligned}\right\}\quad\ldots\ldots(3);$$

hence, if this line be that in question, we have, by combining the equations (2) with these, the conditions

$$1+aa''+bb''=0,\quad 1+a'a''+b'b''=0,$$

which conditions show, that this *minimum* straight line is perpendicular to *both* the lines (1). (See *Anal. Geom.* p. 228.) From these conditions we get

$$a''=\frac{b'-b}{a'b-ab'},\quad b''=\frac{a'-a}{a'b-ab'},$$

by means of which, and the equations (3), the expression for D becomes

$$D=\frac{(z-z')}{a'b-ab'}\sqrt{(a-a')^2+(b-b')^2+(a'b-ab')^2},$$

in which, if we substitute the value of $z-z'$ deduced from (2), we obtain, finally,

$$D=\frac{(b-b')(\alpha-\alpha')-(a-a')(\beta-\beta')}{\sqrt{(a-a')^2+(b-b')^2+(a'b-ab')^2}}.$$

If the numerator of this expression vanish, we shall have $D=0$; so that, in this case, the lines will intersect. Indeed, the condition of intersection of the two lines, (1), we know (*Anal. Geom.* p. 225,) to be

$$(b - b')(a - a') = (a - a')(\beta - \beta').$$

2. Among all rectangular prisms to determine that which, having a given volume, shall have the least possible surface.

Representing the three contiguous edges of the prism by x, y, z, and the volume by a^3, we have

$$u = 2xy + 2xz + 2yz = \text{a minimum},$$

but since

$$xyz = a^3 \quad \therefore \; z = \frac{a^3}{xy}$$

$$\therefore \; u = 2xy + \frac{2a^3}{y} + \frac{2a^3}{x} = \text{a minimum}.$$

Therefore we must have the conditions

$$\frac{du}{dx} = 0, \; \frac{du}{dy} = 0,$$

that is,

$$2y - \frac{2a^3}{x^2} = 0, \; 2x - \frac{2a^3}{y^2} = 0,$$

from which we obtain

$$y = x = a \quad \therefore \; z = a.$$

If these values really correspond to a minimum they must fulfil the conditions

$$[\frac{d^2u}{dx^2}] > 0, \; [\frac{d^2u}{dy^2}] > 0, \; [\frac{d^2u}{dx^2}\,\frac{d^2u}{dy^2} - (\frac{d^2u}{dx\,dy})^2] > 0,$$

and these conditions are fulfilled, since

$$[\frac{d^2u}{dx^2}] = 4, \; [\frac{d^2u}{dy^2}] = 4, \; [\frac{d^2u}{dx\,dy}] = 2.$$

Hence the required prism must be a cube.

In the preceding example we might have concluded, without recurring to these conditions, that the results obtained belong to the required minimum, there being obviously no other maximum or minimum, except that which belongs to $x = 0$, $y = 0$, $z = \infty$, these being the values which cause the differential coefficients to become *infinite*, (see *art.* 50.)

3. To divide a given number, a, into three parts, such that the continued product of the mth power of the first part, the nth power of the second part, and the pth power of the third, may be the greatest possible.

The three parts are $\frac{ma}{m+n+p}$, $\frac{na}{m+n+p}$, $\frac{pa}{m+n+p}$, so that the three parts are to each other as the exponents of the proposed powers.

4. To determine the greatest triangle that can be enclosed by a given perimeter.

The triangle must be equilateral.*

On changing the Independent Variable.

(66.) It is frequently requisite to employ the differential coefficients $\frac{dy}{dx}$, $\frac{d^2y}{dx^2}$, &c., in which x is considered as the principal variable under a change of hypothesis, x, and consequently y, being assumed as a function of some new variable t.

It is therefore of consequence to ascertain what changes take place in the expressions for these coefficients in such cases. This we may do as follows:

Since according to the new hypothesis

$$y = F(x) \text{ and } x = f(t),$$

therefore, (33),

$$\frac{dy}{dt} = \frac{dy}{dx}\frac{dx}{dt} \therefore \frac{dy}{dx} = \frac{dy}{dt} \div \frac{dx}{dt} = \frac{(dy)}{(dx)},$$

where, for brevity, (dy) is put for $\frac{dy}{dt}$ and (dx) for $\frac{dx}{dt}$.

$$\frac{d^2y}{dt^2} = \frac{d^2y}{dx^2}\frac{dx^2}{dt^2} + \frac{dy}{dx}\frac{d^2x}{dt^2}$$

---

* For more examples the student may refer to *Jephson's Fluxional Calculus*, to *Garnier's Calcul Differentiel*, or to *Puissant's Problèmes de Géométrie*.

$$\therefore \frac{d^2y}{dx^2} = \left\{\frac{d^2y}{dt^2} - \frac{(dy)}{(dx)}(d^2x)\right\} \div (dx)^2 = \frac{(d^2y)(dx) - (d^2x)(dy)}{(dx)^3}.$$

In a similar manner we might, if necessary, find the expression for $\frac{d^3y}{dx^3}$. It appears, therefore, that

$$\frac{dy}{dx} = \frac{(dy)}{(dx)}$$

$$\frac{d^2y}{dx^2} = \frac{(d^2y)(dx) - (d^2x)(dy)}{(dx)^3},$$

&c. &c.

If $t = y$ the hypothesis requires that y be considered as the principal and x as the dependent variable. In this case

$$(dy) = \frac{dy}{dy} = 1, (d^2y) = 0, \text{ \&c. } (dx) = \frac{dx}{dy}, (d^2x) = -\frac{d^2x}{dy^2}, \text{ \&c.}$$

$$\therefore \frac{dy}{dx} = \frac{1}{\frac{dx}{dy}} = \frac{1}{(dx)}$$

$$\frac{d^2y}{dx^2} = -\frac{(d^2x)}{(dx)^3},$$

&c. &c.

These formulas will be brought into use in the second section

CHAPTER IX.

ON THE CASES IN WHICH TAYLOR'S THEOREM FAILS.

(67.) It has been shown, in Chapter II., that the general development of the function $F(x+h)$ always proceeds according to the ascending positive powers of h, and the principle upon which this fact has been established is this: viz. that $F(x)$ and $F(x+h)$ must necessarily contain the same number of values; or, in other words, the same radicals that enter $F(x)$ must also enter $F(x+h)$, but no others. Hence we might extend the proposition established in (4), and say, that not only the general development proceeds according to the increasing positive powers of h, but also every particular development, provided the particular value $F(a+h)$ contain the same radicals as $F(a)$ and no more; and provided, moreover, that $F(a)$ is not infinite, which we have seen it must be for the true development of $F(a+h)$ to contain a negative power of h.

(68.) As $F(x+h)$ must contain the same radicals as $F(x)$ and no others, it follows that $F(x+h)-F(x)$ must contain the same radicals as $F(x)$. Now, as multiplying or dividing an expression by any *rational*, quantity, can neither introduce nor destroy radicals in that expression, we infer that the expression for $\frac{F(x+h)-F(x)}{h}$, h being rational, must contain the very same radicals as $F(x)$, and no others, whatever be the value of h; but when $h=0$,

$$\frac{F(x+h)-F(x)}{h}=\frac{dF(x)}{dx};$$

hence the first differential coefficient must contain the same radicals as the function $F(x)$, and, by the same reasoning, the second differential coefficient must contain the same radicals as the first; consequently, the same radicals must enter each differential coefficient that enter into the

original function. If, therefore, there be given to x such a particular value a, that any one of the expressions that may be under radicals in $F(x)$ may become 0, that radical will of course vanish from the function, and consequently from its differential coefficients. But if $a + h$ be substituted for x instead of a, the same radical will necessarily be preserved in $F(a + h)$, although it will still vanish from $F(a)$ and the differential coefficients. It follows, therefore, that $F(a + h)$ will have more values than

$$F(a) + [\frac{dF(x)}{dx}]\ h + [\frac{d^2F(x)}{dx}]\ \frac{h^2}{1 \cdot 2} + \&c.$$

so that this cannot be the form of the true development of $F(a + h)$. It is easy to explain why, in such cases as this, one of the coefficients and indeed all that follow this, must become infinite, when $x = a$. For the exponent of the radical, which vanishes for this value, is diminished by unity at each differentiation, and being fractional, the expression under it will at length appear with a *negative* exponent, and will continue to have a negative exponent in all the succeeding coefficients;* so that these, when a is put for x, become infinite. We have observed above, that the failing cases of which we are speaking, arise from the circumstance of $x = a$ causing an expression to disappear, which is under a radical in $F(x)$. The student must not confound this disappearance of a radical, with that which may arise from a *factor* by which it is multiplied becoming 0 for $x = a$; for though a radical may disappear in this way from $F(x)$ it will not disappear from *all* the differential coefficients, and therefore Taylor's development will hold. Thus if a radical in $F(x)$ is multiplied by $(x - a)^m$, m being a positive whole number, this radical will disappear when $x = a$; but in the mth differential coefficient the factor will be $(x - a)^{m-m}$, which does not vanish when $x = a$, but becomes $= 1$, and thus the radical with which it is connected will appear in this coefficient.

---

* This does not invalidate the previous assertion, that the *same* radicals enter the coefficients, that appear in the original function; for the radical still remains, however we increase or diminish the fractional exponent by integers, for $f(x^{\frac{1}{n} \pm p}) = f(\sqrt[n]{x^{1 \pm np}})$.

We conclude, therefore, that there are but two classes of values for which Taylor's development fails; 1° those which, put for x, render $F(x) = \infty$; that is, those which are roots of the equation $\frac{1}{F(x)} = 0$; and 2° those which substituted for x, cause an expression under a radical, to vanish from $F(x)$, and not from $F(x+h)$. To this latter class belongs the value $x = a$, for every function containing $\sqrt{x-a}$, since the value $x = a$ causes this radical to vanish from $F(x)$, but in $F(x+h)$ it enters as $\sqrt{h}$.

(69.) In order to examine these cases more completely, let in general

$$F(a+h) = A + Bh + Ch^2 + Dh^3 + \ldots . Mh^n + Nh^{n+\frac{1}{p}} + \&c. \ldots . (1)$$

represent the true development of $F(x+h)$ for $x = a$, in which $n + \frac{1}{p}$ denotes a fraction falling between the numbers n and $n+1$. We shall show that the $n+1$th differential coefficient derived from the function $F(x)$ becomes infinite for $x = a$, as also all that follows this; but the preceding coefficients are all finite.

Since in the development (1), h has no fixed value, we may differentiate relatively to h, and we shall have

$$\frac{dF(a+h)}{dh} = B + 2Ch + 3Dh^2 + \ldots . + M_1 h^{n-1} + N_1 h^{n+\frac{1}{p}-1} + \&c.$$

$$\frac{d^2F(a+h)}{dh^2} = 2C + 2 \cdot 3Dh + \ldots \ldots . + M_2 h^{n-2} + N_2 h^{n+\frac{1}{p}-2} + \&c.$$

&c &c.

Now, by (30), these several differential coefficients are the same as

$$\left[\frac{dF(x+h)}{dx}\right], \quad \left[\frac{d^2F(x+h)}{dx^2}\right], \ \&c.$$

the brackets denoting the values when $x = a$.

---

* For brevity, M_n will be here used to denote the nth differential coefficient derived from Mh^n.

Hence, by substitution, in the foregoing equations,

$$[\frac{dF(x+h)}{dx}] = B + 2Ch + 3Dh^2 + \ldots . M_1 h^{n-1} + N_1 h^{n+\frac{1}{p}-1}$$
$$+ \&c. \ldots . (2).$$

$$[\frac{d^2F(x+h)}{dx^2}] = 2C + 2 . 3\overset{\frown}{D}h + \ldots \ldots \ldots M_2 h^{n-2} + N_2 h^{n+\frac{1}{p}-2}$$
$$+ \&c. \ldots . (3).$$

&c. &c.

Putting, now, $h = 0$ in the equations (1), (2), (3), &c. we have the following results:

$$F(a) = A$$

$$[\frac{dF(x)}{dx}] = B$$

$$[\frac{d^2F(x)}{dx^2}] = 2\ C$$

$$[\frac{d^3F(x)}{dx^3}] = 2 . 3\ D$$

$$\vdots$$

$$[\frac{d^nF(x)}{dx^n}] = M_n h^{n-n} + N_n 0^{n-n+\frac{1}{p}} + \&c. = M_n$$

$$[\frac{d^{n+1}F(x)}{dx^{n+1}}] = N_{n+1} 0^{\frac{1}{p}-1} + \&c. = \frac{N_{n+1}}{0} = \infty ,$$

&c. &c.

all the succeeding differential coefficients being obviously infinite, because the exponent $\frac{1}{p} - 1$, which is already negative, continually diminishes by unity.

(70.) It follows, therefore, that if the true development of $F(x + h)$, arranged according to the increasing exponents of h, contain for $x = a$

a fractional power of h, comprised between the powers h^n and h^{n+1}, then the several terms of this development will be correctly determined by Taylor's theorem, as far as the term containing h^n inclusively; but the terms beyond this become infinite, and therefore do not belong to the true development.

If a term, in the true development of $F(a+h)$, contain a negative power of h, this should be the leading term, as the arrangement is according to the increasing exponents; therefore, this first term, when $x=a$ and $h=0$, must be infinite, and consequently all the differential coefficients, (2), (3), &c. must be infinite.

(71.) The converse of these inferences are true, viz. 1°. If, in the general development of $F(x+h)$, the coefficient $\frac{d^{n+1}F(x)}{dx^{n+1}}$ is the first which becomes infinite for a particular value of x, then, in the true development, arranged according to the increasing exponents of h, the term immediately succeeding that which contains h^n, will contain a fractional power of h, the exponent being between n and $n+1$.* For it is obvious, that in order that the $n+1$th may be the first of the coefficients (2), (3), &c. which contain a negative power of h, $h^{n+\frac{1}{p}}$ must be the first fractional power of h which enters the development (1), $n+\frac{1}{p}$ being between n and $n+1$. 2°. If, for a particular value of x, the function $F(x)$ become infinite, then will all the differential coefficients become also infinite; and the true development will

* This is of course on the supposition that the development is possible according to *some* powers or roots of the increment h, with finite coefficients; but when no such development is possible, then the appearance of

$$\left[\frac{d^{n+1}F(x)}{dx^{n+1}}\right]=\infty$$

will indicate that the coefficient of h^{n+1} will itself involve some transcendental function of h not developable according to any powers or roots of h with finite coefficients.

contain a negative power of h, or else a log h, a cot h, &c. For, if the true development of $F(a+h)$ did not contain a negative power of h, nor a log h, a cot h, &c. $F(a)$, which this becomes when $h=0$, could not be infinite; hence, such a function of h must enter, and therefore, as shown above, all the differential coefficients become infinite, for $x=a$. It is, therefore, necessary to examine the function $F(a)$, before we deduce the coefficients from $F(x)$.

(72.) To obtain the true development of the function for those particular values of the variable, which cause Taylor's theorem to fail, the usual course is to recur to the ordinary processes of common algebra, after having substituted $a+h$ for x in $F(x)$.

Suppose, for example, the function were

$$F(x)=2ax-x^2+a\sqrt{x^2-a^2},$$

and that we required the development of $F(x+h)$, for $x=a$.

Taking the differential coefficient, we have

$$\frac{dy}{dx}=2(a-x)+\frac{ax}{\sqrt{x^2-a^2}} \quad \therefore \left[\frac{dy}{dx}\right]=\infty,$$

&c. &c.

As, therefore, the first differential coefficient becomes infinite for the proposed value of x, we conclude that the true development of the function for that value, when arranged according to the increasing exponents of h, has a fractional power of h in the second term, the exponent of this power being between 0 and 1; for it is plain that no transcendental function of h can enter.

Substituting, then, $a+h$ for x in $F(x)$, we have

$$\begin{aligned} F(a+h) &= a^2-h^2+a\sqrt{2ah+h^2} \\ &= a^2-h^2+ah^{\frac{1}{2}}(2a+h)^{\frac{1}{2}}. \end{aligned}$$

Developing $(2a+h)^{\frac{1}{2}}$ by the binomial theorem, we have

$$(2a+h)^{\frac{1}{2}}=(2a)^{\frac{1}{2}}+\frac{h}{2(2a)^{\frac{1}{2}}}-\frac{h^2}{8(2a)^{\frac{3}{2}}}+\&c.$$

consequently,

$$F(a+h) = a^2 + a\,(2a)^{\frac{1}{2}}h^{\frac{1}{2}} + \frac{h^{\frac{3}{2}}}{2\,(2a)^{\frac{1}{2}}} - h^2 - \frac{h^{\frac{5}{2}}}{8\,(2a)^{\frac{3}{2}}} + \&c.$$

Again, let the function be

$$F(x) = \sqrt{x} + (x-a)^2 \log(x-a),$$

and let it be required to find the development of $F(a+h)$. Here

$$F(a) = \sqrt{a} + 0 \times \infty.$$

It becomes necessary, therefore, first to ascertain whether this expression is infinite; for, if it be, we are not to proceed with the differentiation, but to infer, agreeably to art. (70), that the proposed function, and all the differential coefficients, become infinite for $x = a$, and that consequently the true development contains either a negative power of h, or a logarithm of h. Now, by the method explained in (44), we find that, when $x = a$, the true value of

$$(x-a)^2 \log(x-a) = \frac{\log(x-a)}{\frac{1}{(x-a)^2}}$$

is 0. Hence the first term of the development is $\sqrt{a}$, so that the development holds thus far. Differentiating the expression for $F(x)$, we have

$$\frac{dF(x)}{dx} = \frac{1}{2\sqrt{x}} + 2\,(x-a)\log(x-a) + x - a,$$

which, for $x = a$, becomes $\frac{1}{2\sqrt{a}}$; so that the true development of $F(x+h)$ agrees with Taylor's form, as far, at least, as the first two terms. Differentiating again, we have

$$\frac{d^2F(x)}{dx^2} = -\frac{1}{4x\sqrt{x}} + 2\log(x-a) + 3,$$

which, for $x = a$, evidently becomes infinite, inasmuch as $\log 0 = -\infty$. Hence we infer that the true development differs from Taylor's form at

the third term. The first three terms are as follows:

$$(a+h)^{\frac{1}{2}} + h^2 \log h = a^{\frac{1}{2}} + \frac{1}{2}\, a^{-\frac{1}{2}} h + (\log h - \frac{1}{2 \cdot 4}\, a^{-\frac{3}{2}})\, h^2 + \&c.$$

The third term involves $\log h$, which we know is not capable of development according to the powers of h, (see p. 38).

SCHOLIUM.

(73.) In the preceding remarks on the development of functions for particular values of the variable, we have said nothing about the values of h, the increment of that variable, having indeed considered that increment as indeterminate, or rather of arbitrary value. It must, however, be observed that, although the particular value which we give to x does not, in any case, fix the value of h, it may nevertheless fix the limit between which and 0 all the values given to h must be comprised, in order that for particular values of x, Taylor's development may not fail.* This fact is very plain, for, if the development holds for all values of x from $x = a$ up to $x = b$, but fails for $x = b$, in consequence of a differential coefficient becoming infinite, then will the development hold when $a + h$ is substituted for x in $\mathrm{F}(x)$, provided h be taken between the limits $h = 0$ and $h = b - a$, or, more strictly, provided it does not exceed these limits, since throughout this interval $[\mathrm{F}(x)]$ and $[\mathrm{F}(x+h)]$ necessarily involve the same radicals. In like manner, if the development hold for all values of x from $x = b$ down to $x = a$, but fail for $x = a$, then will the development hold when $b - h$ is substituted for x for all values of b from $h = 0$, to $h = b - a$, but it will not hold for the value of h immediately succeeding this last; and it is obvious that h will always be subject to such restrictions unless the development holds, not merely for $x = a$, but universally. When, therefore, we find that for $x = a$ the differential coefficients do not any

* It must be remembered that for Taylor's theorem to hold for $x = a$, $\mathrm{F}(a)$ and $\mathrm{F}(a+h)$ must contain the same radicals.

of them become infinite, all that we can conclude is that the development of $F(a \pm h)$ is according to Taylor's theorem for all values of h between some certain finite value h', which may indeed be indefinitely small, and 0, and it is only when this is not the case that the theorem is said by analysts to *fail*. We have thought it necessary to point out these circumstances to the student, seeing that some authors, from not attending to them, have fallen into very important errors, and have laid down erroneous doctrines with respect to the failing cases of Taylor's theorem. Thus Mr. Jephson, at page 191, vol. i. of his Fluxional Calculus,* a work containing much valuable information, says, "It may further be observed that Taylor's theorem always fails when the assigned value of x causes any of the terms to become imaginary, and that this may take place without causing the function itself to be imaginary; thus take $fx = c + x^2\sqrt{x-a}$ if we suppose $x = 0$, $fx = c$, $f'x = 0$, but $f''x$, $f'''x$ . . . . all contain $\sqrt{-1}$." From this it would appear that Taylor's theorem may fail to give the true development in other cases besides those which cause the differential coefficients to become infinite, which, however, is not true. Whenever, for any particular value of x, Taylor's coefficients become imaginary, we must infer, agreeably to the statement in (4), that the function $F(x+h)$ becomes imaginary for that value of x; h being of course limited, as above explained. In the example just quoted, where

$$f'x = 2x\sqrt{x-a} + \frac{\frac{1}{2}x^2}{\sqrt{x-a}},$$

$$f''x = 2\sqrt{x-a} + \frac{2x}{\sqrt{x-a}} - \frac{\frac{1}{4}x^2}{(x-a)^{\frac{3}{2}}},$$

&c. &c.

the function $f(x+h)$ becomes, when $x = 0$, $f(0+h) = c + h^2\sqrt{h-a}$, and the development is

---

* See note (E) at the end.

$$c + h^2\sqrt{h-a} = c + 0h + \sqrt{-a}\,h^2 + \frac{1}{2\sqrt{-a}}h^3 + \&c.,$$

and this is the true development, for h must not exceed the limits $h = 0$, and $h = a$, since $x = a$ causes the differential coefficient to become infinite, and therefore the development to fail.

With regard to the failing cases of Maclaurin's theorem, it may be observed that they are very different from the failing cases of Taylor's. Whatever be the form of the proposed function, its *general* development, according to Taylor's theorem, never fails; but the failure of Maclaurin's theorem always arises from the *form* of the proposed function, and it is the *general* development that fails, and consequently all the particular cases. For it is obvious that every function will fail to be developable by Maclaurin's theorem, whose form is such that, for $x = 0$, it, or any of the differential coefficients derived from it, becomes infinite.

Before terminating these remarks it may be proper to observe that the student is not to attribute what analysts have been pleased to term the *failing cases* of these theorems to any defect in the theorems themselves; on the contrary they would be very defective if they did not exhibit such cases. All that is meant is, that the function in *particular states* may fail to be developable according to Taylor's series, and under *particular forms* it may fail to be developable according to Maclaurin's series; so that, in fact, these theorems fail to give the true development only when that development is impossible.

(74.) Let us now examine implicit functions, and let us suppose that $x = a$ causes a radical to vanish from $F(x)$ in consequence of a factor of it vanishing; we have seen (68) that such radical will reappear in some of the differential coefficients; suppose it appears in the first, which requires that the factor spoken of be $x - a$, then for $x = a$ this coefficient will have more values than the proposed function, as it contains a radical more. But if the function that $F(x)$ or y is of x be only implicitly given, that is by means of an equation without radicals, we know (52) that the expression for $\frac{dy}{dx}$ will be also without radicals; and from

such an expression it does not at first seem clear how we are to deduce the multiple values alluded to, and which might be obtained by solving the equation for y, and thus introducing the radical. But since the expression for $\frac{dy}{dx}$ appears under the form of a fraction, viz. (52),

$$\frac{dy}{dx} = -\frac{\frac{du}{dx}}{\frac{du}{dy}} \quad \ldots . \quad (1),$$

we readily perceive that one case is possible, and only one, in which this fraction may take a multiplicity of values besides those *implied* in y, viz. the case in which it becomes $\frac{0}{0}$; that is, when the following conditions exist simultaneously, viz.

$$\frac{du}{dx} = 0, \ \frac{du}{dy} = 0 \quad \ldots . \quad (2);$$

so that these conditions are those which must exist for every value of x which destroys a radical in $F(x)$ but not in $\frac{dy}{dx}$.

(75.) Whenever, therefore, any particular value of x destroys a radical in y, but not in $\frac{dy}{dx}$, then the expression (1) must take the form $\frac{0}{0}$, to admit of the necessary multiple values.

The rules laid down in (41) enable us to determine the true value of the fraction (1) in the proposed circumstances, that is, when it takes the form $\frac{0}{0}$, for there is but one independent variable, viz. x. By differentiating numerator and denominator separately, we have, by the article referred to,

$$\left[\frac{dy}{dx}\right] = -\left[\frac{\frac{d^2u}{dx^2} + \frac{d^2u}{dx\,dy}\frac{dy}{dx}}{\frac{d^2u}{dx\,dy} + \frac{d^2u}{dy^2}\frac{dy}{dx}}\right] \quad \ldots . \quad (3);$$

hence,

$$[\frac{d^2u}{dx^2} + 2\,\frac{d^2u}{dx\,dy}\,\frac{dy}{dx} + \frac{d^2u}{dy^2}\,(\frac{dy}{dx})^2] = 0.$$

This being a quadratic equation furnishes *two* values for $[\frac{dy}{dx}]$, which are all that belong to the fraction (1) or to $[\frac{dy}{dx}]$, unless, indeed, both numerator and denominator of (3) also vanish for a value of x given by the conditions (2), in which case we must differentiate again as in art. (41), when the value of $[\frac{dy}{dx}]$ will be given by an equation of the third degree which will furnish *three* values, and so on; and in general, if the fraction, or $[\frac{dy}{dx}]$, admit of n values, the equation which determines them can be obtained only by differentiating n times, which will lead to an equation of the nth degree, and it is plain that the radical destroyed in $F(x)$ must be of the $\frac{1}{n}$th degree, seeing that it gives to $\frac{dy}{dx}$, n additional values.

Suppose, for example, we had the function

$$y = x + (x-a)\sqrt{x-b},$$

$$\therefore \frac{dy}{dx} = 1 + \sqrt{x-b} + \frac{x-a}{2\sqrt{x-b}},$$

and for $x = a$,

$$[y] = a,\ [\frac{dy}{dx}] = 1 \pm \sqrt{a-b};$$

so that the radical which disappears in y appears in $\frac{dy}{dx}$; this, therefore, has *two* values.

Now let the same function be given in an implicit form and without radicals, viz.

$$u=(y-x)^2-(x-a)^2(x-b)=0,$$

$$\therefore \frac{du}{dx}=-2(y-x)-(x-a)(3x-2b+a), \frac{du}{dy}=2(y-x).$$

Since for $x=a$, $y=a$, therefore both these expressions become 0; hence $[\frac{dy}{dx}]=\frac{0}{0}$. Taking, then, the differentials of both numerator and denominator of the fraction,

$$\frac{2(y-x)+(x-a)(3x-2b+a)}{2(y-x)}=\frac{0}{0},$$

we find when $x=a$ that it becomes

$$[\frac{dy}{dx}]=1+\frac{a-b}{[\frac{dy}{dx}]-1},$$

therefore,

$$[\frac{dy}{dx}-1]^2-(a-b)=0 \quad \therefore \quad [\frac{dy}{dx}]=1\pm\sqrt{a-b},$$

as before.

(76.) Suppose now that the radical which disappears from $F(r)$, by reason of a factor, disappears also from $\frac{dy}{dx}$, and that it appears in $\frac{d^2y}{dx^2}$, which is the same as supposing that the vanishing factor is $(x-a)^2$. In this case $\frac{dy}{dx}$ will have the same number of values that $F(x)$ has for $x=a$, but additional values will belong to $\frac{d^2y}{dx^2}$, so that we must have

$$[\frac{d^2y}{dx^2}]=\frac{0}{0};$$

and the true values, when the function is implicit, will all be determined by the principles already employed.

For example: the explicit function

$$y = x + (x-a)^2 \sqrt{x}$$

gives when $x = a$,

$$[y] = a, \ [\frac{dy}{dx}] = 1, \ [\frac{d^2y}{dx^2}] = \pm 2\sqrt{a},$$

and we shall see that the same values are equally given by the implicit function

$$(y-x)^2 = (x-a)^4 x,$$

by applying the foregoing process to $\frac{d^2y}{dx^2}$.

For by successively differentiating, and representing, for brevity, the several coefficients derived from y by p', p'', &c. we have

$$2(p'-1)(y-x) = (x-a)^3(5x-a) \therefore [p'] = 1$$

$$(p'-1)^2 + p''(y-x) = 2(x-a)^2(5x-2a)$$

$$\therefore [p''] = [\frac{2(x-a)^2(5x-2a) - (p'-1)^2}{y-x}] = \frac{0}{0}$$

$$= [\frac{6(x-a)(5x-3a) - 2p'(p'-1)}{p'-1}] = \frac{0}{0}$$

$$= [\frac{60x - 48a - 2p'''(y'-1) - 2p''^2}{p''}] = \frac{12a - 2[p''^2]}{[p'']}$$

$$\therefore 3[p''^2] = 12a \therefore [p''] = \pm 2\sqrt{a},$$

as before.

The two examples following will suffice to exercise the student in this doctrine, which is merely an extension of the principles treated in Chapter V. to implicit functions.

1. Given

$$y^3 = (x-a)^3(x-b),$$

to determine the values of $\frac{dy}{dx}$, when $x = a$,

$$[\frac{dy}{dx}] = \sqrt[3]{a-b}.$$

2. Given

$$(y-x)^2 - (x-a)^4 (x-b) = 0$$

to determine the values of $\frac{dy}{dx}$ and $\frac{d^2y}{dx^2}$, when $x = a$.

$$[\frac{dy}{dx}] = 1, \; [\frac{d^2y}{dy^2}] = 2\sqrt{a-b}.$$

We here terminate the First Section, having fully considered the various particulars relating to the differentiation of functions in general.

SECTION II.

APPLICATION OF THE DIFFERENTIAL CALCULUS TO THE THEORY OF PLANE CURVES.

CHAPTER I.

ON THE METHOD OF TANGENTS.

(77.) We now proceed to apply the Calculus to Geometry, and shall first explain the method of drawing tangents to curves.

The general equation of a secant passing through two points (x', y'), (x'', y''), in any plane curve, is (*Anal. Geom.* p. 197),

$$y - y' = \frac{y' - y''}{x' - x''}(x - x'),$$

$y' - y''$, being the increment of the ordinate or proposed function corresponding to $x' - x''$, the increment of the abscissa or independent variable. The *limit* of the ratio of these increments, by the principles of the calculus, is $\frac{dy'}{dx'}$; that is to say, such is the representation of the ratio or quotient when $x' - x'' = 0$, and, consequently, $y' - y'' = 0$. But when this is the case the secant merges into the tangent. Hence the equation of the tangent, through any point (x', y') of the plane curve, is

$$y - y' = \frac{dy'}{dx'}(x - x') \quad \ldots \quad (1).$$

It appears, therefore, that the differential coefficient $\frac{dy'}{dx'}$, for any pro-

posed point in the curve, has for its value the trigonometrical tangent of the angle included by the linear tangent and the axis of x, that is, provided the axes are rectangular. If the axes are oblique, the same coefficient represents the ratio of the sines of the inclinations of the linear tangent to these axes. (*See Anal. Geom.* p. 19.)

By means of the general equation (1) we can always readily determine the equation of the tangent to any proposed plane curve when the equation of the curve is given, nothing more being required than to determine from that equation the differential coefficient.

Suppose, for example, it were required to find the particular form for the ellipse. We here have to determine $\frac{dy'}{dx'}$ from the equation

$$Ay'^2 + Bx'^2 = A^2B^2,$$

and which is

$$\frac{dy'}{dx'} = -\frac{B^2x'}{A^2y'};$$

therefore the equation of the tangent is

$$y - y' = -\frac{B^2x'}{A^2y'}(x - x'),$$

(x', y') being any point in the curve, and (x, y) any point in the tangent.

Again let it be required to determine the general equation of the tangent to a line of the second order.

By differentiating the general equation

$$Ay'^2 + Bx'y' + Cx'^2 + Dy' + Ex' + F = 0,$$

we have

$$2Ay'\frac{dy'}{dx'} + Bx'\frac{dy'}{dx'} + By' + 2Cx' + D\frac{dy'}{dx'} + E = 0,$$

$$\therefore \frac{dy'}{dx'} = -\frac{2Cx' + By' + E}{2Ay' + Bx' + D};$$

so that the general equation is

$$y - y' = -\frac{2Cx' + By' + E}{2Ay' + Bx' + D}(x - x').$$

(78.) As the normal is always perpendicular to the tangent, its general equation must be, from (1),

$$y - y' = -\frac{1}{\frac{dy'}{dx'}}(x - x') \quad \ldots \quad (2).$$

It is easy now to deduce the expressions for the subtangent and subnormal. For if, in the equation of the tangent, we put $y = 0$, the resulting expression for $x - x'$ will be the analytical value of that part of the axis of x intercepted between the tangent and the ordinate y' of the point of contact, that is to say, it will be the value of the subtangent $T_{,}$ (*Anal. Geom.* p. 84),

$$\therefore T_{,} = \frac{-y'}{\frac{dy'}{dx'}} \quad \ldots \quad (3).$$

If, instead of the equation of the tangent, we put $y = 0$ in that of the normal, then the resulting expression for $x - x'$ will be the value of the intercept between the normal and the ordinate y', that is, it will belong to the subnormal $N_{,}$,

$$\therefore N_{,} = y' \frac{dy'}{dx'} \quad \ldots \quad (4).$$

As to the length of the tangent T, since $T = \sqrt{y'^2 + T_{,}^2}$, we have, in virtue of (3),

$$T = y' \sqrt{1 + \frac{1}{(\frac{dy'}{dx'})^2}} \quad \ldots \quad (5).$$

Also, since the length of the normal, N, is $N = \sqrt{y'^2 + N'^2}$, we have, by (4),

$$N = y' \sqrt{1 + (\frac{dy'}{dx'})^2} \quad \ldots \quad (6).$$

The foregoing expressions evidently apply to any plane curve whatever; that is, to any curve that may be represented by an equation between two variables, whatever be its degree, or however complicated its form.

We shall now give a few examples principally illustrative of the method of drawing tangents to the higher curves, for which purpose we shall obviously require only the formula (3), for it is plain that to any point in a curve we may at once draw a tangent, when the length and position of the subtangent is determined. Or, knowing the point (x', y'), we may, by putting successively $x = 0$ and $y = 0$, in the equation (1), determine the two points in which the required tangent ought to cut the axes, and then draw it through them. If $\frac{dy'}{dx'}$ is 0 at the proposed point, the tangent will be parallel to the axis of x, because, as remarked above, $\frac{dy'}{dx'}$ is the value of the trigonometrical tangent of the inclination of the linear tangent to the axis of x, and for this reason also the tangent will be parallel to the axis of y when $\frac{dy'}{dx'}$ is infinite at the proposed point.

EXAMPLES.

(79.) 1. To draw a tangent to a given point P in the common or conical parabola.

By the equation of the curve

$$y^2 = px,$$

$$\therefore \frac{dy}{dx} = \frac{p}{2y};$$

hence, at the point (x', y'), we have

$$T_1 = y' \div \frac{p}{2y'} = \frac{2y'^2}{p} = 2x'.$$



Hence, having drawn from P, the perpendicular ordinate PM, if we set

off the length, MR, on the axis of x, equal to twice AM, and then draw the line RP, it will be the tangent required.

2. To determine the subtangent and subnormal at a given point (x', y') in the parabola of the nth order, represented by the equation

$$y = ax^n,$$

$$\therefore \frac{dy}{dx} = nax^{n-1},$$

$$\therefore T_{,} = y' \div nax'^{n-1} = \frac{x'}{n}, \quad N_{,} = y' \cdot nax'^{n-1} = na^2x'^{2n-1} \text{ or } \frac{ny'^2}{x'}.$$

3. To determine the subtangent at a given point in the logarithmic, curve.

The equation of this curve related to rectangular coordinates is

$$y = a^x,$$

which shows that if the abscissas x be taken in arithmetical progression, the corresponding ordinates y will be in geometrical progression; so that the ordinates of this curve will represent the numbers, the logarithms of which are represented by the corresponding abscissas, a being the assumed base of the system. Hence, calling the modulus of this base m, we have, by differentiating (13),

$$\frac{dy}{dx} = \frac{1}{m} y,$$

$$\therefore T_{,} = y' \div \frac{y'}{m} = m.$$

Hence the remarkable property that the subtangent is constantly equal to the modulus of the system, whose base is a.

4. To determine the subtangent at a given point in the curve whose equation is

$$x^3 - 3axy + y^3 = 0.$$

Here

$$\left\{\frac{du}{dx}\right\} = 3x^2 - 3ay - 3ax\frac{dy}{dx} + 3y^2\frac{dy}{dx} = 0,$$

$$\therefore \frac{dy}{dx} = \frac{ay - x^2}{y^2 - ax},$$

Hence, at the point (x', y'),

$$T_{,} = \frac{y'^3 - ax'y'}{ay' - x'^2}.$$

5. To draw a tangent to a rectangular hyperbola between the asymptotes, its equation being $xy = a$.

$$T_{,} = x'.$$

6. To determine the subtangent at a given point in a curve whose equation is $y^m = ax^n$, which, because it includes the common parabola, is said to represent parabolas of all orders.

$$T_{,} = \frac{m}{n} x'.$$

7. To determine the subtangent at a given point in a curve whose equation is $x^m y^n = a$, which, because it includes the common hyperbola, is said to belong to hyperbolas of all orders.

$$T_{,} = \frac{n}{m} x'.$$

(80.) If the proposed curve be related to polar coordinates, then the expressions in last article must be changed into functions of these.



Let the curve AP be related to polar coordinates $FP = r$, $PFX = \omega$; then if PR be a tangent at any point, P, and PN the normal, and if RFN be perpendicular to the radius vector FP, the part FR will be the *polar subtangent*, and the part FN the *polar subnormal*. When the pole F and the point P are given, it is obvious that the determination of the subtangent FR will enable us to draw the tangent PR.

The formulas for transforming the equation of a curve from rectangular to polar coordinates, having the same origin, are (*Anal. Geom.*, p. 131,)

$$x = r\cos\omega, \quad y = r\sin\omega, \quad x^2 + y^2 = r^2,$$

and the resulting equation of the curve will have the form

$$r = F(\omega), \quad \text{or } F(r, \omega) = 0,$$

in which we shall consider ω as the independent variable. Now $RF = PF \cdot \tan \angle P = r \tan \angle P$; but, Trigonometry, page 48,

$$\tan \angle P = \frac{\tan \omega - \tan \alpha}{1 + \tan \omega \tan \alpha};$$

that is, since

$$\tan \alpha = \frac{dy}{dx}, \text{ and } \tan \omega = \frac{y}{x},$$

$$\tan \angle P = \frac{\frac{y}{x} - \frac{dy}{dx}}{1 + \frac{y}{x}\frac{dy}{dx}};$$

therefore r times this expression is the value of the polar subtangent. But the differential coefficient $\frac{dy}{dx}$, which implies that x is the principal variable, ought to become, when the variable is changed to ω, (66),

$$\frac{dy}{dx} = \frac{dy}{d\omega} \div \frac{dx}{d\omega} \therefore \tan \angle P = \frac{y\frac{dx}{d\omega} - x\frac{dy}{d\omega}}{x\frac{dx}{d\omega} + y\frac{dy}{d\omega}}.$$

Now, from the above formulas of transformation.

$$\frac{dx}{d\omega} = \frac{dr}{d\omega}\cos\omega - r\sin\omega, \quad \frac{dy}{d\omega} = \frac{dr}{d\omega}\sin\omega + r\cos\omega,$$

$$\therefore y\frac{dx}{d\omega} = r\frac{dr}{d\omega}\sin\omega\cos\omega - r^2\sin^2\omega \ldots\ldots (1).$$

$$x\frac{dy}{d\omega} = r\frac{dr}{d\omega}\sin\omega\cos\omega + r^2\cos^2\omega \ldots\ldots (2).$$

$$x\frac{dx}{d\omega} = r\frac{dr}{d\omega}\cos^2\omega - r^2\sin\omega\cos\omega \ldots\ldots (3).$$

$$y\frac{dy}{d\omega} = r\frac{dr}{d\omega}\sin^2\omega + r^2\sin\omega\cos\omega \ldots\ldots (4).$$

Subtracting (2) from (1), we get the numerator of the foregoing expression for $\tan \angle P$ equal to $-r^2$; and adding (3) and (4), we get the

denominator equal to $r\frac{dr}{d\omega}$; consequently,

$$\tan \angle P = \frac{-r}{\frac{dr}{d\omega}} \therefore RF = \frac{r^2}{\frac{dr}{d\omega}} = \text{subtangent.}$$

Also

$$FN = \frac{FP^2}{FR} = r^2 \div \frac{r^2}{\frac{dr}{d\omega}} = \frac{dr}{d\omega} = \text{subnormal.}$$

It may be proper to observe here that in our figure the tangent from P falls to the left of PF, thus making $\angle P = \omega - a$. If, on the contrary, the tangent fall on the other side, then we should have $\angle P = a - \omega$, in which case

$$\tan \angle P = \frac{r}{\frac{dr}{d\omega}}.$$

(81.) We shall apply these formulas to *spirals*, a class of curves always best represented by polar equations.

8. To determine the subtangent at any point in the *Logarithmic Spiral*, its equation being

$$r = a^{\omega}$$

$$\therefore \frac{dr}{d\omega} = \log a \cdot a^{\omega} = \frac{a^{\omega}}{m},$$

$$\therefore r^2 \div \frac{dr}{d\omega} = mr = T.$$



Hence, if a represent the base of the Napierian system, since the modulus will be 1, the subtangent will be equal to the radius vector, and therefore the angle P will at every point of the curve be equal to 45°, because tan $\angle P = 1$.

Since, by the equation of this curve, $\log r = \omega \log a$, it follows that, if a denote the base of any system, the various values of the angle, or circular arc ω, will denote the logarithms of the numbers represented by the corresponding values of r. Hence, the propriety of the name *logarithmic spiral.*

In this curve

$$\tan \angle \mathrm{P} = r \div \frac{\mathrm{d}r}{\mathrm{d}\omega} = a^{\omega} \div \frac{a^{\omega}}{m} = m;$$

hence the curve cuts all its radii vectores under the same angle; and it is therefore sometimes called the *equiangular* spiral.

9. To determine the subtangent at any point in the *Spiral of Archimedes*, its equation being

$$r = a\omega$$

$$\therefore \frac{\mathrm{d}r}{\mathrm{d}\omega} = a \therefore r^2 \div \frac{\mathrm{d}r}{\mathrm{d}\omega} = a\omega^2 = r\omega = \mathrm{T}_{,},$$

so that FR is equal to the length of the circular arc to radius r, comprehended between FR, FA; when, therefore, $\omega = 2\pi$, the subtangent equals the length of the whole circumference. The spiral of Archimedes belongs to the class of spirals represented by the general equation

$$r = a\omega^{n}.$$

When $n = -1$, we have $r\omega = a$, and the spiral represented is called the *hyperbolic spiral*, on account of the analogy between this equation and $xy = a$. It is also called the *reciprocal spiral*.

10. To determine the polar subtangent at any point in the hyperbolic spiral.

$$\mathrm{T}_{,} = a.$$

11. To determine the polar subtangent at any point in the spiral whose equation is $r = a\omega^{-\frac{1}{2}}$.

$$\mathrm{T}_{,} = 2a\omega^{\frac{1}{2}} = \frac{2a^2}{r}.$$

12. To determine the polar subtangent at any point in the *parabolic spiral*, its equation being $r = a\omega^{\frac{1}{2}}$.

$$\mathrm{T}_{,} = \frac{2r^3}{a^2}.$$

13. To determine the polar subtangent at any point in a spiral whose equation is

$$r = a\omega^{n}.$$

$$\mathrm{T}_{,} = \frac{r^{\frac{n+1}{n}}}{na^{\frac{1}{n}}}.$$

(82.) In these examples the expression for the polar subtangent only is employed; but that for the polar subnormal, deduced immediately from the former at page 150, is also of use. It serves to furnish us very readily with the length of the perpendicular drawn from the pole of the curve to the tangent through any point (r, ω), and, as this is an element frequently introduced into enquiries connected with Physical Astronomy, we shall here show how it may be obtained.

Calling this perpendicular p, it is obvious, from similar triangles, that

$$\mathrm{PN} : \mathrm{PF} :: \mathrm{PF} : p,$$

$$\therefore p = \frac{\mathrm{PF}^2}{\mathrm{PN}}.$$

But $\mathrm{PN} = \sqrt{\mathrm{PF}^2 + \mathrm{FN}^2}$, hence, introducing the expression for FN,

$$p = \frac{r^2}{\sqrt{r^2 + (\frac{\mathrm{d}r}{\mathrm{d}\omega})^2}},$$

$$\text{or } p^2 = \frac{r^4}{r^2 + (\frac{\mathrm{d}r}{\mathrm{d}\omega})^2}.$$

In the three curves of the second order, when the focus is the pole, we have, (*Anal. Geom.* p. 137),

$$r = \frac{2m}{1 + e\cos\omega},$$

from which $\frac{\mathrm{d}r}{\mathrm{d}\omega}$, being obtained and substituted in the expression for p^2, will give

$$p^2 = \frac{4m^2r}{4m - r(1 - e^2)},$$

which, when $e = 1$, that is, when the curve is a parabola, becomes $p^2 = mr$, where m is the distance, FA, of the focus from the vertex. Hence we have, in the parabola,

$$p^2 = \mathrm{FP} \cdot \mathrm{FA}.$$

When $e < 1$, that is, when the curve is an ellipse, in which case $m = \frac{2B^2}{A}$, and $1 - e^2 = \frac{B^2}{A^2}$, we have

$$p^2 = \frac{B^2 r}{2A - r} = \frac{BC^2 . FP}{F'P}.$$

When $e > 1$, that is when the curve is an hyperbola, and consequently $1 - e^2 = -\frac{B^2}{A^2}$, then

$$p^2 = \frac{B^2 r}{2A + r} = \frac{BC^2 . FP}{F'P}.$$

Rectilinear Asymptotes.

(83.) *A rectilinear asymptote* to a curve may be regarded as a tangent of which the point of contact is infinitely distant, so that the determination of the asymptote reduces to the determination of the tangent on the hypothesis that either or both $y' = \infty$, $x' = \infty$, the portions of the axes between the origin and this tangent being, at the same time, one or both finite.

The equation of the tangent being

$$y - y' = \frac{dy'}{dx'}(x - x'),$$

we have, by making successively $y = 0$, $x = 0$, the following expressions for the corresponding parts of the axes of x and y, between the tangent and the origin, viz.

$$x' - \frac{y'}{\frac{dy'}{dx'}}, \text{ and } y' - x'\frac{dy'}{dx'} \quad \ldots \quad (1).$$

If for $x' = \infty$ both these are finite, they will determine two points, one on each axis, through which an asymptote passes. If for $x = \infty$ the first expression is finite and the second infinite, the first will determine a point on the axis of x, and the second will show that a line

through this point, and parallel to the axis of y is an asymptote. If, on the contrary, the second expression is finite, and the first infinite, the asymptote will pass through the point in the axis of y, determined by the finite value, and will be parallel to the axis of x.

When, however, asymptotes parallel to the axes exist, they may generally be detected by merely inspecting the equation, as it is only requisite to ascertain for what values of x, y becomes infinite, or for what values of y, x becomes infinite. Thus, in the equation $xy = a$, $x = 0$ renders $y = \infty$, and $y = 0$ renders $x = \infty$, therefore the two axes are asymptotes. Again, in the equation

$$a^2y^2 - y^2x^2 - bx^4 = 0, \text{ or } y^2 = \frac{bx^4}{a^2 - x^2},$$

it is plain that $x = \pm a$ renders $y = \infty$, we infer, therefore, that the curve represented by this equation has two asymptotes, each parallel to the axis of y, and at the distance a from it.

If both expressions are infinite, there will be no asymptote corresponding to $x = \infty$.

If both expressions are 0, the asymptote will pass through the origin, and its inclination θ, to the axis of x will be determined by $\frac{dy}{dx} = \tan \theta$.

If for $y' = \infty$ one or both of the above general expressions are finite, there will be an asymptote, and its position may be determined as in the foregoing cases.

EXAMPLES.

(84.) 1. Let the curve be the common hyperbola, of which the equation is

$$y = \frac{b}{a}\sqrt{x^2 - a^2},$$

$$\therefore \frac{dy}{dx} = \frac{bx}{a\sqrt{x^2 - a^2}};$$

hence the general expressions (83) are

$$x' - \frac{x^2 - a^2}{x} = \frac{a^2}{x}$$

and

$$-\frac{b}{a}\frac{a^2}{\sqrt{x^2-a^2}}=-\frac{ba}{x\sqrt{1-\frac{a^2}{x^2}}},$$

both of which are 0, when $x=\infty$; hence an asymptote passes through the origin.

Also

$$\frac{dy}{dx}=\pm\frac{b}{a}\cdot\frac{1}{\sqrt{1-\frac{a^2}{x^2}}},$$

which becomes $\pm\frac{b}{a}$, when $x=\infty$, therefore, this being the tangent of the inclination of the asymptotes to the axis of x, they are both represented by the equation

$$y=\pm\frac{b}{a}x.$$

2. To prove that the hyperbola is the only curve of the second order that has asymptotes.

The general equation of a line of the second order, when referred to the principal diameter and tangent through the vertex as axes, is

$$y^2=mx+nx^2,$$

$$\therefore x-\frac{y}{\frac{dy}{dx}}=x-\frac{2y^2}{m+2nx}=\frac{mx+2nx^2-2y^2}{m+2nx}=-\frac{mx}{m+2nx},$$

$$y-x\frac{dy}{dx}=y-\frac{mx+2nx^2}{2y}=\frac{2y^2-mx-2nx^2}{2y}=\frac{mx}{2\sqrt{mx+nx^2}}.$$

Dividing numerator and denominator of each of these expressions by x, they reduce to

$$-\frac{m}{\frac{m}{x}+2n} \quad \text{and} \quad \frac{m}{2\sqrt{\frac{m}{x}+n}};$$

and these, when $x=\infty$, or indeed when $y=\infty$, become

$$-\frac{m}{2n} \quad \text{and} \quad \frac{m}{2\sqrt{n}}.$$

Hence the curve will have asymptotes, provided n be neither 0 nor negative, that is to say, provided the curve be neither a parabola nor an ellipse, but if it be either of these, there can exist no asymptote; therefore the hyperbola is the only line of the second order which has asymptotes.

(85.) When the curve is referred to polar coordinates, then, since the radius vector of the point of contact is infinite when the tangent becomes an asymptote, it follows that if for $r = \infty$ the subtangent is finite; this subtangent may be determined by (80) in terms of ω, and ω may be found from the equation of the curve, so that there will thus be determined a point in the asymptote and its direction, which is all that is necessary to fix its position. There will always be an asymptote if ω is finite, for $r = \infty$. If, for $r = \infty$, ω is also ∞, there exists no asymptote.

3. Let the curve be the hyperbolic spiral.

By ex. 10, art. 81, the subtangent at any point is constant, and equal to a, therefore there must be an asymptote; also by the equation of the curve $\omega = \frac{a}{r} = 0$, when $r = \infty$, therefore the asymptote is perpendicular to the fixed axis, at the distance a from the pole.



Neither the logarithmic spiral, nor the spiral of Archimedes have an asymptote.

4. Let the spiral whose equation is

$$r = \frac{a\omega^2}{\omega^2 - 1} = \frac{a}{1 - \omega^{-2}}$$

be proposed, which admits of a rectilinear asymptote, because $\omega = 1$ renders $r = \infty$. The *direction*, therefore, of the asymptote is ascertained, and consequently the direction of the infinite radius vector, since they must be parallel. It remains, therefore, to determine the subtangent, or distance of the asymptote from the pole

$$\frac{dr}{d\omega} = -\frac{2a\omega^{-3}}{(1 - \omega^{-2})^2} = -\frac{2r^2}{a\omega^3},$$

$$\therefore r^2 \div \frac{dr}{d\omega} = r^2 \div \frac{2r^2}{a\omega^3} = \frac{a\omega^3}{2} = \frac{a}{2} = T,$$

because $\omega = 1$ when $r = \infty$.

(86.) Although we do not propose to treat fully in this place of *curvilinear* asymptotes, yet we may remark in passing, that if r should be finite although ω be infinite, it will prove that the spiral must be continued for an infinite number of revolutions round its pole, before it can meet the circumference of a circle whose radius is this finite value. In such a case, therefore, the spiral has a *circular asymptote*. If, moreover, the value of r for $\omega = \infty$ be greater than the value of r for every other value of ω, the spiral will be included within its circular asymptote, but, otherwise, it will be without this circle.

5. Thus in the spiral whose equation is

$$(r^2 - ar)\,\omega^2 - 1 = 0, \text{ or } \omega = \frac{1}{\sqrt{r^2 - ar}},$$

ω is infinite when $r = a$, and for all less values of r, ω is imaginary; hence the spiral can never approach so near to the pole as $r = a$, till after an infinite number of revolutions; so that the circumference whose radius is a is *within* the spiral, and is asymptotic.

If, on the contrary, the equation had been

$$\omega = \frac{1}{\sqrt{ar - r^2}},$$

then also $r = a$ gives $\omega = \infty$, but for all other real values of ω, r is less than a, so that this spiral is enclosed by its asymptote circle, the radius of which is a.

6. To determine the rectilinear asymptote to the logarithmic curve.

The axis of x.

7. To determine the equation of the asymptote to the curve whose equation is

$$y^3 = ax^2 + x^3.$$

The equation is $y = x + \frac{1}{3}a$.

8. To determine the rectilinear asymptote to the spiral whose equation is

$$r = a\omega^{-\frac{1}{2}}.$$

The fixed axis is the asymptote.

9. To determine whether the spiral shown to have a rectilinear asymptote in ex. 4 has also a circular asymptote.

The circle whose centre is the pole and radius $= a$ is an asymptote.

(87.) Before terminating the present chapter, it will be necessary to exhibit the expression for the differential of the arc of any plane curve, as we shall have occasion to employ this expression in the next chapter.

Let us call the arc AB of any plane curve s, and the coordinates of B, x, y; let also BD be a tangent at B, and BC any chord, then if BE, ED are parallel to the rectangular axes, BC will be increment of the arc s, corresponding to $BE = h$, the increment of the abscissa x.



Now, putting tan $DBE = a$, we have

$$ED = ha \therefore BD = \sqrt{h^2 + h^2a^2}$$

and

$$\frac{BD + DC}{BC} = \frac{\sqrt{h^2(1 + a^2)} + ha - CE}{\sqrt{h^2 + CE^2}}.$$

This ratio continually approaches to $\frac{CE}{CE}$, or to unity, as h diminishes, and this it actually becomes when $h = 0$. Consequently, since the arc BC is always, when of any definite length, longer than the chord BC, and shorter than BD + DC,* it follows that, when $h = 0$, the ratio of the arc to either of these must be unity; therefore

$$\text{in the limit } \frac{\text{arc BC}}{\text{chord BC}} = 1 \therefore \frac{\text{arc BC}}{h} \div \frac{\text{chord BC}}{h} = 1,$$

but

$$\frac{\text{chord BC}}{h} = \frac{\sqrt{h^2 + CE^2}}{h} = \sqrt{1 + \frac{CE^2}{h^2}},$$

and CE is the increment of the ordinate y, corresponding to the increment h of the abscissa x; hence, when $h = 0$, the ratio becomes

---

* See Young's *Elements of Plane Geometry*, p. 204.

$$\frac{ds}{dx} \div \sqrt{1+\frac{dy^2}{dx^2}} = 1,$$

$$\therefore \frac{ds}{dx} = \sqrt{1+\frac{dy^2}{dx^2}}.$$

If any other independent variable be taken instead of x, then, denoting the several differential coefficients relatively to this new variable by (dx), (dy), (ds), we have (66)

$$(ds) = \sqrt{(dx)^2 + (dy)^2}.$$

At the point where $\frac{dy}{dx} = 0$, $\frac{ds}{dx} = 1$, or $(ds) = (dx)$.

From the expression for the length of the normal at page 145, it follows that

$$\frac{ds}{dx} = \frac{N}{y}.$$

CHAPTER II.

ON OSCULATION, AND THE RADIUS OF CURVATURE OF PLANE CURVES.

(88.) Let

$$y = f(x),\ Y = F(x),$$

be the equations of two plane curves, in the former of which we shall suppose the constants a, b, c, &c. to be known, and therefore the curve itself to be determinate; while in the latter we shall consider the constants A, B, C, &c. to be unknown, or arbitrary, and therefore the *species* only of the curve given. The constants which enter into the equation of a curve are usually called the *parameters*.

If, now, x take the increment h, and the corresponding ordinates y', Y' be developed, we shall have, by Taylor's theorem,

$$y' = y + \frac{dy}{dx}h + \frac{d^2y}{dx^2}\frac{h^2}{1.2} + \frac{d^3y}{dx^3}\frac{h^3}{1.2.3} + \&c. \ldots . (1),$$

$$Y' = Y + \frac{dY}{dx}h + \frac{d^2Y}{dx^2}\frac{h^2}{1.2} + \frac{d^3Y}{dx^3}\frac{h^3}{1.2.3} + \&c. \ldots . (2).$$

The parameters which enter (2) being arbitrary, they may be determined so as to fulfil as many of the conditions

$$y = Y, \frac{dy}{dx} = \frac{dY}{dx}, \frac{d^2y}{dx^2} = \frac{d^2Y}{dx^2}, \&c. \ldots . (3),$$

as there are parameters, but obviously not more conditions.

We shall thus have the values of A, B, C, &c. in terms of x, and of the fixed parameters a, b, c, &c.; which values, substituted in (2), will cause so many of the leading terms in both series to become *identical*, whatever be the value of x. Other corresponding terms of the two series may, indeed, be rendered also identical, but this can take place only accidentally, not necessarily. Hence, whatever particular value we now give to x, the resulting values of the corresponding coefficients will necessarily agree to the extent mentioned, that is, as far as the n first terms, if there are n constants originally in (2); and this is true, even if such particular value of x render any of the coefficients infinite, inasmuch as they are *always identical* as far as these terms, but no further.

We know, however, that in those cases where any of the coefficients become infinite, (1) and (2) will fail to represent the true developments of the ordinates y', Y' at the proposed points. Nevertheless, as the two series have been rendered identical, as far as n terms, should they both fail within this extent, the terms which supply these in the true development, must necessarily be identical. (*See note C at the end.*)

Now the greater number of leading terms in the two developments, which are identical, the nearer will the developments themselves approach to identity, provided, at least, h may be taken as small as we please; for if $n - 1$ terms in each are identical, we may represent the difference of the two developments by

$$A_n h^\alpha + S - (A'_n h^{\alpha'} + S'),$$

$$\text{or } A_n h^\alpha - A'_n h^{\alpha'} + S - S' \ldots . (4),$$

where S, S′ represent the sums of the remaining terms in each series after the nth. Hence, h^α being the lowest power of h which enters this expression, for the difference it follows from (47), that a value may be given to h small enough to cause the term $A_n h^\alpha$ to become greater than all the other terms in (4), and consequently, for this small value,

$$A_n h^\alpha - A'_n h^{\alpha'} > S - S',$$

and, therefore, the whole difference (4) is greater than twice S — S′, but when the nth term is the same in both developments, as well as the preceding terms, then the difference (4) is reduced simply to S — S′, which we have just seen to be less than (4). Consequently the developments approach nearer to identity, for all values of h between some certain finite value h' and 0, as the number of identical leading terms become greater.

When the first of the conditions (3) exist, the curves have a common point; when the second also exists they have a common tangent at that point, and are consequently in contact there, and the contact will be the more intimate, or the curves will be the closer in the vicinity of the point, as the number of following conditions become greater; so that of all curves of a given species, *that* will touch any fixed curve at a proposed point with the closest contact whose parameters are *all* determined agreeably to the conditions (3). No other curve of the same species can, from what is proved above, approach so nearly to coincidence with the proposed, in the immediate vicinity of the point of contact, as this; so that no other of that species can pass between this and the proposed. A curve, thus determined, is said to be, in reference to the proposed curve, its *osculating curve* of the given species.

(89.) It appears, from what has now been said, that there may be different *orders* of contact at any proposed point. The first two of the conditions (3) must exist for there to be contact at all; therefore, when only these exist, the contact is called *simple contact*, or *contact of the*

first order; if the next condition also exist, the contact is of the *second order*, and so on; and it is obvious, that of any given species, *the osculating curve* will have the highest order of contact, at any proposed point, in a given curve. If the curve, given in species, has n parameters, the highest order of contact will be the $n-1$th, unless, indeed, the same values of these parameters that fulfil the n conditions (3), should happen also to fulfil the $n+1$th, the $n+2$th, &c.; but this, as observed before, can take place only accidentally, and cannot be predicted of any proposed point, although we see it is possible for such points to exist.

(90.) At those points in the proposed curve, for which Taylor's development does not fail, contact of an even order is both contact and intersection, and contact of an odd order is without intersection; before proving this, however, we may hint to the student that contact is not opposed to intersection, for two curves are said to be in contact at a point, when they have a common tangent at that point; and yet, as we are about to show, one of these curves may pass between the tangent and the other, and so intersect where they are admitted to be in contact. To prove the proposition, let us take the difference (4), which, when Taylor's theorem holds, is

$$(A_n - A'_n)\,h^\alpha + S - S' \ldots . \quad (5),$$

A_n A'_n being here the $n-1$th differential coefficients. If these are *odd*, the contact is of an even order; also α being odd, h^α will have contrary signs for $h = +h'$ and $h = -h'$, and, therefore, since for these small values of h, the sign of the whole expression (5) is the same as that of the first term, the difference of the ordinates corresponding to $x+h$, and to $x-h$, will be the one positive and the other negative, so that the two curves must necessarily cross at the point whose abscissa is x.



But if α is even, the contact is of an odd order, and the difference (5) between the ordinates of the two curves corresponding to the same abscissa, $x+h$, will, for a small value of h, have the same sign, whether h be positive or negative; so that, in this case, the curves do not cross each other at the point of contact.

(91.) The student must not fail to bear in remembrance, that the proposition just established comprehends only those points of the proposed curve, at which none of the differential coefficients become infinite, from the first to that immediately beyond the coefficient which fixes the order of the contact. For it is only upon the supposition that the true development, within these limits, proceeds according to the ascending integral and positive powers of h, that the foregoing conclusions respecting the signs of the difference (5) can be fairly drawn. *(See note C.)*

(92.) From the principles of osculation now established, it is evident that any plane curve being given, and any point in it chosen, we may always find what particular curve, of any proposed species, shall touch at that point with the closest contact, or which shall most nearly coincide with the given curve in the immediate vicinity of the proposed point. Thus, an *ellipse* or a *parabola* being given, and a point in it proposed, we may determine the *circle* that shall approach more nearly to coincidence with that ellipse or parabola in the vicinity of the proposed point, than any other circle, and which will therefore better represent the curvature of the given curve at the proposed point than any other. On account of its simplicity and uniformity, the circle is the curve employed to estimate, in this way, the curvature of other curves at proposed points; that is, the curvature is estimated by the curvature of the *osculating circle*, or rather as the curvature of a circle increases as the radius diminishes, and vice versâ, it is usual to adopt, as a fit representation of the curvature, the reciprocal of the radius.

The osculating circle is called the *circle of curvature*, and its radius the *radius of curvature;* and, from what has been said above, it follows that the determination of the curvature at any point in a proposed curve, reduces itself to the determination of this radius: to this, therefore, we shall now proceed.

Radius of Curvature.

PROBLEM I.

(93.) To determine the radius of curvature at any proposed point of a given curve.

The general equation of a circle being

$$(x-\alpha)^2+(y-\beta)^2=r^2,$$

it becomes determined as soon as we fix the values of the parameters α, β, r, and these may be determined, so as to fulfil any three independent conditions, but not more. In the case before us, the conditions to be fulfilled are those of (3), art. (87): that is to say, putting p', p'', &c. for the successive differential coefficients derived from $Y=F(x)$, the equation of the given curve, the conditions to be fulfilled are

$$y=Y,\quad \frac{dy}{dx}=p',\quad \frac{d^2y}{dx^2}=p'',$$

in order that the resulting values of α, β, r, may belong to the equation of the osculating circle. Now

$$\frac{dy}{dx}=-\frac{x-\alpha}{y-\beta},\quad \frac{d^2y}{dx^2}=-\frac{1}{y-\beta}-\frac{(x-\alpha)^2}{(y-\beta)^3}=-\frac{r^2}{(y-\beta)^3};$$

hence, putting y for its equal Y, the three equations for determining α, β, and r, are

$$(x-\alpha)^2+(y-\beta)^2=r^2 \quad \ldots\ (1),$$

$$(x-\alpha)+p'(y-\beta)=0 \quad \ldots\ (2),$$

$$p''(y-\beta)^3=-r^2 \quad \ldots\ (3).$$

From the second equation

$$(x-\alpha)^2=p'^2(y-\beta)^2.$$

Substituting this in the first,

$$(p'^2+1)\,(y-\beta)^2=r^2.$$

Adding this last to the third, there results

$$y-\beta=-\frac{p'^2+1}{p''},$$

which, substituted in (2), gives

$$x-\alpha=\frac{p'(p'^2+1)}{p''}\quad\therefore\quad r^2=\frac{(p'^2+1)^3}{(-p'')^2}.$$

Consequently, $\qquad \alpha=x-\frac{p'(p'^2+1)}{p''},\quad \beta=y+\frac{p'^2+1}{p''},$

$$r=\frac{(p'^2+1)^{\frac{3}{2}}}{-p''}=-\frac{\frac{ds^3}{dx^3}}{p''}=-\frac{N^3}{y^3p''}.$$ (See p. 159).

These equations completely determine the osculating circle, whenever the coordinates x, y of the proposed point are given.

Should this point be such as to render $p'=0$, then the expression, for the radius of curvature at that point, becomes $r=-\frac{1}{p''}$, and the tangent at the proposed point must be parallel to the axis of x (78).

The sign of p'' will always make known the direction of curvature; for, if a linear tangent be drawn at the proposed point (x, y), we shall have for the ordinate of the tangent, due to the abscissa $x+h$, the expression $y+p'h$; and for that of the curve, due to the same abscissa, $y+p'h+\frac{1}{2}p''h^2+$ &c.; and it is obvious that if p'' is *plus*, this latter ordinate, for a small value of h, must exceed the other; and thus the tangent will proceed from the point in a direction between the curve and axis of x, that is, the curve will, at that point, be *convex* towards the axis of x. If, on the contrary, p'' is *minus*, then the ordinate of the curve will be less than that of the tangent; so that the curve proceeds from the point between the tangent and axis of x, and is thus *concave* towards that axis.

Should $p''=0$ at the proposed point, r will be infinite, whether $p'=0$ or not, so that the osculating circle then becomes a straight line; as, therefore, this straight line has contact of the second order, the parts of the curve in the vicinity of the point will lie on contrary sides of it, as in



the annexed diagram (90); that is, supposing p''' is neither 0 nor ∞. If $p''' = 0$, and the next following differential coefficient neither 0 nor ∞, the contact will be of an order which is unaccompanied by intersection. The curve will be *convex* at the point if the first finite differential coefficient be of an even order and $+$; and it will be *concave* when the sign of this coefficient is $-$.

A point at which the tangent intersects the curve, or at which the curve changes from convex to concave, is called *a point of inflexion*, or, *a point of contrary flexure*. The analytical indications of such points will be more fully enquired into when we come to speak of the *singular points* of curves.

(94.) By referring to equation (2) above, which has place even when the contact is but of the first order, we learn that the centre (α, β) of every touching circle, is always on the normal at the point of contact; for that equation is the same as

$$(\beta - y) = -\frac{1}{\frac{dy}{dx}}(\alpha - x).$$

We shall now apply the general expression, for the radius of curvature, to a few particular cases.

EXAMPLES.

(95.) 1. Required the radius of curvature, at any point in a parabola.

Differentiating the equation of the curve, $y^2 = 4mx$, we have

$$2yp' = 4m \quad \therefore \; p' = \frac{2m}{y},$$

$$2yp'' + 2p'^2 = 0 \quad \therefore \; p'' = -\frac{p'^2}{y} = -\frac{4m^2}{y^3},$$

$$\therefore \; r = -\frac{N^3}{y^3 p''} = \frac{N^3}{4m^2}.$$

As the expression for the normal diminishes with x, the vertex is the point of greatest curvature, r being there equal to $2m$, half the parameter; moreover, p'' being negative, the curve is always *concave* to the axis of x.

2. To determine the radius of curvature at any point in an ellipse.

By differentiating the equation

$$a^2y^2 + b^2x^2 = a^2b^2,$$

we have

$$a^2yp' + b^2x = 0 \;\therefore\; p' = -\frac{b^2x}{a^2y},$$

$$a^2yp'' + a^2p'^2 + b^2 = 0 \;\therefore\; p'' = -\frac{b^2 + a^2p'^2}{a^2y} = -\frac{b^4}{a^2y^3},$$

$$\therefore\; r = -\frac{(p'^2+1)^{\frac{3}{2}}}{p''} = \frac{(a^4y^2 + b^4x^2)^{\frac{3}{2}}}{a^6y^3} \cdot \frac{a^2y^3}{b^4} = \frac{(a^4y^2 + b^4x^2)^{\frac{3}{2}}}{a^4b^4} \quad \ldots\ (1).$$

or, using the expression for r in terms of the normal, we have

$$r = -\frac{N^3}{y^3p''} = \frac{a^2}{b^4}N^3 \quad \ldots\ (2).^*$$

Since *(Anal. Geom.*, p. 87),

$$aN = bb' \;\therefore\; r = \frac{b'^3}{ab} \quad \ldots\ (3).$$

At the vertex of the transverse axis $r = \frac{b^2}{a}$ = semiparameter = the normal, *(Anal. Geom.*, p. 69).

At the vertex of the conjugate $r = \frac{a^2}{b}$.

From equations (2) and (3) it follows that, in the ellipse, the radius of curvature varies as the cube of the normal, or as the cube of the diameter parallel to the tangent through the proposed point.

It is often desirable to obtain r as a function of λ, the angle included between the normal and the transverse axis. For this purpose we have, since

---

* This expression will become identical with (1) if we substitute for N^3 its value given in the Analytical Geometry, p. 84, viz.

$$N^3 = \left\{\frac{a^4y^2 + b^4x^2}{a^4}\right\}^{\frac{3}{2}}.$$

$x^2 = a^2(1 - \frac{y^2}{b^2})$, $y^2 = N^2 \sin^2\lambda$, and (*Anal. Geom.* p. 84)

$$N^2 = \frac{b^4}{a^4} x^2 + y^2,$$

$$N^2 = \frac{b^4}{a^2}(1 - \frac{N^2 \sin^2\lambda}{b^2}) + N^2 \sin^2\lambda,$$

$$\therefore N^2 \{1 - (1 - \frac{b^2}{a^2}) \sin^2\lambda\} = \frac{b^4}{a^2};$$

but, (*Anal. Geom.*, p. 68,)

$$1 - \frac{b^2}{a^2} = e^2,$$

$$\therefore N = \frac{b^2}{a} \cdot \frac{1}{(1 - e^2 \sin^2\lambda)^{\frac{1}{2}}},$$

$$\therefore r = \frac{a^2}{b^4} N^3 = \frac{b^2}{a(1 - e^2 \sin^2\lambda)^{\frac{3}{2}}} = \frac{a(1 - e^2)}{(1 - e^2 \sin^2\lambda)^{\frac{3}{2}}}.$$

The curvature $\frac{1}{r}$ is greatest at the vertex of the transverse, and least at the vertex of the conjugate axis. At the former point $r = \frac{b^2}{a}$, and at the latter $r = \frac{a^2}{b}$, as shown in the preceding page.

(96.) The present is a very important problem, being intimately connected with enquiries relative to the figure of the earth.

By means of the last expression for r, the ratio of the polar and equatorial diameters of the earth, may be readily deduced, when we know the lengths of a degree of the meridian in two known latitudes, L, l, for these lengths may, without error, be considered to coincide with the osculating circles through their middle points; and, since similar arcs of circles are as their radii, we have, by putting M, m for the measured degrees, and R, r for the corresponding radii,

$$R : r :: M : m,$$

but

$$R = \frac{a(1-e^2)}{(1-e^2\sin^2 L)^{\frac{3}{2}}}, \text{ and } r = \frac{a(1-e^2)}{(1-e^2\sin^2 l)^{\frac{3}{2}}},$$

therefore, since $mR = Mr$, we have

$$\frac{m}{(1-e^2\sin^2 L)^{\frac{3}{2}}} = \frac{M}{(1-e^2\sin^2 l)^{\frac{3}{2}}},$$

or

$$m^{\frac{2}{3}}(1-e^2\sin^2 l) = M^{\frac{2}{3}}(1-e^2\sin^2 L),$$

$$\therefore e^2 = 1 - \frac{b^2}{a^2} = \frac{M^{\frac{2}{3}} - m^{\frac{2}{3}}}{M^{\frac{2}{3}}\sin^2 L - m^{\frac{2}{3}}\sin^2 l} *,$$

$$\therefore \frac{a}{b} = \sqrt{\left\{\frac{M^{\frac{2}{3}}\sin^2 L - m^{\frac{2}{3}}\sin^2 l}{m^{\frac{2}{3}}\cos^2 l - M^{\frac{2}{3}}\cos^2 L}\right\}} = \sqrt{\left\{\frac{\sin^2 L - (\frac{m}{M})^{\frac{2}{3}}\sin^2 l}{(\frac{m}{M})^{\frac{2}{3}}\cos^2 l - \cos^2 L}\right\}}.$$

If $l = 0$; that is, if the degree m is measured at the equator, then

$$\frac{a}{b} = \sqrt{\left\{\frac{\sin^2 L}{(\frac{m}{M})^{\frac{2}{3}} - \cos^2 L}\right\}},$$

$$= \frac{\sin L}{\sqrt{\left\{(\frac{m}{M})^{\frac{2}{3}} - \cos^2 L\right\}}}.$$

---

* From this expression for e^2, in terms of the measured quantities M, m, L, l, the actual lengths of the equatorial and polar radii may be found, thus:

$$\text{Since } r \times 3{\cdot}1416 = 180m \quad \therefore r = \frac{180m}{3{\cdot}1416}.$$

$$\therefore \frac{180m}{3{\cdot}1416} = \frac{a(1-e^2)}{(1-e^2\sin^2 l)^{\frac{3}{2}}},$$

$$\therefore a = \frac{180m(1-e^2\sin^2 l)^{\frac{3}{2}}}{3{\cdot}1416(1-e^2)},$$

whence may be deduced b.

3. To determine the radius of curvature at any point in the logarithmic curve, its equation being $y = a^x$.

$$r = \frac{(m^2 + y^2)^{\frac{3}{2}}}{my}, \text{ } m \text{ being the modulus.}$$

4. To determine the radius of curvature at any point in the cubical parabola, its equation being $y^3 = ax$.

$$r = \frac{(9y^4 + a^2)^{\frac{3}{2}}}{6a^2y}.$$

PROBLEM II.

(97.) To determine those points in a given curve, at which the osculating circle shall have contact of the third order.

It is here required to find for what points of a given curve the values of α, β, r, determined by the first three conditions (3), art. (87), satisfy also the fourth condition.

The differential coefficient p''', as derived from equation (3), p. 129, is

$$p''' = -\frac{3p''p'}{y - \beta},$$

and this must agree with the p''' derived from the equation of the proposed curve, at those points where the contact is of the third order; that is, the abscissas of these points will all be given by the roots of the equation

$$(y - \beta)\, p''' + 3p''\, p' = 0,$$

and it may be easily shown, that the points which satisfy this equation are those of greatest and least curvature; for, since

$$r = -\frac{(p'^2 + 1)^{\frac{3}{2}}}{p''},$$

$$\therefore \frac{dr}{dx} = \frac{-3(p'^2 + 1)^{\frac{1}{2}} p' p''^2 + (p'^2 + 1)^{\frac{3}{2}} p'''}{p''^2},$$

and when r is a maximum or a minimum, this expression is equal

to 0 (49); hence

$$-3p'p''^2+(p'^2+1)p'''=0,$$

or, dividing by p'' and recalling the value of $y-\beta$, deduced in (93), we have, finally,

$$(y-\beta)p'''+3p''p'=0;$$

which being the same equation as that deduced above, it follows that at the points of maximum and minimum curvature, the contact is of the third order.

SCHOLIUM.

(98.) In the preceding investigations we have always considered x to be the independent variable, because the expression for the radius of curvature has been obtained comformably to this hypothesis. But if any other quantity is taken for the independent variable, the foregoing expression for r will not apply; therefore, in order to give the greatest generality possible to the formula for the radius of curvature, we shall now suppose any arbitrary quantity whatever to be the independent variable, x and y being functions of it. Hence, instead of p' and p'', we shall have (66)

$$\frac{(dy)}{(dx)} \text{ and } \frac{(d^2y)(dx)-(d^2x)(dy)}{(dx^2)},$$

the parentheses serving to intimate that the independent variable, according to which the differentials of the functions x, y are taken, is arbitrary; and the differential of which, when chosen, is, with its proper powers to be introduced as denominators of the above differentials. Making therefore, these substitutions in the expression for r, it becomes

$$r=-\frac{\{(dy)^2+(dx)^2\}^{\frac{3}{2}}}{(d^2y)(dx)-(d^2x)(dy)};$$

or, since (86)

$$(ds)=\sqrt{(dy)^2+(dx)^2},$$

whatever be the independent variable,

$$r = -\frac{(ds)^3}{(d^2y)(dx) - (d^2x)(dy)} \quad \ldots\ldots (1).$$

(99.) This expression is of the utmost generality, and will furnish a correct formula for every hypothesis respecting the independent variable. Thus, if x be chosen for the independent variable, then $(dx) = 1$ and $(d^2x) = 0$, and the formula in that case is

$$r = -\frac{\frac{ds^3}{dx^3}}{\frac{d^2y}{dx^2}} \quad \ldots\ldots (2),$$

being the same as that at first given as it ought to be. If y be the independent variable, then $(dy) = 1$ and $(d^2y) = 0$; so that, upon this hypothesis, the formula is

$$r = \frac{\frac{ds^3}{dy^3}}{\frac{d^2x}{dy^2}} \quad \ldots\ldots (3).$$

If s be the independent variable, then $(ds) = 1$; that is,

$$\sqrt{(dx)^2 + (dy)^2} = 1,$$

and the differential of this being $= 0$, we have

$$\therefore (d^2y) = -\frac{(dx)}{(dy)}(d^2x) \quad \ldots\ldots (4);$$

substituting this in the denominator of (1), there results

$$r = \frac{(dy)}{(d^2x)} \div \sqrt{(dx)^2 + (dy)^2} = \frac{(dy)}{(d^2x)} = \frac{\frac{dy}{ds}}{\frac{d^2x}{ds^2}} \quad \ldots\ldots (5).$$

By squaring (1) on this last hypothesis, we have

$$r^2 = \frac{1}{(d^2y)(dx) - (d^2x)(dy)^2};$$

but, since from (4)

$$(d^2y)(dy) + (d^2x)(dx) = 0,$$

it may be added to the denominator of this expression for r^2 without affecting its value; so that

$$r^2 = \frac{1}{\{(d^2y)(dx) - (d^2x)(dy)\}^2 + \{(d^2y)(dy) + (d^2x)(dx)\}^2}$$

$$= \frac{1}{(dy)^2 + (dx)^2 \times (d^2y)^2 + (d^2x)^2}$$

$$= \frac{1}{(d^2y)^2 + (d^2x)^2} \quad \therefore r = \frac{1}{\sqrt{(\frac{d^2y}{ds^2})^2 + (\frac{d^2x}{ds^2})^2}} \quad \ldots\ldots (6).$$

(100.) We shall now proceed to determine a suitable formula for *polar curves*.

If the circle whose equation is (1) p. 164, be transformed from rectangular to polar coordinates, the pole being at the origin of the primitive axes, and the axis of x being the fixed line from which the variable angle ω of the radius vector γ is measured, we shall have (*Anal. Geom.* p. 131),

$$(\gamma \cos \omega - \alpha)^2 + (\gamma \sin \omega - \beta^2) = r^2 \ldots\ldots (1),$$

If, therefore, we differentiate on the supposition that ω is the independent and γ the dependent variable, and denote the first and second differential coefficients by $p_,$ and $p_{,,}$, we shall have

$$(\gamma \cos \omega - \alpha)(p_, \cos \omega - \gamma \sin \omega) + (\gamma \sin \omega - \beta)(p_, \sin \omega + \gamma \cos \omega) = 0 \ldots (2)$$

$$(p_, \cos \omega - \gamma \sin \omega)^2 + (\gamma \cos \omega - \alpha)(p_{,,} \cos \omega - 2p_, \sin \omega - \gamma \cos \omega) +$$

$$(p_, \sin \omega + \gamma \cos \omega)^2 + (\gamma \sin \omega - \beta)(p_{,,} \sin \omega + 2p_, \cos \omega - \gamma \sin \omega) = 0 \ldots (3)$$

If from the two latter equations we determine the values of $\gamma \sin \omega - \beta$ and $\gamma \cos \omega - \alpha$, and substitute them in (1), we shall obtain the following expression for r in functions of γ and its differential coefficients, viz.

$$r = \frac{(\gamma^2 + p_,^2)^{\frac{3}{2}}}{\gamma^2 + 2p_,^2 - \gamma p_{,,}} \quad \ldots . \quad (4);$$

but we shall arrive at this expression more readily by first deducing from the equations

$$y = \gamma \sin \omega, \; x = \gamma \cos \omega$$

the differential coefficients

$$\frac{dy}{d\omega} = \gamma \cos \omega + p_, \sin \omega = (dy)$$

$$\frac{dx}{d\omega} = -\gamma \sin \omega + p_, \cos \omega = (dx)$$

$$\frac{d^2y}{d\omega^2} = -\gamma \sin \omega + 2p_, \cos \omega + p_{,,} \sin \omega = (d^2y)$$

$$\frac{d^2x}{d\omega^2} = -\gamma \cos \omega - 2p_, \sin \omega + p_{,,} \cos \omega = (d^2x),$$

and then substituting them in the general formula (1), p. 172.

Since (80) the expression for the normal PN is

$$N = (\gamma^2 + p_,^2)^{\frac{1}{2}},$$

we may put the above expression for r under the form

$$r = \frac{N^3}{r^2 + 2p_,^2 - \gamma p_{,,}} \quad \ldots . \quad (5),$$

5. To determine the radius of curvature at any point in the logarithmic spiral

$$y = a^\omega$$

$$\frac{d\gamma}{d\omega} = \frac{a^\omega}{m} = \frac{\gamma}{m} = p_,$$

$$\frac{d^2\gamma}{d\omega} = \frac{\gamma}{m^2} = p_{,,}.$$

Hence

$$r = \frac{(\gamma^2 + p_i^2)^{\frac{3}{2}}}{\gamma^2 + 2p_i^2 - \gamma p_{ii}} = (\gamma^2 + p_i^2)^{\frac{1}{2}} = \gamma\sqrt{1 + \frac{1}{m^2}} =$$

$$\gamma\sqrt{1 + \frac{1}{\tan^2 \mathrm{P}}} \ (art.\ 80) = \gamma \operatorname{cosec} \mathrm{P}.$$

It appears, therefore, that the radius of curvature is always equal to the normal.

6. To determine the radius of curvature at any point in the curve whose equation is

$$\gamma = 2\cos\omega \pm 1$$

$$\therefore r = \frac{(5 \pm 4\cos\omega)^{\frac{3}{2}}}{9 \pm 6\cos\omega}.$$

(101.) The general expression for the radius of curvature at any point (γ, ω) of a curve referred to polar coordinates, may be concisely exhibited in terms of γ and p, the radius vector and perpendicular from the pole upon the tangent.

For differentiating the expression for p at (82), and substituting p_i for $\frac{\mathrm{d}r}{\mathrm{d}\omega}$, and γ for r, we have

$$\frac{\mathrm{d}p}{\mathrm{d}\omega} = \frac{2\gamma p_i p_i^2 + \gamma^3 p_i - \gamma^2 p_i p_{ii}}{(\gamma^2 + p_i^2)^{\frac{3}{2}}},$$

which is obviously equal to the reciprocal of the expression for r in equation (4), multiplied by γp_i. Consequently,

$$r = \frac{\gamma p_i}{\frac{\mathrm{d}p}{\mathrm{d}\omega}} = \frac{\gamma \frac{\mathrm{d}\gamma}{\mathrm{d}\omega}}{\frac{\mathrm{d}p}{\mathrm{d}\omega}};$$

that is, (p. 127,)

$$r = \gamma \frac{\mathrm{d}\gamma}{\mathrm{d}p}.$$

If from the centre of curvature a perpendicular were drawn to the radius vector, it is plain that the angle included between this perpen-

dicular and the radius of curvature would be equal to P, the angle included between the radius vector and tangent.* The perpendicular here supposed to be drawn, would obviously bisect that chord of the osculating circle which passes through the pole, and which, for distinction, is called the *chord of curvature.*

Hence, from the similar right-angled triangles, we should have the following equality, viz.

$$\tfrac{1}{2} \text{ chord of curvature} = r\frac{p}{\gamma} = p\frac{\mathrm{d}\gamma}{\mathrm{d}p},$$

an expression which is of use in Astronomy.†

CHAPTER III.

ON INVOLUTES, EVOLUTES, AND CONSECUTIVE CURVES.

(102.) If osculating circles be applied to *every* point in a curve, the locus of their centres is called the *evolute* of the proposed curve, this latter being called the *involute.*

The equation of the evolute may be determined by combining the equation of the proposed curve with the equations (2), (3), p. 129, containing the *variable* coordinates α, β of the centre. As these three equations must exist simultaneously for every point of contact (x, y), the two quantities x, y may be eliminated, and, therefore, a resulting equation obtained, containing only α and β, which equation, therefore will express the general relation between α and β for every point (x, y); in other words, it will represent the locus of the centres of the osculating circles.

* Although neither this perpendicular, nor that from the pole to the tangent, called p, are actually exhibited in the figure at page 148, yet the student, bearing in mind that the centre of curvature is always on the normal, will find no difficulty in supplying these two lines.

† See Young's "Treatise on Mechanics," p. 212.

Or, representing the equation of the proposed curve by $y = \mathrm{F}(x)$, we shall have to eliminate x and y from the equations (p. 130)

$$\alpha = x - \frac{p'(p'^2+1)}{p''},\ \beta = y + \frac{p'^2+1}{p''}$$

$$y = \mathrm{F}(x),$$

when the resulting equation in α, β will be that of the evolute.

EXAMPLES.

(103.) To determine the evolute of the common parabola

$$y^2 = 4mx \therefore p' = \frac{2m}{y} \therefore p'' = -\frac{4m^2}{y^3},$$

$$\therefore 1 + p'^2 = \frac{y^2+4m^2}{y^2} = 1 + \frac{m}{x},\ \frac{p'}{p''} = -\frac{y^2}{2m},$$

$$\therefore \alpha = x + \frac{y^2}{2m} + 2m = 3x + 2m \therefore x = \frac{\alpha - 2m}{3},$$

$$\beta = y - \frac{y^3}{4m^2} \therefore y = \frac{-y^3}{m\frac{y^2}{x}} = \frac{-2x^{\frac{3}{2}}}{m^{\frac{1}{2}}} \therefore x = \left(\frac{m\beta^2}{4}\right)^{\frac{1}{3}},$$

$$\therefore \beta^2 = \frac{4}{27m}(\alpha - 2m)^3,$$

which is the equation of the evolute. If the origin be removed to that point in the axis of x whose abscissa is $2m$, then the equation becomes

$$\beta^2 = \frac{4}{27m}\alpha^3,$$

the locus of which is called the *semicubical parabola*. It passes through the origin because $\beta = 0$ when $\alpha = 0$; therefore the focus of the proposed involute is in the middle, between its vertex and the vertex of the evolute. *(Anal. Geom.* art. 100.) The curve con-



sists of two branches symmetrically situated with respect to the axis of x, or of α, and lies entirely to the right of the origin, for every positive value of α gives two equal and opposite values of β, and for negative values of α and β is impossible. It is easy to see, therefore, that the form of the curve is that represented in the margin.

2. To determine the evolute of the ellipse.

By example 2, page 167, we have

$$p' = -\frac{b^2x}{a^2y},\ p'' = -\frac{b^4}{a^2y^3},$$

$$\therefore 1 + p'^2 = \frac{a^4y^2 + b^4x^2}{a^4y^2},\ \frac{p'}{p''} = \frac{xy^2}{b^2},$$

$$\therefore \alpha = x - \frac{x(a^4y^2 + b^4x^2)}{a^4b^2},\ \beta = y - \frac{y(a^4y^2 + b^4x^2)}{a^2b^4}.$$

Now, since, by the equation of the curve,

$$a^2y^2 = a^2b^2 - b^2x^2 \text{ or } b^2x^2 = a^2b^2 - a^2y^2,$$

$$\therefore a^4y^2 + b^4x^2 = b^2(a^4 - c^2x^2) \text{ or } = a^2(b^4 + c^2y^2),$$

c^2 being put for $a^2 - b^2$. Hence, by substitution,

$$\alpha = \frac{c^2x^3}{a^4},\ \beta = -\frac{c^2y^3}{b^4},$$

$$\therefore x^2 = (\frac{\alpha^2a^8}{c^4})^{\frac{1}{3}},\ y^2 = (\frac{\beta^2b^8}{c^4})^{\frac{1}{3}}.$$

Substituting these values in the equation of the involute, we have

$$a^2(\frac{\beta^2b^8}{c^4})^{\frac{1}{3}} + b^2(\frac{\alpha^2a^8}{c^4})^{\frac{1}{3}} = a^2b^2,$$

or, finally, dividing all the terms by $\frac{a^2b^2}{c^{\frac{4}{3}}}$, we obtain for the evolute the equation

$$(b\beta)^{\frac{2}{3}} + (a\alpha)^{\frac{2}{3}} = c^{\frac{4}{3}} = (a^2 - b^2)^{\frac{2}{3}}.$$



If $\alpha = 0$, then $\beta = \pm \frac{c^2}{b}$, so that the curve meets the axis of y in two points, c, d, equidistant from the origin O. If $\beta = 0$, then $\alpha = \pm \frac{c^2}{a}$, so that it also meets the axis of x in two points, b, a, equidistant from 0. If α is numerically greater than $\frac{c^2}{a}$ the ordinates become imaginary, and if β is numerically greater than $\frac{c^2}{b}$ the abscissas become imaginary; therefore the curve is limited by the four points a, b, c, d, and touches the axes at those points. It consists, therefore, of four branches symmetrically situated as in the figure.

3. To determine the evolute of the rectangular hyperbola, its equation between the asymptotes being $xy = a^2$.

The equation of the evolute is

$$(\beta + \alpha)^{\frac{2}{3}} - (\beta - \alpha)^{\frac{2}{3}} = (4a)^{\frac{2}{3}}.$$

THEOREM.

(104.) Normals to the curve are tangents to the evolute.

Let the equations of the curve and of its evolute be

$$y = F(x) \text{ and } \beta = \Phi(\alpha).$$

The normal to the curve is

$$y - y' = -\frac{1}{p'}(x - x').$$

The tangent to the evolute is

$$\beta - \beta' = \frac{d\beta}{d\alpha}(\alpha - \alpha').$$

These lines both pass through the point (α, β) since the centre of curvature is always on the normal. We have therefore to prove that

$$\frac{d\beta}{d\alpha} = -\frac{1}{p'}.$$

In order to this let us put for shortness $\frac{p'^2+1}{p''} = f$, then, (page 165,)

$$\alpha = x - p'f, \qquad \beta = y + f,$$

$$\therefore \frac{d\alpha}{dx} = 1 - p''f - p'\frac{df}{dx} = -p'^2 - p'\frac{df}{dx},$$

$$\frac{d\beta}{dx} = p' + \frac{df}{dx} \quad \therefore \quad \frac{d\beta}{d\alpha} = -\frac{1}{p'},$$

which was to be proved.

THEOREM.

(105.) The difference of any two radii of curvature is equal to the arc of the evolute comprehended between them.

Differentiating the equation

$$(y-\beta)^2 + (x-\alpha)^2 = r^2,$$

on the hypothesis that α is the independent variable, we have

$$(y-\beta)\frac{dy}{d\alpha} + (x-\alpha)\frac{dx}{d\alpha} - (y-\beta)\frac{d\beta}{d\alpha} - (x-\alpha) = r\frac{dr}{d\alpha}.$$

But equation (2), page 164,

$$(y-\beta)\frac{dy}{d\alpha} + (x-\alpha)\frac{dx}{d\alpha} = 0;^*$$

---

* It will be remembered that

$$p' = \frac{dy}{dx} = \frac{\frac{dy}{d\alpha}}{\frac{dx}{d\alpha}}$$ (See p. 127).

hence

$$-(y-\beta)\frac{d\beta}{d\alpha}-(x-\alpha)=r\frac{dr}{d\alpha};$$

but by last article,

$$y-\beta=(x-\alpha)\frac{d\beta}{d\alpha},$$

$$\therefore (x-\alpha)^2\left(\frac{d\beta^2}{d\alpha^2}+1\right)=r^2 \quad \ldots \quad (1).$$

and

$$-(x-\alpha)\left(\frac{d\beta^2}{d\alpha^2}+1\right)=r\frac{dr}{d\alpha} \quad \ldots \quad (2).$$

Dividing (2) by the square root of (1), we have

$$-\sqrt{\frac{d\beta^2}{d\alpha^2}+1}=\frac{dr}{d\alpha},$$

that is (87),

$$-\frac{ds}{d\alpha}=\frac{dr}{d\alpha} \therefore -s=r\pm \text{a constant},$$

for otherwise there could not be $-\frac{ds}{d\alpha}=\frac{dr}{d\alpha}$.

Hence if r, r' be the radii of curvature of any two points, and s, s' the corresponding arcs of the evolute, then

$$r\pm\text{const}=-s$$

$$r'\pm\text{const}=-s'$$

$$r-r'=s'-s,$$

so that the difference of the two radii is equal to the arc of the evolute comprehended between them; therefore, if a string fastened to one extremity of this arc be wrapped round it and continued in the direction of the tangent at the other extremity as far as the involute curve, the portion of the string thus coinciding with the tangent will by

(104) be the radius of curvature at that point P of the involute curve which it meets, and, consequently, by the above property, if the string be now unwound, P will trace out the involute.

On Consecutive Lines and Curves.

(106.) Every equation between two variables may always be considered as the analytical representation of some plane curve, given in species by the degree of the equation, and determinable both in form and position by the constants which enter it, provided, that is, these constants are fixed and determinate. If, however, the equation contains an arbitrary or indeterminate constant α, then, by assuming different values for α, the equation will represent so many different curves varying in form and position, but all belonging to the same *family of curves*.

Now if we consider the form and position of one of these curves to be fixed by the condition $\alpha = \alpha'$, another, intersecting this in some point, (x', y'), may be determined from a new condition $\alpha = \alpha' + h$; and if h be continually diminished, this latter curve will approach more and more closely to the fixed curve, and will at length coincide with it. During this approach, the point of intersection (x', y') necessarily varies, and becomes fixed in position only when the varying curve becomes coincident with the fixed curve. In this position the point is said to be the intersection of *consecutive curves*, so that what mathematicians call consecutive curves, are, in reality, coincident curves, and the point which has been denominated their point of intersection may be determined as follows:

(107.) Let

$$F(x, y, \alpha') = 0 \ldots . (1)$$

represent any plane curve, α' being a parameter, and for any intersecting curve of the same family let α' become $\alpha' + h$, then, since however numerous these intersecting curves may be, the x, y of the intersections belong also to the equation (1); it follows that as far as these points are concerned, the only quantity in equation (1) which varies is α', therefore, considering x, y as constants in reference to these points, we

have, by Taylor's theorem,

$$F(x, y, x' + h) = F(x, y, x') + \frac{dF(x, y, x')}{dx'} h +$$

$$\frac{d^2F(x, y, x')}{dx'^2} \frac{h^2}{1 \cdot 2} + \&c.$$

but $F(x, y, x') = 0$, therefore

$$\frac{F(x, y, x' + h)}{h} = \frac{dF(x, y, x')}{dx'} + \frac{d^2F(x, y, x')}{dx'^2} \frac{h}{1 \cdot 2} + \&c.$$

but $F(x, y, x' + h) = 0$, therefore

$$0 = \frac{dF(x, y, x')}{dx'} + \frac{d^2F(x, y, x')}{dx'^2} \frac{h}{1 \cdot 2} + \&c.$$

hence, when the curves are consecutive, that is, when $h = 0$, we have the following conditions, viz.

$$\left.\begin{aligned} F(x, y, x') &= 0 \\ \frac{dF(x, y, x')}{dx'} &= 0 \end{aligned}\right\} \cdots\cdots (2),$$

to determine x and y.

Suppose, for example, it were required to determine the point of intersection of consecutive normals in any plane curve.

Representing the equation of the curve by

$$y' = F(x'),$$

and any point in the normal by (x, y), we have for the equation of the normal,

$$y - y' = -\frac{1}{p'}(x - x') \text{ or } (y - y')p' + x - x' = 0.$$

This corresponds to the first of equations (2), x' being the parameter; hence, differentiating with respect to x', of which y' is a function given by the equation of the curve, we have

$$(y-y')\,p'-p'^2-1=0,$$

$$\therefore y=y'+\frac{p'^2+1}{p'},$$

$$\therefore x=x'-\frac{p'(p'^2+1)}{p'};$$

hence (93) consecutive normals intersect at the centre of curvature.

(108.) If we eliminate the variable parameter x' by means of the equations (2), the resulting equation will belong to every point of intersection given by every curve of the family

$$u=\mathrm{F}(x,y,x')=0 \ldots\ldots (1),$$

and its consecutive curve; for whatever value we suppose x' to take in the equations (2), the result of the elimination will obviously be always the same. Hence this resulting equation represents the locus of all the intersections, and we may show that at these same intersections this locus touches every individual curve in the family. The equation (1), where x' represents a function of x, y, determined by the second of the conditions (2) in last article, is obviously the equation of the locus of which we are speaking, and the same equation, when x' takes all possible values from 0 to $\pm\infty$, furnishes the family of curves, which we are now to show are all touched by this locus. Taking any one of this family, and differentiating its equation (1), x' being constant, we have

$$\mathrm{d}u=\frac{\mathrm{d}u}{\mathrm{d}x}\,\mathrm{d}x+\frac{\mathrm{d}u}{\mathrm{d}y}\,\mathrm{d}y=0.$$

Differentiating also the equation (1) of the locus, x' being given by the second condition of (2) in last article, we have

$$\mathrm{d}u=\frac{\mathrm{d}u}{\mathrm{d}x}\,\mathrm{d}x+\frac{\mathrm{d}u}{\mathrm{d}y}\,\mathrm{d}y+\frac{\mathrm{d}u}{\mathrm{d}x'}\,\mathrm{d}x'=0,$$

but by the condition just referred to $\dfrac{\mathrm{d}u}{\mathrm{d}x'}=0$, at the point where the curves whose equations we have just differentiated meet in this conse-

cutive position; hence, since at those points each of these equations give the same value for $\frac{dy}{dx}$, it follows that they have contact of the first order; we infer, therefore, that the equation (1), when x' is determined from the second of the conditions (2), last article, *represents a curve which touches and envelopes, or is enveloped by, the entire family of curves represented by the equation* (1), x' being any arbitrary constant. Thus, as we already know, the locus of the intersections of normals with their consecutive normals is a curve which touches them all at their points of intersection, being the evolute of the curve to which the normals belong.

The following examples will further illustrate this theory.

EXAMPLES.

(109.) 1. To determine the curve which touches an infinite series of equal circles, whose centres are all situated on the same circumference.

Let the equation of the fixed circle be

$$\alpha^2 + \beta^2 = R^2 \ldots\ldots (1,)$$

the centre of that circle being the origin of the coordinates.

The equation of the moveable circle, (α, β) being the varying centre, will be

$$(x-\alpha)^2 + (y-\beta)^2 - r^2 = 0.$$

But as (α, β) is always on the fixed circumference (1), we must always have between α and β the relation

$$\alpha^2 + \beta^2 = R^2 \therefore \beta = \sqrt{R^2 - \alpha^2}.$$

Hence by substitution,

$$(x-\alpha)^2 + (y - \sqrt{R^2-\alpha^2})^2 - r^2 = 0 = u \ldots\ldots (2).$$

Here α is the variable parameter, and, consequently, the equation represents the entire series of moveable circles; if, therefore, we differentiate

this equation with respect to a, and equate the result, $\frac{du}{da}$, to zero, this new equation, combined with (2), will enable us to eliminate a, and the result of this elimination will, by the preceding theory, be the analytical representation of the curve which touches all the moveable circles at those points where each is intersected by its consecutive circle. Now

$$\frac{du}{da} = -(x-a) + (y - \sqrt{R^2-a^2})\,(R^2-a^2)^{-\frac{1}{2}}a$$

$$= -(x-a) + \frac{ay}{\sqrt{R^2-a^2}} - a = 0,$$

$$\therefore -x\sqrt{R^2-a^2} + ay = 0,$$

$$\therefore x^2(R^2-a^2) = a^2y^2,$$

$$\therefore R^2x^2 = a^2(x^2+y^2),$$

$$\therefore a = \frac{Rx}{\sqrt{x^2+y^2}};$$

this substituted in (2), or in its development,

$$x^2 - 2ax + y^2 - 2y\sqrt{R^2-a^2} + R^2 = r^2,$$

gives

$$x^2 + y^2 - \frac{2Rx^2}{\sqrt{x^2+y^2}} - 2y\sqrt{\{R^2 - \frac{R^2x^2}{x^2+y^2}\}} + R^2 = r^2,$$

or

$$x^2 + y^2 - \frac{2R(x^2+y^2)}{\sqrt{x^2+y^2}} + R^2 = r^2,$$

or

$$x^2 + y^2 - 2R\sqrt{x^2+y^2} + R^2 = r^2,$$

whence

$$\sqrt{x^2+y^2} - R = \pm r,$$

and, consequently,

$$x^2 + y^2 = (R \pm r)^2,$$

an equation representing *two* circles, whose radii are respectively $R + r$

and $R - r$. Hence the series of circles are touched and enveloped by two circular arcs, having these radii, and the same centre as the fixed circle.

2. Between the sides of a given angle are drawn an infinite number of straight lines, so that the triangles formed may all have the same surface; required the curve to which every one of these lines is a tangent.

Let the given angle be θ, and, taking its sides for axes, we have, for the equation of every variable line,

$$y = \alpha x + \beta \ldots . (1),$$

and, putting successively, $y = 0$ and $x = 0$, the resulting expressions for x and y denote the sides of the variable triangle, including the given angle, so that these sides are $-\frac{\beta}{\alpha}$ and β: hence, calling the constant surface s, we have

$$s = -\frac{\beta^2}{2\alpha} \sin\theta \therefore \alpha = -\frac{\beta^2 \sin\theta}{2s};$$

hence the equation (1) is the same as

$$y = -\frac{\beta^2 \sin\theta}{2s} x + \beta \ldots . (2),$$

where β is considered as an arbitrary constant. But if for this arbitrary constant we substitute the function of x, arising from the condition $\frac{dy}{d\beta} = 0$, then (2) will represent the locus of the intersections of each variable line, with its consecutive line, which locus touches them all. Differentiating then with regard to β, we have

$$-\frac{\beta \sin\theta}{s} x + 1 = 0 \therefore \beta = \frac{s}{x \sin\theta},$$

this substituted in (2), gives, for the equation of the sought curve,

$$y = -\frac{s}{2x^2 \sin\theta} x + \frac{s}{x \sin\theta},$$

or rather

$$xy = \frac{s}{2 \sin\theta};$$

hence the curve is an hyperbola, having the sides of the given angle for asymptotes.

3. The centres of an infinite number of equal circles are all situated on the same straight line: required the line which touches them all.

Ans. They are touched by two parallels to the line of centres.

4. From every point in a parabola lines are drawn, making the same angle with the tangent that the tangent makes with the diameter at the point; required the line touching them all?

Ans. They are touched by a point, viz. the focus, in which therefore they all meet.

CHAPTER IV.

ON THE SINGULAR POINTS OF CURVES, AND ON CURVILINEAR ASYMPTOTES.

Multiple Points.

(110.) If several branches of a curve meet in one point, whether by intersecting or touching each other, that point is called a multiple point. In the former case the point is said to be of the first species, and in the latter of the second species, and we propose here to enquire how, by means of the equation of a curve, these points, if any, may be detected.

Multiple points of the first species. When the curve has multiple points of the first species, we readily arrive at the means of determining their position from the consideration that at such points there must be as many rectilinear tangents as there are touching branches, and, consequently, as many values for $\frac{dy}{dx}$, the tangent of the inclination of

any tangent through the point (x, y) to the axis of x; so that the equation of the curve being freed from radicals and put under the form

$$F(x, y) = 0,$$

its multiple points of the first species will all be given analytically by the equation

$$p' = -\frac{du}{dx} \div \frac{du}{dy} = \frac{0}{0},$$

so that no systems of values for x and y can belong to multiple points of the first species, but such as satisfy the conditions

$$\frac{du}{dx} = 0, \quad \frac{du}{dy} = 0,$$

as well as the equation of the curve. Having, therefore, determined all such systems of values by solving the two last equations, the true values of p' for each system will be ascertained by proceeding as in (41), and those systems only will belong to multiple points of the first species that give multiple values to p'. Let us apply this to an example or two.

EXAMPLES.

(111.) 1. To determine whether the curve represented by the equation

$$ay^3 - x^3y - bx^3 = 0,$$

has any intersecting branches,

$$\frac{du}{dx} = -3x^2(y + b), \quad \frac{du}{dy} = 3ay^2 - x^3.$$

At the points where branches intersect we must have

$$3x^2(y + b) = 0, \quad 3ay^2 - x^3 = 0,$$

$$\therefore x = 0, \quad y = 0,$$

or

$$x = \sqrt[3]{3ab^2}, \quad y = -b;$$

this second system of coordinates do not satisfy the proposed equation, and therefore do not mark any point in the curve; the first system, which is admissible, shows that if there exist any multiple point it must be at the origin. Hence, to ascertain the true value of p' at this point, we have, by differentiating both numerator and denominator in the expression for p',

$$[p'] = [\frac{3x^2(y+b)}{3ay^2 - x^3}] = \frac{0}{0}$$

$$= [\frac{6x(y+b) + 3x^2p'}{6ayp' - 3x^2}] = \frac{0}{0}$$

$$= [\frac{6(y+b) + 12p'x + 3x^2p''}{6ayp'' + 6ap'^2 - 6x}] = \frac{6b}{6a[p']^2},$$

$$\therefore [p'] = \sqrt[3]{\frac{b}{a}};$$

therefore, as this has but one *real* value, the curve has no intersecting branches.

2. To determine whether the curve represented by the equation

$$x^4 + 2ax^2y - ay^3 = 0$$

has intersecting branches

$$\frac{du}{dx} = 4x(x^2 + ay) = 0, \quad \frac{du}{dy} = a(2x^2 - 3y^2) = 0.$$

There is but one system of values that can satisfy these three equations, viz.

$$x = 0, \quad y = 0;$$

so that if there are intersecting branches they must intersect at the origin. To determine, therefore, whether at this point p' has multiple values, we have

$$[p'] = [\frac{4x(x^2 + ay)}{a(3y^2 - 2x^2)}] = \frac{0}{0}$$

$$= [\frac{6x^2 + 2ay + 2axp'}{3ayp' - 2ax}] = \frac{0}{0}$$

$$= \frac{4a[p']}{3a[p']^2 - 2a},$$

$$\therefore\ 3a\,[p']^3 - 6a\,[p'] = 0,$$

$$\therefore\ [p'] = 0, \quad \text{or } [p'] = \pm\sqrt{2};$$

hence *three* branches of the curve intersect at the origin; the tangent to one of them at that point is parallel to the axis of x, and the tangents to the other two are symmetrically situated with respect to the axis of y, since they are inclined to the axis of x, at angles whose tangents are $+\sqrt{2}$ and $-\sqrt{2}$.

(112.) Should the values of p', corresponding to any values of x and y, which satisfy the equation of the curve, be all imaginary, we must infer that, although such a system of values belong to a point of the locus, yet that point must be detached from the other points of the locus; for since, if the abscissa of this point be increased by h, the development of the ordinate will agree with Taylor's development, as far, at least, as the second term for all values of h, between some finite value and 0, it follows that all the corresponding ordinates between these limits must be imaginary, so that the proposed point is isolated, having no geometrical connexion with the curve, although its coordinates satisfy the equation. Such a point is called *a conjugate point.*

(113.) From what has now been said, it appears that, by having the equation of a plane curve given, those points in it where branches intersect, as also those which are entirely detached from the curve, although belonging to its equation, may always be determined by the application of the differential calculus, and independently of all considerations about the failing cases of Taylor's theorem, except, indeed, those connected with the theory of vanishing fractions. We shall now seek the analytical indications of

Multiple Points of the Second Species.

(114.) The several species of multiple points, or those where branches of the curve touch each other, the differential calculus does not furnish the means of readily determining from the implicit equation of the

curve. We know that at such a point, p' cannot admit of different values, since the branches have one common tangent, and we know, moreover, that if Taylor's theorem does not fail at that point, we shall, by successively differentiating, at length arrive at a coefficient which, being put under the form $\frac{0}{0}$, the different values will indicate so many different touching branches; for if no coefficient gave multiple values for the proposed coordinates x', y', then the ordinates corresponding to the abscissas between the limits x' and $x \pm h$, h being some finite value, would each have but one value, and, therefore, different branches could not proceed from the point (x', y'). But we have no means of ascertaining *à priori* which of the coefficients furnishes the multiple value. When, however, the equation of the curve is explicit, then the multiple points of either species are very easily determined. Thus, if the equation of the curve be

$$y = (x-a)^{\frac{3}{2}}\sqrt{x-b} + c,$$

we at once see that $x = a$ destroys the radical in y and p', that reappears in p''; therefore, at the point corresponding to this abscissa, there will be but one tangent, and yet two branches of the curve proceed from it on account of the double value of p''. Hence the point is a *double* point of the second species, the branches have contact of the first order, and, because $p' = 0$, the common tangent is parallel to the axis; if the radical had been of the third degree, the point corresponding to the same abscissa would have been a *triple* point, &c. It appears, therefore, that when the equation of the curve is solved for y, there will exist a multiple point, if in the expression for $\frac{y}{x}$ a radical is multiplied by the factor $(x-a)^m$. If $m = 1$, the branches of the curve intersect at the point whose abscissa is $x = a$, because then p' at that point takes the same values as the radical, but if $m > 1$, then the branches touch, because then the radical is destroyed in p' for $x = a$; in both cases the index of the radical will denote the number of branches which meet in the point. Such, therefore, are the geometrical significations of the cases discussed in (75) and (76).



Cusps, or Points of Regression.

(115.) A *cusp*, or *point of regression*, is that particular kind of *double* point of the second species in which the two touching branches terminate, and through which they do not pass, so that on one side of such a point, viz. on that where the branches lie, the ordinate has a double value, and on the other side the contiguous ordinate has an imaginary value.



The cusp represented in the first figure, where the branches are one on each side the common tangent, is called a cusp of the first kind, and that in the second figure, where the branches lie both on one side, a cusp of the second kind.

(116.) It is obvious that cusps can exist only at those points, the particular coordinates of which cause Taylor's theorem to fail; for if Taylor's theorem did not fail at such a point, then the ordinates in the vicinity, corresponding both to $x + h$ and to $x - h$, would be both possible or impossible at the same time. We are not, however, to infer that when the adjacent ordinates are on the one side of any point real, and on the other side imaginary, that a cusp *necessarily* exists at that point; for it is plain that the same analytical indications are furnished by the point which limits *any* curve in the direction of the axis of x, or at which the tangent is perpendicular to that axis, as in the third figure. It becomes important, therefore, in seeking particular points of curves, to be able to distinguish the point which limits the curve in the direction of the axes from cusps.

(117.) Now, at the limits, the tangents to the curve are parallel to the axes, the limits are therefore determined by the equations $\frac{dy}{dx} = \infty$ and $\frac{dy}{dx} = 0$; and they fulfil, moreover, the following additional con-

ditions, viz. 1°, the ordinate, or abscissa, whichever it may be, that is parallel to the tangent, immediately *beyond* the limit, must be imaginary; but if it be ascertained that this is not the case, the point is not a limit but a cusp of the first kind, posited as in the annexed figures, or else a point of inflexion; the latter when the contiguous ordinates are the one greater, and the other less than that at the point. 2°, Besides the first condition, there must exist also this, viz. that immediately *within* the limit the double ordinate or abscissa, whichever may be parallel to the tangent, must have one of its values greater and the other less than at the point; but if both are greater or both less, the point is not a limit but a cusp of the second kind, posited as in the annexed figures. Hence, when the branches forming the cusp touch the abscissa or the ordinate of the point, they may be discovered by seeking among the values which satisfy the equations $\frac{dy}{dx} = 0$



and $\frac{dy}{dx} = \infty$, those which do not fulfil both the foregoing conditions.

Let us illustrate this by examples.

EXAMPLES.

(118.) 1. To determine whether the curve whose equation is

$$(y-b)^3 = (x-a)^2$$

has a cusp at the point where the tangent is parallel to the axis of y.

By differentiating

$$\frac{dy}{dx} = \frac{2}{3} \cdot \frac{x-a}{(y-b)^2},$$

this becomes infinite for $y = b$, therefore the point to be examined is (a, b). In order to this, substitute $a \pm h$ for x in the proposed equation, and we have, for the contiguous ordinates,

$$y = b + h^{\frac{3}{3}},$$

which is not imaginary either for $+ h$ or $- h$; the point (a, b) is therefore a cusp of the first kind, and posited as in the figure, since the contiguous values of y are both greater than b.



2. To determine whether the curve whose equation is

$$y - a = (x - b)^{\frac{1}{3}} + (x - b)^{\frac{2}{4}}$$

has a cusp at the point where the tangent is parallel to the axis of y.

Here the coefficient $\frac{dy}{dx}$ becomes infinite for $x = b$, therefore the point to be examined is (b, a). Substituting $b + h$ for x, we have

$$y = a + h^{\frac{1}{3}} + h^{\frac{2}{4}}.$$

For negative values of h this is imaginary, therefore the curve lies entirely to the right of the ordinate $y = a$, so that the condition 1° pertaining to a limit is fulfilled. To the right of this ordinate the two values of y, corresponding to a value of h, ever so small, are *both* greater than $y = a$, so that the condition 2° is not fulfilled; the point (b, a) is therefore a cusp of the second kind, and posited as in the cut.



3. To determine the point of the curve whose equation is

$$(y - a - x)^4 = (x - b)^3,$$

at which the tangent is parallel to the axis of y.

The differential coefficient becomes infinite for $x = b$, therefore the point to be examined is $(b, a + b)$. Substituting $b + h$ for x,

$$y = (a + b) + h^{\frac{3}{4}} + h,$$

negative values of h render this imaginary, therefore the condition 1° is fulfilled; positive values give two values for y, and as h may be taken so small that $h^{\frac{3}{4}}$ may exceed h, and since, moreover, the two values of $h^{\frac{3}{4}}$ are the one positive and the other negative, it follows that the real ordinate contiguous to the point has one value greater, and

the other less, than that at the point of contact; hence the condition 2° is also fulfilled, and thus the point marks the limit of the curve, which, therefore, lies to the right of the ordinate, through $x = b$

(119.) Having thus seen how to determine those cusps where the branches touch an ordinate or abscissa, we shall now seek how to discover those at which the tangent is oblique to the axes. The true development of the ordinate contiguous to such a cusp must be of the form

$$y' + \frac{dy'}{dx'} h + Ah^{\alpha} + \beta h^{\beta} + \&c.$$

and the corresponding ordinate of the tangent will be

$$y' + \frac{dy'}{dx'} h;$$

hence, subtracting this from the former, and calling the difference Δ, we have

$$\Delta = Ah^{\alpha} + \beta h^{\beta} + \&c.$$

(120.) Now, in order that the point $(x'\ y')$ may be a cusp, this difference, for a small value of h, must have two values; and to be a cusp of the first kind, these two values must obviously have opposite signs; but since h may be so small that Ah^{α} may exceed the sum of all the following terms, h^{α} must have two opposite values; hence, α must be a fraction with an *even* denominator, and, conversely, if α be a fraction with an even denominator, the point $(x',\ y')$ will be a cusp of the first kind. Hence, at such a point, $\frac{d^2y}{dx^2}$ is either 0 or ∞ : 0 if $\beta > 2$, and ∞ if $\beta < 2$.

(121.) In order that the cusp may be of the second kind, both values of Δ must have the same sign, therefore h^{α} cannot admit of opposite values for the same value of h, consequently α must in this case be either a whole number, or else a fraction with an *odd* denominator;

and conversely, if α be either a whole number or a fraction with an odd denominator, the point (x', y) will be a cusp of the second kind, provided, of course, that Δ has *two* values. The position of the branches will depend on the sign of A.

We shall now give an example or two.

(122.) 4. To determine whether the curve whose equation is

$$y = x \pm x^{\frac{2}{2}}$$

has a cusp.

Here y is possible for positive values of x, and imaginary for all negative values; hence there *may* be a cusp at the origin. To ascertain this, put h for x, in the equation, and we have, for the contiguous ordinate, the value

$$y = h \pm h^{\frac{2}{2}}.$$

The coefficient of h being $1 = \frac{dy'}{dx'}$, we see that the tangent to the curve at the origin is inclined at 45° to the axes, and, since $\frac{2}{2}$ has an even denominator, the origin is a cusp of the first kind.

5. To determine whether the curve whose equation is

$$y - a = x + bx^3 + cx^{\frac{5}{2}}$$

has a cusp.

Here y is imaginary for all negative values of x, therefore the point $(0, a)$ *may* be a cusp. Substituting h for x, we have

$$y = a + h + bh^2 + ch^{\frac{5}{2}}.$$

As before, the tangent is inclined at 45° to the axes, and, since the exponent of the third term is a whole number, and the whole expression admits of two values, in consequence of the even root $h^{\frac{5}{2}}$, it follows that the proposed point is a cusp of the second kind. The branches are situated to the right of the axis of y, because h must be positive, and they are above the tangent because bh^2 is positive.

6. To determine whether the curve whose equation is

$$(2y + x + 1)^2 = 2(1 - x)^3$$

has a cusp.

Here values of x greater than 1 are obviously inadmissible, and to the value $x=1$ corresponds $y=-1$; hence the point having these coordinates *may* be a cusp. Substituting $1+h$ for x, we have

$$y=-1+\tfrac{1}{2}h+h^{\frac{5}{3}},$$

therefore the tangent to the curve at the proposed point has the trigonometrical tangent of its inclination to the axis of x equal to $\frac{1}{2}$, and since the fraction $\frac{5}{3}$ has an even denominator, the point is a cusp of the first kind. Because h is negative, the branches are to the left of the ordinate to the point which is below the axis of x, because this ordinate is negative.



Points of Inflexion.

(123.) Points of inflexion have been defined at (93), and we have there shown that a point of this kind always exists when its abscissa causes all the differential coefficients to vanish between the first and the nth, provided the nth be odd and become neither 0 nor ∞. The simplest indication therefore of a point of inflexion is $[\frac{d^2y}{dx^2}]=0$, and $[\frac{d^3y}{dx^3}]$ neither 0 nor ∞; such indications, however, cannot be furnished by any point at which the tangent is parallel to the axis of y, since in this case $[\frac{dy}{dx}]$ and all the following coefficients become infinite. Neither can these indications take place at any point, for which Taylor's theorem fails after the third term. It becomes, therefore, of consequence, in examining particular points of a curve, to be able to detect the existence of points of inflexion by some general method, independently of the differential coefficients beyond the first. The only general method of doing this is that which we have already employed for the discovery of cusps, and which consists simply in examining the course of the curve in the immediate vicinity, and on each side the point in question. Points of inflexion are somewhat similar to cusps, each having some of the analytical peculiarities common to both, and to the limiting points of curves, as already hinted at in (116.) But the characteristic property

of a point of inflexion is, that the adjacent ordinates on each side are the one greater and the other less than the ordinate at the point. This peculiarity distinguishes a point of inflexion from a limit, inasmuch as at a limit the ordinate immediately beyond is imaginary; and it distinguishes it from a cusp of the first kind, inasmuch as at such a cusp the adjacent ordinates are either both greater or both less than at the point, or else, as is the case when the tangent at the point is oblique to the axes, one of these ordinates is imaginary, the other double. We have then first to ascertain at what points of the curve inflexions *may* exist, or to find what points are given by the conditions



$$\frac{d^2y}{dx^2} = \frac{P}{Q} = 0 \text{ or } \infty,$$

or, which is the same thing, what points are given by the separate conditions

$$P = 0, \quad Q = 0,$$

we are then, by examining the course of the curve in the vicinity of each point, to determine to which of them really belong the characteristic of an inflexion.

Thus the means of distinguishing points of inflexion being sufficiently clear, we shall proceed to a few examples.

EXAMPLES.

1. To determine whether the curve whose equation is

$$y = b + (x - a)^3$$

has a point of inflexion where the tangent is parallel to the axis of x.

Here

$$p' = 3(x - a)^2,$$

and when the tangent is parallel to the axis of x, $p' = 0$, $\therefore x = a$ and $y = b$, at the proposed point. In the vicinity, $x = a + h$,

$$\therefore y = b + h^3,$$

which is *greater* than b, the ordinate of the point when h is positive, and less when h is negative; the point (a, b) is therefore a point of inflexion.

2. To determine whether the curve whose equation is $y^3 = x^5$ or $y = x^{\frac{5}{3}}$ has an inflexion at any point.

$$p' = \tfrac{5}{3}x^{\frac{2}{3}},\ p'' = \tfrac{2}{3}\cdot\tfrac{5}{3}x^{-\frac{1}{3}},$$

this becomes ∞ for $x = 0$, therefore a point of inflexion *may* exist at the origin. Putting h for x, we have

$$y = h^{\frac{5}{3}},$$

which is greater than 0, the ordinate of the point, when h is positive, and less when h is negative; hence there is an inflexion at the origin. Also the equation of the tangent being $y = \frac{5}{3}x^{\frac{2}{3}}$, the ordinates corresponding to $x = \pm h$ are both less than those given by the above equation; hence the curve lies above the tangent to the right of the origin, and below it to the left, as in the figure.

3. To determine whether the curve whose equation is

$$y - x = (x - a)^{\frac{5}{3}}$$

has a point of inflexion

$$p' = 1 + \tfrac{5}{3}(x-a)^{\frac{2}{3}},\ p'' = \tfrac{2}{3}\cdot\tfrac{5}{3}(x-a)^{-\frac{1}{3}},$$

this becomes infinite for $x = a$, therefore a point of inflexion *may* exist at the point (a, a). In the vicinity of this point $x = a + h$,

$$\therefore\ y = a + h + h^{\frac{5}{3}},$$

which is greater than a when h is positive, and less when h is negative; hence (a, a) is a point of inflexion. As the corresponding ordinates of the tangent $y = a \pm h$, one, viz. $y = a + h$, is less than that of the curve, and the other greater, the curve bends, as in the figure.

On Curvilinear Asymptotes.

(124.) Two plane curves, having infinite branches, are said to be asymptotes to each other, when they approach the closer to each other as the branches are prolonged, but meet only at an infinite distance.*

Hence, since the expression for the difference of the ordinates corresponding to the same abscissa in two such curves becomes less and less, as the abscissa becomes greater and greater, and finally becomes 0, when the abscissa becomes ∞, it follows that that expression can contain none but *negative* powers of x, without the addition of any constant quantity. For, if a positive power of x entered the expression for the difference, that expression would become not 0 but ∞, when $x = \infty$, and, if there were a quantity independent of x, the difference would be reduced to this quantity, and not to 0, for $x = 0$. Hence two curves are asymptotes to each other, when the general expression for the difference of the ordinates corresponding to the same abscissa is

$$\Delta = A'x^{-\alpha} + B'x^{-\beta} + C'x^{-\gamma} + \&c. \ldots\ldots (1),$$

or when the general expression for the difference of the abscissas corresponding to the same ordinate is

$$\Delta = A'y^{-\alpha} + B'y^{-\beta} + C'y^{-\gamma} + \&c. \ldots\ldots (2),$$

and conversely, when the curves are asymptotes to each other; one or both these forms must have place.

If for one of the curves whose corresponding ordinates are supposed to give the difference (1) there be substituted another, which would reduce that difference to

$$B'x^{-\beta} + C'x^{-\gamma} + \&c.$$

this new curve would be an asymptote to both, and would obviously,

---

* Spirals meet their asymptotic circles only after an infinite number of revolutions; these we do not consider here, having adverted to them at (86).

throughout its course, continually approach nearer to that which it has been compared to, than the one for which we have substituted it does. In like manner, if a third curve would further reduce the difference (1) to

$$C'x^{-\gamma} + \&c.$$

this third curve would approach the first still nearer, and all the four would be asymptotes to each other. It appears, therefore, that every curve of which the ordinate may be expanded into an expression of the form

$$y = Ax^{a} + Bx^{b} + \ldots . A'x^{-\alpha} + B'x^{-\beta} + \&c. \ldots . (3).$$

admits of an infinite number of asymptotes.

Since the general expression for the ordinate of a straight line is $y = Ax + B$, for the difference between this ordinate and that of a curve at the point whose abscissa is x, to have the form (1), the equation of the curve must be

$$y = Ax + B + A'x^{-\alpha} + B'x^{-\beta} + \&c. \ldots . (4),$$

this equation, therefore, comprehends all the curves that have a rectilinear asymptote, and among them the common hyperbola, whose equation is

$$y = \pm \frac{B}{A}(x^2 - A^2)^{\frac{1}{2}} = \mp \frac{B}{A}x \mp \tfrac{1}{2}ABx^{-1} + \&c.$$

The curves included in the equation (4) are therefore called *hyperbolic curves.*

The other curves comprised in the more general equation (3), not admitting of a rectilinear asymptote, are called *parabolic curves.*

The common hyperbola, we see by the above equation, admits of the two rectilinear asymptotes $y = \pm \frac{B}{A}x$, and of an infinite number of hyperbolic asymptotes.

As an example of this method of discovering rectilinear and curvilinear asymptotes, let the equation

$$my^3 - xy^3 = mx^3$$

be proposed. The development of y in a series of descending powers of x, is (Ex. 9, p. 68,)

$$y = -m - \frac{m^4}{x^3} - \&c.$$

therefore the curve has one rectilinear asymptote, parallel to the axis of x, its equation being $y = -m$; the hyperbolic asymptote next to this, and which lies closer to the curve, is of the fourth order, its equation being

$$yx^3 + mx^3 + m^4 = 0.$$

Again, let the equation of the proposed curve be

$$y = \frac{b}{(x^2 - a^2)^{\frac{1}{2}}}$$

$$= bx^{-1} + \&c. \ldots . \quad (1);$$

also, since

$$x^2 - a^2 = \frac{b^2}{y^2} \therefore x = a + \tfrac{1}{2} \cdot \frac{b^2}{a} y^{-2} + \&c. \ldots . \quad (2).$$

From (1) it appears that the curve has a rectilinear asymptote, coincident with the axis of x, its equation being $y = 0$; the hyperbola whose asymptotes coincide with the axes is also an asymptote, its equation being $xy = b$. From (2) it appears that the curve has another rectilinear asymptote, parallel to the axis of y, its equation being $x = a$; the hyperbola next to this is of the third order. If we consider the radical, in the proposed equation, to admit of either a positive or a negative value, then there will be two rectilinear asymptotes, parallel to the axis of y and equidistant from it, as also two hyperbolic asymptotes, symmetrically situated between the axes.

SECTION III.

ON THE GENERAL THEORY OF CURVE SURFACES AND OF CURVES OF DOUBLE CURVATURE.

CHAPTER I.

ON TANGENT AND NORMAL PLANES.

PROBLEM I.

(125.) **To determine the equation of the tangent plane at any point on a curve surface.**

Let (x', y', z') represent any point on a curve surface of which the equation is

$$z = F(x, y),$$

then the tangent plane will obviously be determined, when two linear tangents through this point are determined. Let us then consider, for greater simplicity, the two linear tangents respectively parallel to the planes of xz, zy; their equations are

$$\left.\begin{aligned} z - z' &= a(x - x') \\ y &= y' \end{aligned}\right\} \quad \ldots\ldots (1)$$

and

$$\left.\begin{aligned} z - z' &= b(y - y') \\ x &= x' \end{aligned}\right\} \quad \ldots\ldots (2).$$

But since these are tangents to the plane curves, which are the sections through (x', y', z'), parallel to the planes of xz, zy, therefore (77)

$$a = \frac{dz'}{dx'} = p', \quad b = \frac{dz'}{dy'} = q'.$$

Moreover, the traces of the plane through the lines (1), (2), upon the planes of xz, zy, being parallel to the lines themselves, a and b must be the same in the traces as in these lines; and, since they are the same in the plane as in its traces, it follows that the equation of this plane must be

$$z - z' = p'(x - x') + q'(y - y') \ldots\ (3),$$

in which the partial differential coefficients p', q', express the trigonometrical tangents of the inclinations of the vertical traces to the axes of x and y respectively.

For the angle which the horizontal trace makes with the axis of x, we have, by putting $z = 0$, in (3),

$$\text{tangent of inclination} = \frac{p'}{q'}.$$

(126.) If the equation of the surface is given under the form

$$u = F(x, y, z) = 0 \ldots\ (4),$$

then the expressions for the total differential coefficients derived from u considered as a function, first of the single variable x, and then of the single variable y, are (57)

$$\left\{\frac{du}{dx}\right\} = \frac{du}{dx} + \frac{du}{dz} p' = 0$$

$$\left\{\frac{du}{dy}\right\} = \frac{du}{dy} + \frac{du}{dz} q' = 0,$$

from which we get the values

$$p' = -\frac{\frac{du}{dx}}{\frac{du}{dz}}, \quad q' = -\frac{\frac{du}{dy}}{\frac{du}{dz}};$$

hence, by substituting these expressions in (3), the equation for the tangent plane becomes

$$(z-z')\frac{du}{dz}+(x-x')\frac{du}{dx}+(y-y')\frac{du}{dy}=0 \quad \ldots \quad (5).$$

PROBLEM II.

(127.) To determine the equation of the normal line at any point of a curve surface.

We have here merely to express the equation of a straight line, perpendicular to the plane (3), and passing through the point of contact (x', y', z').

Now the projections of this line must be perpendicular to the traces of the tangent plane, or to the lines (1), (2); hence the equations of these projections must be

$$\left.\begin{aligned} x-x'+p'(z-z')&=0\\ y-y'+q'(z-z')&=0 \end{aligned}\right\},$$

which together, therefore, represent the normal.

(128.) If we represent by α, β, γ, the inclinations of this line to the axes of x, y, z, respectively, then (*Anal. Geom.*, p. 226,)

$$\cos\alpha=\frac{-p'}{\sqrt{p'^2+q'^2+1}}$$

$$\cos\beta=\frac{-q'}{\sqrt{p'^2+q'^2+1}}$$

$$\cos\gamma=\frac{1}{\sqrt{p'^2+q'^2+1}}.$$

(129.) If the equation of the surface be given under the form (4), last problem, then, in these expressions for the inclinations, we must,

instead of p' and q', write their values as before determined from that equation. If, for brevity, we put

$$v = \frac{1}{\sqrt{\frac{du^2}{dx^2} + \frac{du^2}{dy^2} + \frac{du^2}{dz^2}}},$$

the expressions for the cosines will then be

$$\cos\alpha = v\frac{du}{dx}, \quad \cos\beta = v\frac{du}{dy}, \quad \cos\gamma = v\frac{du}{dz}.$$

As every plane which contains the normal line must be perpendicular to the tangent plane, it is obvious that there exists an infinite number of normal planes to any point of a surface.

PROBLEM III.

(130.) To determine the equation of the tangent line to any point of a curve of double curvature.

We have already indicated (*Anal. Geom.*, p. 311,) how this equation is to be determined:

Let

$$y = f(x), \quad z = F(x) \ldots . (1),$$

be the equations of the projections of the proposed curve, on the planes of xy, xz; and let (x', y', z') be the point to which the linear tangent is to be drawn, which point will be projected into (x', y') and (x', z'), on the plane curves (1); therefore tangents through them to these plane curves, will be represented by the two equations

$$\left.\begin{aligned} y - y' &= p'(x - x') \\ z - z' &= q'(x - x') \end{aligned}\right\} \ldots . (2);$$

these, therefore, together, represent the required tangent in space.

PROBLEM IV.

(131.) To determine the equation of the normal plane at any point in a curve of double curvature.

The equation of any plane passing through a proposed point, is *(Anal. Geom.*, p. 231,)

$$A(x-x')+B(y-y')+C(z-z')=0 \quad \ldots \quad (1);$$

and for the traces of this plane on the planes of xy, xz, we have, by putting in succession $z=0$, $y=0$, the equations

$$y-y'=-\frac{A}{B}(x-x')+\frac{C}{B}z',$$

$$z-z'=-\frac{A}{C}(x-x')+\frac{B}{C}y',$$

but since these two traces are respectively perpendicular to those marked (2), last problem,

$$\therefore \frac{B}{A}=p', \quad \frac{C}{A}=q';$$

hence the equation (1) becomes

$$x-x'+p'(y-y')+q'(z-z')=0 \quad \ldots \quad (2),$$

which represents the normal plane sought.

CHAPTER II.

ON CYLINDRICAL SURFACES, CONICAL SURFACES, AND SURFACES OF REVOLUTION.

(132.) These surfaces have been considered in the *Analytical Geometry*, and the general equations of the first two classes have been deduced on the hypothesis that the directrix is always a plane curve. We shall now suppose the directrix to be any curve situated in space, and investigate the differential equations of these surfaces, as also of surfaces of revolution in general.

Cylindrical and Conical Surfaces.

PROBLEM I.

To determine the equation of cylindrical surfaces in general.

Let the equations of the generating straight line be

$$\left.\begin{aligned} x &= az + \alpha \\ y &= bz + \beta \end{aligned}\right\} \quad \therefore \quad \left\{\begin{aligned} \alpha &= x - az \\ \beta &= y - bz \end{aligned}\right. \quad \ldots\ (1),$$

and the equations of any curve in space, considered as the directrix,

$$F(x, y, z) = 0, \quad f(x, y, z) = 0 \quad \ldots\ (2).$$

Now for every point in this directrix, all these equations exist simultaneously; moreover, the constants a, b, are fixed, since the inclination of the generating line does not vary, but the constants α, β, are not fixed, since the *position* of the generating line does vary. If, then, we eliminate x, y, z, from the above equations, there will enter, in the resulting equation, only the constants a, b, and the indeterminates α, β; hence, solving this equation, for β we shall get a result of the form $\beta = \phi(\alpha)$; consequently, if we now substitute in this the values of α and β given

above, in terms of x, y, z, we shall have this general relation among these variables, viz.

$$y - bz = \phi(x - az) = 0 \ldots\ldots (3),$$

which is the equation of cylindrical surfaces in general, the function ϕ depending entirely on the nature of the directrix, and a, b being given by the generatrix.

(133.) Now, by differentiation, this function may be eliminated (58), hence,

$$\frac{-bp'}{1-bq'} = \frac{1-ap'}{-aq'},$$

$$\therefore\ ap' + bq' = 1 \text{ or } a\frac{dz}{dx} + b\frac{dz}{dy} = 1 \ldots\ldots (4),$$

which is the general differential equation of cylindrical surfaces.

(134.) The same equation may be immediately deduced from the general equation of a tangent plane, to the cylindrical surface. Thus, the equation of any tangent plane, through a point (x', y', z') being

$$z - z' = p'(x - x') + q'(y - y'),$$

the condition necessary for it to be *always* tangent to the cylinder on which this point is situated, is merely that it may be always parallel to its generatrix (1), and this condition, expressed analytically, is (*Anal. Geom.* p. 232,)

$$ap' + bq' - 1 = 0 \ldots\ldots (4);$$

this is, therefore, the relation which must have place between the partial differential coefficients derived from the equation of the surface, in order that that surface may be cylindrical, and it agrees with the relation before established.

If, in this equation, we write for p', q', their values deduced from the equation $u = 0$ of any cylindrical surface, as exhibited in (126), it becomes

$$a\frac{du}{dx} + b\frac{du}{dy} + \frac{du}{dz} = 0 \ldots\ldots (5).$$

PROBLEM II.

(135.) Given the equation of the generatrix, to determine the cylindrical surface which envelopes a given curve surface.

Since the cylinder envelopes the given surface, the curve of contact is common to both, therefore every tangent plane to the cylinder touches the enveloped surface in that curve. The equation of any of these tangent planes is

$$z - z' = p'(x - x') + q'(y - y'),$$

whether p' and q' be derived from the equation of the surface, and take those particular values which restrict them to the curve of contact, or whether p' and q' be derived from the equation of the cylinder, and preserve their general values; because in the one case the contact of each tangent is confined to a point in the curve of contact, and in the other case the contact extends along the whole length of the cylinder. Hence, for the curve of contact, the condition (5) must have place, as well as for the entire surface of the cylinder. The mode of solution is, therefore, obvious; we must deduce p' and q' from the equation of the given surface, and substitute them in (5); the result, combined with the equation of the given surface, will obviously represent the curve for which p' and q' are common to both surfaces; that is to say, we shall thus have the equations of the directrix, and that of the generatrix being also given, the particular cylindrical surface becomes determined.

(136.) If the proposed curve surface be of the second order, then the equation (5) will necessarily be of the first degree in x, y, z, and will, therefore, represent a plane; so that the combination of this, with the equation of any surface, must necessarily represent a plane section of that surface; we infer, therefore, that if any cylindrical surface circumscribe a surface of the second order, the curve of contact will always be a plane curve, and consequently of the second order, and therefore the cylinder itself must be of the second order.

As an example, let it be required to determine the cylindrical surface having the generatrix (1) in Problem I., and enveloping the ellipsoid, whose equation is

$$u = Ax^2 + By^2 + Cz^2 - 1 = 0 \quad \ldots \quad (1).$$

The directrix being on this surface, is included in the equation (1), and the same directrix being on the enveloping cylinder, is included in the equation (5) of last problem; hence the combination of these two equations represents the directrix itself to the exclusion of all other points; that is, the equations of the directrix are

$$\left.\begin{aligned} Ax^2 + By^2 + Cz^2 - 1 &= 0 \\ Aax + Bby + Cz &= 0 \end{aligned}\right\} \quad \ldots \quad (2),$$

which equations correspond with those marked (2) in Problem I.; and to get the cylindrical surface sought, it remains for us to proceed as directed in Problem I. Eliminating x, y, z, from the two equations above, combined with those of the generatrix in Problem I., there results the following relation between α and β, viz.

$$(A\alpha^2 + B\beta^2 - 1)(Aa + Bb + C) = (Aa\alpha + Bb\beta)^2;$$

in which equation we are to substitute, agreeably to Problem I., the values of α and β, given by the generatrix, in terms of x, y, z. The result of this substitution is

$$\{A(x - az)^2 + B(y - bz)^2 - 1\}(Aa^2 + Bb^2 + C) =$$
$$\{Aa(x - az) + Bb(y - bz)\}^2$$

which is therefore the equation of the cylindric surface sought. In order to reduce it to a more commodious form introduce $Cz - Cz$ within the brackets of the second member, which will then be

$$\{(Aax + Bby + Cz) - (Aa^2 + Bb^2 + C)z\}^2$$

and by actually squaring and transposing, the equation ultimately takes the following form, viz.

$$(Ax^2 + By^2 + Cz^2 - 1)(Aa^2 + Bb^2 + C) = (Aax + Bby + Cz)^2,$$

showing that the cylinder which it represents touches the proposed ellipsoid along the curve situated in the plane (2) above, viz. in

$$Aax + Bby + Cz = 0.$$

PROBLEM III.

(137.) To determine the general equation of conical surfaces.

Let (x', y', z') be the vertex of the conical surface, then, since the generatrix always passes through this point, its equations, in any position, will be

$$\left.\begin{aligned} x - x' &= a(z - z') \\ y - y' &= b(z - z') \end{aligned}\right\} \ldots\ (1),$$

$$\therefore a = \frac{x - x'}{z - z'}, \; b = \frac{y - y'}{z - z'}.$$

Also, let the equations of the directrix be

$$F(x, y, z) = 0, \; f(x, y, z) = 0 \ldots\ (2),$$

then, since for every point in this line, the equations (1) and (2) exist together, we may eliminate the variables x, y, z; the result will be an equation, containing the fixed constants x', y', and the indeterminates a, b; therefore, solving this equation for b, we shall have $b = \phi(a)$. Hence, substituting for a, b, their values in terms of x, y, we have

$$\frac{y - y'}{z - z'} = \phi\left(\frac{x - x'}{z - z'}\right),$$

for the equation of conical surfaces in general, the function ϕ depending entirely on the directrix.

(138.) Eliminating the function ϕ, by differentiating each member of this equation with respect to x and y, and dividing the results as in (58), we have

$$\frac{(y - y')\, p'}{z - z' - (y - y')\, q'} = \frac{z - z' - (x - x')\, p'}{(x - x')\, q'},$$

which reduces to

$$z - z' = p'(x - x') + q'(y - y'),$$

the differential equation of conical surfaces in general.

(139.) This same equation, like that of cylindrical surfaces, may be obtained more readily by the consideration of the tangent plane, which, as it always passes through the vertex (x', y', z') is, in every position, represented by the equation

$$z - z' = p'(x - x') + q'(y - y'),$$

this relation, therefore, must exist between the partial differential coefficients p', q', for every point of the surface, in order that it may be conical.

As in Problem I. if for p', q', we substitute their values derived from the implicit equation of any conical surface, the differential equation becomes

$$(x - x')\frac{du}{dx} + (y - y')\frac{du}{dy} + (z - z')\frac{du}{dz} = 0.$$

PROBLEM IV.

(140.) Given the position of the vertex, to determine the equation of the conical surface that envelopes a given curve surface.

Since the cone envelopes the proposed surface, the curve of contact is common to both, so that the tangent planes to the cone touch also the given surface, according to this curve. The equation, therefore, of the tangent plane

$$z - z' = p'(x - x') + q'(y - y') \ldots . (1),$$

holds equally for any point on the conical surface, and for any point in the curve of contact. Hence, if the values of p', q', be derived from the equation of the given surface, and substituted in (1), this, combined with the equation of the given surface, must represent the curve common to both surfaces, that is, the directrix of the cone. Therefore, the vertex and directrix being known, the equation of the required conical surface becomes determinable.

(141.) If the given curve surface be of the second order, the equation (1) will be also of the second order; but, nevertheless, the combination

of these two equations will be that of a plane, for a surface of the second order may be generally represented by the equation

$$\left.\begin{aligned} &Ax^2 + By^2 + Cz^2 \\ &+ 2Dyz + 2Exz + 2Fxy \\ &+ 2Gx + 2Hy + 2Jz \end{aligned}\right\} = K \quad \ldots \quad (2),$$

which gives

$$\frac{dz}{dx} = -\frac{Ax + Fy + Ez + G}{Ex + Dy + Cz + J}$$

$$\frac{dz}{dy} = -\frac{Fx + By + Dz + H}{Ex + Dy + Cz + J},$$

substituting these values for p' and q', in the equation (1), and subtracting from the result the equation (2), we have

$$\left.\begin{aligned} &(Ax' + Fy' + Ez' + G)x \\ &+ (Fx' + By' + Dz' + H)y \\ &+ (Ex' + Dy' + Cz' + J)z \\ &+ Gx' + Hy' + Jz' + K \end{aligned}\right\} = 0 \quad \ldots \quad (3),$$

which is the equation of a plane; therefore, the conical surface which circumscribes a surface of the second order, must itself be also of the second order.

(142.) The above proof is from *Monge (Application de l'Analyse à la Géométrie)*, but it may be rendered much more concise, by assuming the axes of reference so as to give the general equation of the surface a simpler form. Thus, let the axis of x pass through the centre, if the surface have a centre, or be parallel to its diameters if it have not, and let the other two axes be parallel to the conjugates to this, the form of the equation will then be

$$Az^2 + By^2 + Cx^2 + 2Fx = G \quad \ldots \quad (4),$$

$$\therefore \frac{dz}{dx} = -\frac{F + Cx}{Az}, \quad \frac{dz}{dy} = -\frac{By}{Az}.$$

These values, substituted for p' and q' in (1), convert that equation into

$$z - z' + \frac{(F + Cx)(x - x')}{Az} + \frac{By(y - y')}{Az} = 0,$$

or

$$Az^2 + By^2 + Cx^2 + Fx - Az'z - By'y - Cx'x - Fx' = 0.$$

The difference between this and (4) is

$$Az'z + By'y + Cx'x + Fx + Fx' = G,$$

the equation of a plane.

(143.) Referring again to *Monge's* process, we may remark, that if we accent the constants in the general equation (2), it may be taken as the representative of another surface of the second order, for which the plane of contact with a circumscribing cone, whose summit coincides with that of the former cone, will be represented by the equation

$$\left.\begin{aligned} &(A'x' + F'y' + E'z' + G')\,x \\ &+ (F'x' + B'y' + D'z' + H')\,y \\ &+ (E'x' + D'y' + C'z' + J')\,z \\ &+ G'x' + H'y' + J'z' + K' \end{aligned}\right\} = 0 \quad \ldots \quad (5).$$

Now, although equation (3) be multiplied by an indeterminate constant, p, the result will still represent the same plane, and this plane will obviously be identical to that represented by (5), provided the coefficients of the variables x, y, z, are the same in both equations, that is to say, provided we have the conditions

$$p(Ax' + Fy' + Ez' + G) = A'x' + F'y' + E'z' + G'$$

$$p(Fx' + By' + Dz' + H) = F'x' + B'y' + D'z' + H'$$

$$p(Ex' + Dy' + Cz' + J) = E'x' + D'y' + C'z' + J'$$

$$p(Gx' + Hy' + Jz' + K) = G'x' + H'y' + J'z' + K'.$$

As therefore the four quantities, x', y', z', p, are arbitrary, they may be

determined so that these conditions shall be fulfilled, the four equations being just sufficient to fix the values of these four quantities, and as each of them enters only in the first degree, they will each have but one value. It follows, therefore, that there is a certain point, and only one, from which, as a vertex, if tangent cones be drawn to two given surfaces of the second order, their planes of contact shall coincide. The common vertex will be at the intersection of those diameters to each of which the plane of contact is conjugate; since it has been shown above that the vertex of the tangent cone is always situated on that diameter of the surface to which the plane of contact is conjugate.*

(144.) We may here observe, that as we have not fixed the origin of the axes to any particular point on the diameter which has been taken for the axis of x, nor indeed the diameter itself, we may consider the diameter to be that passing through the vertex (x', y', z') of the cone, and this point to be the origin; in which case x', y', z' will each be 0, and the equation of the plane through the curve of contact will then be simply

$$Fx = G \therefore x = \frac{G}{F},$$

hence, the plane through the curve of contact is conjugate to the diameter through the vertex of the cone. If this vertex be supposed infinitely distant, the same result will belong to the circumscribing cylinder, viz. that the plane of the curve of contact is conjugate to the diameter parallel to the generatrix of the cylinder.

Surfaces of Revolution.

(145.) The surfaces of revolution, considered in the Analytical Geometry, comprise those only in which the revolving curve is always situated in the plane of the fixed axis. We shall here treat of surfaces

---

* For these and other kindred properties, the student is referred to *Mr. Davies's* paper on *Geometry of Three Dimensions*, in *Leybourn's Repository*, vol. 5.

of revolution in general, the revolving curve being any how situated with respect to the axes. Sections of the surface, in the plane of the axis, are called *meridians*.

PROBLEM V.

(146.) To determine the equation of surfaces of revolution in general.

Let the equations of the generating curve be

$$F(x, y, z) = 0,\ f(x, y, z) = 0 \ldots\ (1),$$

and those of the fixed axis

$$\left.\begin{aligned} x &= az + \alpha \\ y &= bz + \beta \end{aligned}\right\} \ldots\ (2),$$

then, since the characteristic property of surfaces of revolution is, that every section perpendicular to the fixed axis is a circle, we shall have first to determine a plane perpendicular to the line (2), and then to express the condition that this plane, combined with the surface, always represents a circle whose centre is on (2). Now, the equation of the required plane is (*Anal. Geom.* p. 235,)

$$z + ax + by = c \ldots\ (3),$$

and the condition is, that it must give the same section as if it were to cut a sphere, whose centre we may fix at pleasure, but whose radius will vary with the section, that is, it will depend upon c in equa. (3). Assuming the centre of this sphere at the point where the line (2) pierces the plane of xy, its equation will be (*Anal. Geom.* p. 241,)

$$(x - \alpha)^2 + (y - \beta)^2 + z^2 = r^2 \ldots\ (4).$$

Hence, supposing r to be the proper function of c, the equations (1), (3), (4), must all have place together; hence we may eliminate x, y, z, and thus determine what the relation between r and c must necessarily be, to render these equations coexistent. The result of the elimination will obviously lead to $c = \phi(r^2)$, hence, substituting for c and r their

values in terms of the variables, we have, finally,

$$z + ax + by = \phi\{(x-\alpha)^2 + (y-\beta)^2 + z^2\} \ldots\ldots (5),$$

for the relation which must always exist among the co-ordinates of every point, in every circular section. This, therefore, is the equation of surfaces of revolution in general.

(147.) If the fixed axis be taken for the axis of z, then a, α; b, β, are each 0, therefore, in this case, the general equation becomes

$$z = \phi(x^2 + y^2 + z^2) \ldots\ldots (6),$$

which, solved for z, gives the form

$$z = \psi(x^2 + y^2) \ldots\ldots (7).$$

(148.) There is one case of this general problem, viz. that where the generatrix is a straight line, revolving round the axis of z, but not in the same plane with it, that deserves particular notice.

Let us take, for axis of x, the shortest distance between the axis of z and the generating line; then this axis will be perpendicular to both (65); the equations, therefore, of the line will be

$$x = \pm a,\ y = \pm bz,$$

also, for any variable section perpendicular to the axis of z,

$$z = c,\ \ x^2 + y^2 + z^2 = r^2.$$

Eliminating x, y, z, we have

$$a^2 + b^2c^2 + c^2 = r^2,$$

for c and r^2 putting their values above, we have

$$a^2 + b^2z^2 = x^2 + y^2.$$

By putting successively $x = 0$, $y = 0$, in this equation, the resulting forms belong to hyperbolas; hence the surface is the *hyperboloid of re-*

volution of a single sheet. The equation of the hyperbola corresponding to $x = 0$ is

$$z = \frac{y}{b}\sqrt{1 + \frac{a^2}{y^2}},$$

so that $y = \pm bz$ is the equation of the asymptotes (see *Anal. Geom.* p. 99,) hence the generating straight line, in its first position, is in a plane with, and parallel to, one or other of the asymptotes of that hyperbola in its first position, which would generate by revolving round the axis of z, the same surface as the line; these two lines, therefore, continue parallel during the revolution of both; the one, viz. the asymptote, generating the conical surface asymptotic to the hyperboloid generated by the other line, viz. the line

$$x = \pm a,\ y = bz,$$

or

$$x = \pm a,\ y = -bz,$$

and it therefore follows that these four lines will be the sections made on the surface by two tangent planes to the asymptotic cone, drawn through any diametrically opposite points in its surface; these will cut each other on the surface, two and two, and include an angle equal to that between the asymptotes; so that the surface may be generated by the revolution of either of these intersecting lines.

We shall shortly see that hyperboloids of one sheet, in general, admit of two distinct modes of generation by the motion of a straight line.

(149.) Eliminating the indeterminate function ϕ, in (5), which depends on the nature of the generating curve (1) by differentiation, as, in the preceding problems, we find

$$\frac{p' + a}{q' + b} = \frac{x - \alpha + p'z}{y - \beta + q'z},$$

from which results the partial differential equation

$$(y - \beta - bz)\,p' - (x - \alpha - az)\,q' - b\,(x - \alpha) - a\,(y - \beta) = 0 \quad . \quad (1),$$

and when the axis of z coincides with that of revolution, this becomes

$$yp' - xq' = 0.$$

The differential equation of surfaces of revolution may also be obtained from the consideration of the normal, which must always cut the axis of revolution, being situated in the meridian plane. Thus, the equations to the normal are (127)

$$\left.\begin{aligned} x - x' + p'(z - z') &= 0 \\ y - y' + q'(z - z') &= 0 \end{aligned}\right\}$$

and as these must exist simultaneously with the equations (2), we may eliminate x, y, z, and the result will necessarily be the required relation between p', q', and the variable co-ordinates x', y', z', of any point on the surface.

PROBLEM VI.

(150.) A given curve surface revolves round a given axis, to determine the surface which touches and envelopes the moveable surface in every position.

The enveloping surface touches the moveable one in every position; if, therefore, we take any particular position of the latter, their combination will give the curve of contact; this curve being common to both surfaces, the tangent planes, at all its points, are common to both surfaces; hence, the values of p', q', which vary only with the tangent plane, are the same for both surfaces, as far as this common curve is concerned, and it is evidently by the revolution of this curve round the fixed axis that the enveloping surface is generated. Hence, to determine this curve, we must deduce p', q', from the given equation, substitute them in the general equation (1) of surfaces of revolution, since there is a line on some such surface to which they belong, as well as to the given surface; and then, to determine what this line really is, it will be necessary merely to combine this last result with the equation of the given surface: we shall thus obtain the equations of the generating curve, and the position of the fixed axis being previously known, the enveloping surface is determinable by Prob. V.

(151.) As an illustration of this, let us suppose a spheroid to revolve about any diameter, to find the equation of the surface enveloping it in every position.

Let the surface be referred to the principal diameters of the spheroid, then the equations of any other diameter will be

$$x = az,\ y = bz \ldots . \ (1),$$

and the spheroid itself may be represented by the equation

$$x^2 + y^2 + n^2z^2 = m^2,$$

from which we derive

$$p' = -\frac{1}{n^2} \cdot \frac{x}{z},\ q' = -\frac{1}{n^2} \cdot \frac{y}{z},$$

substituting these values in the general equation, for all surfaces of revolution round the proposed axis (1), that is in the equation

$$(y - bz)\,p' - (x - az)\,q' + ay - bx = 0,$$

and we have

$$(ay - bx)\,(1 - \frac{1}{n^2}) = 0,$$

$$\therefore\ ay = bx,$$

hence, combining this with the given equation, we have, for the generating curve of the envelope, the equations

$$\left.\begin{aligned} x^2 + y^2 + n^2z^2 &= m^2 \\ ay &= bx \end{aligned}\right\} \ldots . \ (2),$$

hence, the envelope itself is to be determined thus. We must eliminate x, y, z, by means of (2), and the equations

$$\left.\begin{aligned} z + ax + by &= c \\ x^2 + y^2 + z^2 &= r^2 \end{aligned}\right\} \ldots . \ (3),$$

of any circular section, the result will be

$$(r^2n^2 - m^2)(a^2 + b^2) = \{c\sqrt{n^2 - 1} - \sqrt{m^2 - r^2}\}^2,$$

putting for r and c their values in terms of x, y, z, we have, finally,

$$\{n^2(x^2 + y^2 + z^2) - m^2\}(a^2 + b^2),$$

$$= \{(z + ax + by)\sqrt{n^2 - 1} - \sqrt{m^2 - x^2 - y^2 - z^2}\}^2,$$

which is the equation of the enveloping surface.*

CHAPTER III.

ON THE CURVATURE OF SURFACES IN GENERAL.

(152.) The simplest method of contemplating surfaces is by considering them as produced by the motion of a line straight or curved, which, in all its positions, is subject to a fixed law. Viewed under this aspect, surfaces seem to divide themselves into two distinct and very comprehensive classes, viz. those whose generatrices must necessarily be a curve, and those whose generatrices may be a straight line. If, in this latter class of surfaces, the law which regulates the generating straight line be such that through any two of its positions, however close a plane may always be drawn, then it is obvious, that in every such surface, if a plane through the generatrix in any position, but not through any other points of the surface, that is, if a tangent plane, be drawn, this plane, if supposed perfectly flexible, might be wrapped round the surface, without being twisted or torn; or, on the contrary, the surface itself might be unrolled, and would then coincide in all its points with the plane. Surfaces of this kind are, therefore, very properly distin-

* This solution is from *Hymers's Geometry of Three Dimensions*, p. 145.

guished by the name *Developable Surfaces;* the simplest of these are the cone and cylinder.

(153.) We see, therefore, that these surfaces are such that a plane may be drawn through any two positions of the generatrix, and which, if turned round one position, supposed fixed, will pass through all the intermediate positions of the other. But if the law of generation is such that this cannot have place for any two positions, however close, then the tangent plane, through one position, could plainly never be brought to pass also through another position, however near, without being *twisted.* Such surfaces, therefore, are properly designated by the name *Twisted Surfaces.*

These two kinds of surfaces will be separately discussed hereafter; the particulars in the present chapter relate to curve surfaces in general.

Osculation of Curve Surfaces.

(154.) Let the equations of two curve surfaces be

$$z = f(x, y), \quad Z = F(x, y),$$

when referred to the same axes of coordinates. The first of these surfaces we shall suppose fixed, both in magnitude and position, by the constants a, b, c, &c., which enter its equation, being fixed. The second surface we shall suppose fixed only in form, by the form of its equation being given, but indeterminate as to magnitude and position, on account of the arbitrary constants A, B, C, &c. which enter its equation.

Let now the variables x and y take the increments h and k; then, for the first surface, we have (60)

$$z' = z + \frac{dz}{dx}h + \frac{dz}{dy}k + \tfrac{1}{2}\left(\frac{d^2z}{dx^2}h^2 + 2\frac{d^2z}{dx\,dy}hk + \frac{d^2z}{dy^2}k^2\right) + \&c.;$$

and for the second,

$$Z' = Z + \frac{dZ}{dx}h + \frac{dZ}{dy}k + \tfrac{1}{2}\left(\frac{d^2Z}{dx^2} + 2\frac{d^2Z}{dx\,dy}hk + \frac{d^2Z}{dy^2}k^2\right) + \&c.$$

or, more briefly,

$$z' = z + p'h + q'k + \tfrac{1}{2}(r'h^2 + 2s'hk + t'k^2) + \&c.$$

$$Z' = Z + P'h + Q'k + \tfrac{1}{2}(R'h^2 + 2S'hk + T'k^2) + \&c.$$

Now the constants A, B, C, &c. being arbitrary, we may determine one of them in functions of x, y, and the known constants, so that the condition

$$z = Z \quad \text{or} \quad f(x, y) = F(x, y)$$

may be fulfilled. Such a value substituted for the constant in the equation $Z = F(x, y)$, will cause all the surfaces represented by this equation to have a point (x, y, z) in common with the given surface. If two more of the arbitrary constants be determined from the conditions

$$p' = P', \quad q' = Q',$$

the resulting values of these constants being also substituted in the same equation, the surfaces then represented will, in consequence, all have a tangent plane at the point (x, y, z) in common with the fixed surface. Therefore, that this may be the case, three arbitrary constants, at least, must enter the proposed equation, and the contact which they determine is called *contact of the first order*. *Contact of the second order* requires that the following additional conditions be fulfilled, viz.

$$r' = R', \quad s' = S', \quad t' = T',$$

requiring three more arbitrary constants to be determined, and so on; and that surface, all of whose arbitrary constants are determined agreeably to these conditions, will, for reasons similar to those assigned at (88) for plane curves, touch the proposed surface more intimately than any other surface of the same order. It is called the *osculating surface* of that order.

If the touching surface be a sphere, then, since in its equation there

can enter only *four* disposable constants, the contact cannot be so high as the second order, seeing that for this there must be *six* disposable constants; but as contact of the first order would leave still one constant arbitrary, it follows that an infinite number of spheres may have simple contact with a surface at any proposed point, yet one of these *may* be determined that shall be strictly the osculating sphere, or which shall touch more intimately all round the point of contact than any other.

Curvature of different Sections.

PROBLEM I.

(155.) At any point on a curve surface to find the radius of curvature of a normal section.

For greater simplicity, let us suppose the plane of xy to coincide with the tangent plane at the proposed point, then the axis of z will coincide with the normal, and all the normal sections will be vertical. Let the plane of the proposed section be inclined at an angle θ to the plane of xz, then the angle which its trace x' on the plane of xy makes with the axis of x will obviously be θ; and the x, y of this trace will also be the x, y of the section. Now by equation (1) at (98) the length of the radius of curvature ρ at the proposed point where, (87), $(dx') = (ds)$ and $\frac{(dz)}{(dx')} = 0$, is

$$\rho = \frac{\frac{ds^2}{dx^2}}{\frac{d^2z}{dx^2}} = \frac{\frac{ds^2}{dx^2}}{r' + 2s'\frac{dy}{dx} + t'\frac{dy^2}{dx^2}}.$$

But (87)

$$\frac{dx'^2}{dx^2} = \frac{ds^2}{dx^2} = 1 + \frac{dy^2}{dx^2} = 1 + \tan^2\theta;$$

hence, by substitution,

$$\rho = \frac{1 + \tan^2\theta}{r' + 2s'\tan\theta + t'\tan^2\theta}$$

$$= \frac{1}{r'\cos^2\theta + 2s'\cos\theta\sin\theta + t'\sin^2\theta} \quad \ldots\ (1).$$

For the radius of curvature ρ', of a second normal section inclined at an angle $\theta + 90°$ to the plane of xz, we have, by putting $\theta + 90°$ for θ,

$$\rho' = \frac{1}{r' \sin^2\theta - 2s' \cos\theta \sin\theta + t' \cos^2\theta} \quad \ldots\ (2),$$

consequently,

$$\frac{1}{\rho} + \frac{1}{\rho'} = r' + t' \quad \ldots\ (3);$$

so that *the sum of the curvatures of any two normal sections through the same point at right angles to each other, is a constant quantity.*

(156.) Consequently, when one of these curvatures is the *greatest possible*, the other must be the *least possible;* that is, at every point on a curve surface, *the sections of greatest and least curvature are always perpendicular to each other*, which beautiful theorem was first discovered by *Euler*, and is demonstrated by most writers on curve surfaces, though in a manner far less simple than that above.

(157.) To determine the values of the radii of curvature of any perpendicular sections at their point of intersection, let the plane of xz be made to coincide with one of them, by turning round the normal; that is to say, let $\theta = 0$, then the foregoing expressions for ρ and ρ' become

$$\rho = \frac{1}{r'}, \quad \rho' = \frac{1}{t'} \quad \ldots\ (4).$$

(158.) But to determine the expressions for the radii of *greatest* snd *least* curvatures, without causing the vertical planes of coordinates to coincide with the sections, we must know the inclinations of these sections to the vertical planes; that is, we must know the angle θ. To find this from the property $\frac{1}{\rho} = max.$ or $min.$, we have, taking θ for the independent variable in the expression for ρ,

$$\frac{\mathrm{d}\frac{1}{\rho}}{\mathrm{d}\theta} = -2r' \cos\theta \sin\theta + 2s'(\cos^2\theta - \sin^2\theta) + 2t' \sin\theta \cos\theta = 0 \quad \ldots\ (5);$$

hence, dividing by $2\sin^2\theta$, we have

$$\cot^2\theta + \frac{t'-r'}{s'}\cot\theta - 1 = 0 \quad \ldots\ldots (6),$$

or if we divide by $2\cos^2\theta$ instead of by $2\sin^2\theta$, we shall have

$$\tan^2\theta + \frac{r'-t'}{s'}\tan\theta - 1 = 0 \quad \ldots\ldots (7),$$

from either of which equations we may obtain the two inclinations sought. From the first we have

$$\cot\theta = \frac{r'-t' \pm \sqrt{(r'-t')^2 + 4s'^2}}{2s'};$$

from the second,

$$\tan\theta = \frac{t'-r' \pm \sqrt{(r'-t')^2 + 4s'^2}}{2s'},$$

the upper sign corresponding to the maximum, and the under sign to the minimum.

The first expression for ρ in (1) may be written thus:

$$\rho = \frac{\cot^2\theta + 1}{r'\cot^2\theta + 2s'\cot\theta + t'},$$

whence

$$\frac{1}{\rho} = \frac{1}{1+\cot^2\theta}\{r'\cot^2\theta + 2s'\cot\theta + t'\},$$

but from (6),

$$t' = s'\tan\theta - s'\cot\theta + r';$$

hence, by substitution,

$$\frac{1}{\rho} = \frac{1}{1+\cot^2\theta}\{r'(1+\cot^2\theta) + s'(\tan\theta + \cot\theta)\}$$

$$= \frac{1}{1+\cot^2\theta}\{r'(1+\cot^2\theta) + s'\tan\theta(1+\cot^2\theta)\}$$

$$= r' + s' \tan\theta = r + \frac{t' - r' \pm \sqrt{(r' - t')^2 + 4s'^2}}{2}$$

$$= \frac{t' + r' \pm \sqrt{(r' - t')^2 + 4s'^2}}{2}.$$

Consequently, for the radii of greatest and least curvature we have the expressions

$$\left.\begin{aligned} R &= \frac{2}{r' + t' - \sqrt{(r' - t')^2 + 4s'^2}} \\ r &= \frac{2}{r' + t' + \sqrt{(r' - t')^2 + 4s'^2}} \end{aligned}\right\} \quad \ldots . \quad (8).$$

These are called the *principal radii of curvature* at the proposed point, and the sections themselves the *principal sections* through that point.

(159.) If we know the principal radii and the inclination ϕ of any normal section to a principal section through the point, the radius of curvature of the normal section at that point may be expressed in terms of these known quantities. For, bringing the vertical coordinate planes into coincidence with the planes of principal section, we have $\theta = 0$, and, consequently, as appears from equation (6), last article $s' = 0$; and, since (4)

$$\frac{1}{r} = r', \quad \frac{1}{R} = t',$$

we have (1)

$$\rho = \frac{1}{r' \cos^2\phi + t' \sin^2\phi} = \frac{Rr}{r \sin^2\phi + R \cos^2\phi} \quad \ldots . \quad (9),$$

$$\therefore \frac{1}{\rho} = \frac{\sin^2\phi}{R} + \frac{\cos^2\phi}{r} \quad \ldots . \quad (10).$$

It is plain from this expression that if R and r have the same sign, ρ will have that sign for every section through the proposed point, which is the same as saying that if the principal sections are both convex or both concave, every other section through the same point will be similarly convex or concave, and, therefore, also the entire surface at that point. In such a case the minimum radius must be absolutely

shorter than any other radius of curvature at the point, and the maximum radius longer than any other.

(160.) If the two principal radii have not only the same sign, but the same length, then the foregoing expression gives always $\rho = R$ whatever be the inclination ϕ; so that then all the normal sections have the same curvature, and all are *principal sections*, as is the case with the sphere, and with the ellipsoid of revolution, the paraboloid of revolution, &c. at those points through which the fixed axis passes. Such points are called by the French *ombilics;* they evidently exist for every point of the surface at which r', s', t', fulfil the conditions

$$r' - t' = 0, \quad s' = 0,$$

which are necessary and sufficient to make $R = r$; it being remembered that, in all our preceding analytical conditions, the axis of z coincides with the normal at the point considered.

(161.) If the surface belong to the second of the classes mentioned in (152), then no point can be assumed on it through which a straight line may not be drawn; and, as the curvature of this line is 0, it follows that the curvature of the section perpendicular to it must equal the sum of the curvatures of any two perpendicular sections through the same point.

(162.) Let us now suppose that the principal radii R, r have different signs, as r positive and R negative, which will be the case if one of these sections be convex and the other concave, we shall then have

$$\rho = \frac{Rr}{R\cos^2\phi - r\sin^2\phi},$$

which becomes infinite when

$$r\sin^2\phi = R\cos^2\phi, \text{ or when } \tan\phi = \pm\sqrt{\frac{R}{r}},$$

but for all positive and negative values of ϕ between this and 0, ρ will be positive, while beyond these limits ρ will be negative.

It appears, therefore, that if from the origin two straight lines are drawn in the tangent plane, inclined to the axis of x at the angles $\tan\phi = +\sqrt{\frac{R}{r}}$ and $\tan\phi = -\sqrt{\frac{R}{r}}$, these will coincide with the surface; all the sections between the sides of the two opposite angles thus formed will be convex, all the sections between the sides of the other two opposite supplementary angles will be concave; so that the two straight lines which we have seen may be drawn from the proposed point to coincide with the surface, separate the convexity from the concavity at that point.

(163.) In order to determine whether the principal radii at any point are both of the same sign or not, we may observe that the expressions (8) for these radii at art. (158) may be put under the forms

$$\left.\begin{aligned} R &= \frac{2}{r' + t' - \sqrt{(r' + t')^2 - 4(r't' - s'^2)}} \\ r &= \frac{2}{r' + t' + \sqrt{(r' + t')^2 - 4(r't' - s'^2}} \end{aligned}\right\} \quad \ldots\ldots (10),$$

from which forms we immediately see that the radii will have the same sign, viz. positive if $r't' - s'^2 > 0$, and contrary signs if $r't' - s'^2 < 0$; this last condition, therefore, exists in the case just considered.

(164.) We shall terminate these remarks by showing that a paraboloid of the second order may always be found, such that its vertex being applied to any point in any curve surface, the normal sections through that point shall have the same curvature for both surfaces.

For, take the planes of the principal sections for those of xz, yz, then the radii of these sections being R, r we know that a paraboloid, whose vertex is at the origin, will in reference to the same axes be represented by the equation (*Anal. Geom.* p. 263,)

$$z = \frac{x^2}{2r} \pm \frac{y^2}{2R},$$

r and R being the semi-parameters of the sections of the paraboloid on

the planes of xz, yz. Now the equation of a normal section of this paraboloid, by a plane whose inclination to that of xz is ϕ, will be obtained by substituting in this equation $x' \cos \phi$ for x, $x' \sin \phi$ for y, z remaining the same for all normal sections (*Anal. Geom.* p. 281); hence, the equation of the section in question is

$$z = \left(\frac{\cos^2 \phi}{2r} \pm \frac{\sin^2 \phi}{2R}\right) x'^2 \therefore x'^2 = \frac{2Rr}{R \cos^2 \phi \pm r \sin^2 \phi} z.$$

so that the semi-parameter, and, consequently, the radius of curvature (94) of this parabolic or hyperbolic section, is

$$\frac{Rr}{R \cos^2 \phi \pm r \sin^2 \phi},$$

the very same as the radius of curvature of the corresponding section of the proposed surface, be this what it may (159). Hence, this paraboloid has the same curvature in every direction that the proposed surface has at the origin of the coordinates. As a tangent plane at the point of osculation touches both surfaces, and as both have the same curvature there, it follows that if a secant plane, parallel to and at a small distance from the tangent, continually move towards it, the varying section will approach nearer and nearer in form to a curve of the second order, in which it will finally vanish when the secant comes to coincide with the tangent; and the direction of the axes of this evanescent curve will be the same as those of greatest and least curvature through the point. *Dupin*, who first noticed this curious circumstance, calls the curve to which the section tends the *indicatrix* of the surface at that point. A surface may thus be regarded as covered with elliptic, parabolic, and hyperbolic points; the first class including circular points which are the "ombilics" or *umbilici* of the surface.

It is demonstrable that on every surface there exists a curve of double curvature along which the points are all parabolic; and which curve divides the surface into two regions, in one of which the points are all hyperbolic, and in the other all elliptic. Moreover, in the hyperbolic region there exists a curve of double curvature, along which the hyperbolic points are equilateral; and this curve subdivides the hyperbolic region into two, such that the angle of the asymptotes is acute through-

out the one, and obtuse throughout the other. For the investigation of these curious particulars we must refer the student to a paper on the "Courbure des Surfaces," by M. Gergonne, in the "Annales de Mathématiques," 1831.

PROBLEM II.

(165.) To determine the radius of curvature at any point in an oblique section.

Take the tangent to the section through the point as axis of x, the point itself for the origin, and the axis of z' in the plane of the section; then, calling the *normal* the axis of z, the normal section through the axis of x, s, and the oblique section s', we have, at the proposed point (87), $(ds) = (ds')$. Now at the same point $\gamma = \frac{(ds')^2}{(d^2z')}$, (98), but if the axis of z' be transferred to the axis of z, then

$$z = z' \cos\theta \therefore (d^2z) = (d^2z') \cos\theta;$$

hence, by substitution,

$$\gamma = \frac{(ds)^2}{(d^2z)} \cos\theta = \rho \cos\theta \ldots\ldots (1),$$

where γ is the radius of the oblique section, and ρ the radius of the normal section through the tangent to the former; so that γ *is the projection of* ρ *on the plane of the oblique section*, which remarkable property is the *theorem of Meusnier*.

It immediately follows from this theorem, that, if with the radius of any normal section of a curve surface a sphere be described, and through the tangent to that section at the normal point, planes be drawn, cutting both the sphere and the proposed surface, every section of the sphere will be an osculating circle to the corresponding section of the surface; because, if the normal radius of the sphere be projected on any of these sections, the projection will obviously be the radius of that section, and the same projection is, by the above theorem, the radius of curvature of the corresponding section of the proposed surface.

Lines of Curvature and Radii of Spherical Curvature.

(166.) In speaking of plane curves we have already explained (106) what is to be understood by *consecutive normals* and *consecutive curves*. We propose, in the present article, to consider the intersections of any normal at a point of a curve surface, with its consecutive normal; but here it must be remarked that consecutive normals to curve surfaces do not necessarily intersect, as in plane curves; for, before coinciding, these normals, although ever so close, need not be both in the same plane; and, in such a case, when they become consecutive, or coincide, they coincide throughout at once, having even then no point in common that before coinciding was a point of intersection. Hence such consecutive normals have no point of intersection. If, however, upon any curve surface there can be traced a line, such that the normal to the surface at every point of it is intersected by the consecutive normal, that line will have peculiar properties. Such a line is called, by *Monge*, *a line of curvature*.

PROBLEM III.

(167.) To determine the lines of curvature through any point on a curve surface.

Let the surface be referred to any rectangular axes whatever, then $(x', y', z',)$ being any point on it, we have, for the equations of the normal,

$$\left.\begin{aligned} &(A) \qquad x - x' + p'(z - z') = 0 \\ &(B) \qquad y - y' + q'(z - z') = 0 \end{aligned}\right\} \quad \ldots\ldots (1).$$

Let now the independent variables x', y', take any increments h, k; the equations of the normal to the corresponding point will be

$$\left.\begin{aligned} &A + \frac{dA}{dx'}h + \frac{dA}{dy'}k + \&c. = 0 \\ &B + \frac{dB}{dx'}h + \frac{dB}{dy'}k + \&c. = 0 \end{aligned}\right\} \quad \ldots\ldots (2).$$

Now, if the normals (1), (2) intersect, their equations must exist simultaneously; therefore, since A = 0, B = 0,

$$\left.\begin{aligned} \frac{dA}{dx'} + \frac{dA}{dy'}\,\frac{k}{h} + \&c. = 0 \\ \frac{dB}{dx'} + \frac{dB}{dy'}\,\frac{k}{h} + \&c. = 0 \end{aligned}\right\} \quad \ldots\ldots (3)$$

The coordinates $(x, y, z,)$ of the intersection of the proposed normals will be obtained by the combination of the four equations (1) and (3) in terms of x', y', z', which are fixed, and of the increments k, h. But from four equations three unknowns may be always eliminated, and the result of this elimination will be an equation between the other quantities; hence then there exists a *constant* relation between the increments k, h, when the normals intersect; these increments are therefore *dependent;* consequently the y', x', of which these are the increments, must be dependent;* therefore, when the normals are consecutive, that is, when $h = 0$, the equations (3) become

$$\left.\begin{aligned} \frac{dA}{dx'} + \frac{dA}{dy'}\,\frac{dy'}{dx'} = 0 \\ \frac{dB}{dx'} + \frac{dB}{dy'}\,\frac{dy'}{dx'} = 0 \end{aligned}\right\} \quad \ldots\ldots (3'),$$

or, by substituting for A and B their values (1),

$$1 + p'\left(p' + q'\,\frac{dy'}{dx'}\right) + (z' - z)\left(r' + s'\,\frac{dy'}{dx'}\right) = 0 \quad \ldots\ldots (4),$$

$$\frac{dy'}{dx'} + q'\left(p' + q'\,\frac{dy'}{dx'}\right) + (z' - z)\left(s' + t'\,\frac{dy'}{dx'}\right) = 0 \quad \ldots\ldots (5),$$

---

* If this should appear doubtful to the student, its truth may be shown by removing the axes of x, y, to the proposed point, in which position h, k, will be the variable coordinates of the line of curvature, and these will merely take a constant when the axes are replaced in their first position.

from which, eliminating $z' - z$, we have the following equation for determining $\frac{dy'}{dx'}$

$$\{(1+q'^2)s' - p'q't'\}\frac{dy'^2}{dx'^2} + \{(1+q'^2)r' - (1+p'^2)t'\}\frac{dy'}{dx'} -$$

$$(1+p'^2)s' + p'q'r' = 0,$$

or, which is the same thing,

$$\left(\frac{1+q'^2}{t'} - \frac{p'q'}{s'}\right)s't'\frac{dy'^2}{dx'^2} + \left(\frac{1+q'^2}{t'} - \frac{1+p'^2}{r'}\right)r't'\frac{dy'}{dx'}$$

$$-\left(\frac{1+p'^2}{r'} - \frac{p'q'}{s'}\right)r's' = 0 \ldots . \quad (6).$$

This being a quadratic equation furnishes two values for $\frac{dy'}{dx'}$, the tangent of the inclination of the projection of the line of curvature, through (x', y', z'), on the plane of xy, to the axis of x. Hence, there are two directions in which lines of curvature can be drawn through any proposed point; and if in (6) we substitute for p', q', &c. their general values in functions of x, y, that equation will then be the differential equation which belongs to the projections of *every* pair of lines of curvature; so that every line on a curve surface which at all its points satisfies this equation, will be a line of curvature.

In the foregoing equation if we assume

$$\frac{1+p'^2}{r'} = \frac{p'q'}{s'} = \frac{1+q'^2}{t'},$$

the tangent $\frac{dy'}{dx'}$ will not be restricted by that equation to particular values, inasmuch as the equation then becomes identical, and will therefore be satisfied for *any* value of $\frac{dy'}{dx'}$. Hence, from the points at which the foregoing conditions have place, there issue lines of cur-

vature in every direction. These points are "*ombilics*," and the conditions which they must fulfil become the same as those already noticed at (160), when, as is the case there, $p' = 0, q' = 0$.

(168.) Between every pair of lines of curvature there exists a very remarkable relation: it is that they are always *at right angles to each other*. To prove this it will only be necessary to place the coordinate planes, which have hitherto been arbitrary, so that the plane of xy may coincide with, or at least be parallel to, the tangent plane at the point to be considered, in which case p' and q' are both 0; and, consequently, the equation (6) becomes

$$\frac{dy'^2}{dx'^2} + \frac{r' - t'}{s'} \frac{dy'}{dx'} - 1 = 0 \quad \ldots \quad (7),$$

therefore, calling the two roots or values of $\frac{dy'}{dx'}$, $\tan\theta$ and $\tan\theta'$, we have, by the theory of equations,

$$\tan\theta \tan\theta' = -1,$$

which proves that the projections of the two lines of curvature through the origin, are perpendicular to each other, and consequently the lines themselves are perpendicular to each other.

Moreover, the equation (7), if divided by $\frac{dy'^2}{dx'^2} = \tan^2\theta$, becomes identical to equation (6), page 228, which determines the inclinations of the principal sections; hence, the lines of curvature through any point, always touch the sections of greatest and least curvature at that point. Also, in the same hypothesis, with respect to the disposition of the coordinate planes, $z' = 0$, as well as p', q'; therefore the equation (4) or (5) gives

$$z = \frac{1}{r' + s' \tan\theta}, \quad \text{or} = \frac{\tan\theta}{s' + t' \tan\theta},$$

but if the plane of xz coincide with a plane of principal section, it will, as we have just seen, touch the line of curvature, and then $\theta = 0$, so that

$$z = \frac{1}{r'} \text{ or } \frac{1}{t'},$$

and these are precisely the expressions found at (159), for the two radii of curvature of the principal sections at the proposed point, in reference to the same axes; hence we infer, that the consecutive normals to the surface at any point, intersect at the same points as the consecutive normals to the principal sections. These points of intersection are no other than the centres of curvature of the surface at the proposed point, for if spheres be described from these centres to pass through the proposed point, they will touch there, since both have the same normal, and therefore the same tangent plane; and these two spheres have the same curvature as the surface in the two directions of the lines of curvature, since consecutive normals to the surface in these directions, cut that through the point at the centres of these spheres; also the plane sections, tangential to these directions, have the corresponding sections of the spheres for their osculating circles, since the consecutive normals, at their point of contact, also intersect at these centres; therefore, the radii of curvature of the surface at any point, coincides entirely with the radii of curvature of the principal sections through that point, so that (160) if the radii are both equal at any point, the curvature of the surface is uniform all round that point.

(169.) The annexed figure is intended to give an idea of the disposition of the lines of curvature on the surface (S,) drawn through points P,P′, &c. PT, P′T′, &c. are the normals to the surface at those points, and as each is intersected by its consecutive normal, the locus TT′ . . . of these intersections is a curve. The locus too of the normals PT, P′T′, &c. themselves, form a surface, throughout perpendicular to the proposed; this surface, thus generated by the motion of a straight line PT along the curve PP′ . . . and each position intersecting its consecutive position, is obviously a *developable surface*; one of whose edges is the line of curvature PP′ . . . and the other the line of centres TT′ . . . which latter is called the *edge of regression of the developable*



surface. Proceeding, in like manner, along the other line of curvature through P, we have another developable normal surface, whose edge of regression is the locus of the centres of curvature belonging to this second line of curvature. Applying similar considerations to *every* point on the surface (S), we shall thus have an infinite number of developable normal surfaces at right angles to each other, and which will obviously form together two continuous volumes; and the edges of regression will, in like manner, form two continuous surfaces, or *sheets*, being the locus of all the centres of curvature. These surfaces, therefore, bear the same relation to the original surface, as that which in plane curves we have called the evolute bears to the involute.

It would be quite incompatible with the pretensions of the present volume to extend any further our enquiries into the properties of *lines of curvature*. For more detailed information respecting these remarkable lines, the student must study the illustrious author by whom they were first considered, MONGE, in his *Application de l'Analyse à la Géométrie*, a work abounding with the most profound and beautiful speculations on the subject of curve surfaces and curves of double curvature; and which, together with the *Developpements de Géométrie of Dupin*, constitute a complete body of information on a very attractive and important branch of mathematical study, the cultivation of which, however, has been almost entirely neglected hitherto in this country.*

Radii of Spherical Curvature.

(170.) We have already seen that the radii of spherical curvature or simply the radii of curvature at any point of a surface, are identical to the radii of the principal sections through that point; and have given tolerably commodious formulas for the calculation of these radii, when the axes to which the surface is referred originate at the proposed point, the plane of xy being coincident with the tangent plane, and the axis of

* The only English Mathematician, I believe, who has produced public proof of his having given much attention to these enquiries, is *Mr. Davies*, of Bath, whose papers on surfaces, &c. in *Leybourn's Repository*, I have already had occasion to refer to.

z with the normal at that point. We have also seen that when these radii are determined, a paraboloid may also be determined, having its vertex at the proposed point and its curvature in all directions round that point and in its immediate vicinity, the same as the curvature of the surface; so that be the surface ever so complicated, its curvature at any particular point will be correctly presented to us by the vertex of a determinable paraboloid. All this, however, supposes the radii of curvature of the surface at this point to be known; it remains therefore, to show how these radii may be determined whatever be the position of the coordinate axes.

PROBLEM IV.

(171.) Given the coordinates of a point on a curve surface to determine the radii of curvature at that point.

Let (x', y', z') be the point on the surface, and (x, y, z) either of the sought points on the normal corresponding to the centres of curvature; then the radius R from either will be given by the expression

$$R^2 = (x - x')^2 + (y - y')^2 + (z - z')^2.$$

Since at the proposed point the equations (1) and (3), at art. (167), must exist simultaneously with this, we have, by substituting in this the values of $(x - x')$, $(y - y')$, as given by (1),

$$R = (z - z')\sqrt{1 + p'^2 + q'^2}.$$

Now, if from (4), (5), we eliminate the unknown $\frac{dy'}{dx'}$, we shall have

$$(z - z')^2 (r't' - s'^2) + (z - z')\{(1 + q'^2) r' - 2p'q's' + (1 + p'^2) t'\} + (1 + p'^2 + q'^2) = 0;$$

or, putting according to Monge,

$$g = r't' - s'^2$$

$$h = (1 + q'^2) r' - 2p'q's' + (1 + p'^2) t'$$

$$k^2 = 1 + p'^2 + q'^2,$$

the equation for determining $z - z'$ becomes

$$(z-z')^2 + \frac{h}{g}(z-z') + \frac{k^2}{g} = 0 \quad \ldots\ldots (1),$$

and the roots of this substituted in the equation

$$R = (z-z')\,k,$$

give

$$R = \frac{k}{2g}\left(h \pm \sqrt{h^2 - 4gk^2}\right) \quad \ldots\ldots (2)$$

$$= \frac{2k^2}{h \pm \sqrt{h^2 - 4gk^2}} \quad \ldots\ldots (3).$$

(172.) Thus the radii of curvature are determined; and the directions of the lines of curvature, and therefore also of the principal sections, are determined by Problem III.; consequently, the radius of curvature of an oblique section, any how inclined to coordinate planes, any how situated with respect to the surface, may now be determined by help of the formulas (9) and (1) at pages 229 and 233. It appears from (3) that the surface will be convex or concave in the direction of a line of curvature in the immediate vicinity of the point, according as $g > 0$ or $g < 0$. If $g = 0$ the equation (3) shows that one of the radii will be infinite.

When the functions of x', y', z', represented by p', q', r', s', t', are complicated, the expressions just deduced for the radii of curvature will obviously be complicated in the extreme. They are, however, easily manageable when the proposed surface is of the second order, as *Dupin* has shown in his *Developpements* for both classes of these surfaces. We shall here give the solution for surfaces which have not a centre, that is, for paraboloids; the process for the other class, or for central surfaces, being exactly the same, but the reductions rather longer.

PROBLEM V.

(173.) To determine the radii of curvature at any point in a paraboloid.

The general equation of paraboloids being

$$\frac{x^2}{A} + \frac{y^2}{B} = 2z,$$

we have

$$p' = \frac{x}{A},\ q' = \frac{y}{B} \therefore 1 + p'^2 + q'^2 = \frac{x^2}{A^2} + \frac{y^2}{B^2} + 1 = k^2$$

$$r' = \frac{1}{A},\ s' = 0,\ t' = \frac{1}{B} \therefore r't' - s'^2 = \frac{1}{AB} = g$$

$$\therefore h = (1 + \frac{x^2}{A^2})\ \frac{1}{B} + (1 + \frac{y^2}{B^2})\ \frac{1}{A} = \frac{A + B + 2z}{AB}.$$

Hence, generally, whatever be the paraboloid, we have, for the coefficients in equation (1) pa. 241, the values

$$\frac{h}{g} = A + B + 2z,\ \frac{k^2}{g} = AB\ (\frac{x^2}{A^2} + \frac{y^2}{B^2} + 1),$$

and consequently for R we have

$$R = \sqrt{\frac{x^2}{A^2} + \frac{y^2}{B^2} + 1} \times$$

$$\{\frac{A+B+2z}{2} \pm \sqrt{(\frac{A + B + 2z}{2})^2 - AB\ (\frac{x^2}{A^2} + \frac{y^2}{B^2} + 1)}\}.$$

The sum of the two radii are, therefore,

$$R + r = \sqrt{\frac{x^2}{A^2} + \frac{y^2}{B^2} + 1} \times (A + B + 2z),$$

but (128) the first of these factors is the reciprocal of the cosine of the inclination α of the normal at the point $(x, y, z,)$ to the axis of z,

$$\therefore (R + r) \cos \alpha = A + B + 2z,$$

which is the expression for the sum of the projections of the radii of curvature on the axis of z; A, B being the semi-parameters of the sections on the planes of xz, yz (*Anal. Geom.* p. 263). If the point be at the vertex, then $x = 0$, $y = 0$, $z = 0$, and the values of R then become

$$R = \frac{A+B}{2} \pm \sqrt{\left(\frac{A+B}{2}\right)^2 - AB} = \frac{A+B}{2} \pm \frac{A-B}{2}$$

$$\therefore R = A,\ r = B,$$

and these are also the radii of curvature of the two parabolic sections on the planes of xz, yz (95), so that these sections which we have already called the principal sections in the Analytical Geometry, are really the principal sections, or those of greatest and least curvature. A similar process leads to similar inferences for central surfaces of the second order.

CHAPTER IV.

ON TWISTED SURFACES.*

(174.) We have already stated (152) a twisted surface to be one whose generatrix is a straight line moving in such a manner along its directrices that it continually changes the plane of its motion.

The present chapter will be devoted to the consideration of this class of surfaces. Proceeding from the simpler kinds to the more general, we shall first examine the surfaces whose directrices are straight lines as

* This is the class of surfaces called by the French *Surfaces Gauches*, and which, together with the class of developable surfaces, they include under the general name of *Surfaces Réglées*, expressive of their mode of generation by straight lined generatrices. There has recently appeared, in

well as the generatrices; then those having one of its directrices a curve, afterwards those having two curvilinear directrices, and lastly those having three directrices of any kind.

Twisted Surfaces having Rectilinear Directrices only.

PROBLEM I.

(175.) To determine the surfaces generated by a straight line moving parallel to a fixed plane, and along two rectilinear directrices not situated in one plane.

Let the fixed plane, called the directing plane, be taken for that of xy, and the plane parallel to the two directrices for that of xz; then the equations of these directrices will be

$$(1) \ldots \left\{ \begin{matrix} x = az + a \\ y = \beta \end{matrix} \right\} \text{ and } \left\{ \begin{matrix} x = a'z + a' \\ y = \beta' \end{matrix} \right. \ldots (2),$$

and the generatrix, being parallel to the plane of xy, will be represented by the equations

$$z = b, \; y = mx + n \ldots (3).$$

As this line has always a point in common with (1), the four equations (1), (3) exist together, therefore, eliminating x, y, z, we have, among the variable parameters, the relation

$$\beta = m(ab + a) + n \ldots (4),$$

the parameters a, a, β, being fixed by the position of the directrices, but the others being variable.

In like manner, since the line (3) has also always a point in common with (2), the four equations (2), (3) exist together; therefore, eliminating

---

Leybourn's Repository, No. 22, a very masterly enquiry into the history of these surfaces, from the pen of *Mr. Davies*, wherein the claims of the English to the first consideration of "rule surfaces" is fully established.

x, y, z, we get for a second relation among the three arbitrary parameters the equation

$$\beta' = m(a'b + \alpha') + n \quad \ldots \quad (5).$$

By means of the two relations (4) and (5) among the parameters which enter (3), we may eliminate them and thus obtain the sought equation in x, y, z. Subtracting each from (3), we have

$$y - \beta = m(x - az - \alpha)$$

$$y - \beta' = m(x - a'z - \alpha'),$$

eliminating m we obtain, finally,

$$(a - a')yz + (\alpha - \alpha')y + (a'\beta - a\beta')z + (\beta' - \beta)x = a\beta' - a'\beta \ldots\ldots (6)$$

for the equation of the surface, which is therefore of the second order. Let us now enquire what particular kind of surfaces of the second order this equation includes. By applying the criteria (3), p. 292, *Anal. Geom.*, we find that the surfaces are not central, they must, therefore, be paraboloids. By putting $x = k$ we find in the resulting equation for any section parallel to the plane of yz, that the squares of the variables are absent, therefore, *Anal. Geom.*, p. 166, these sections are all hyperbolas. We infer, therefore, that the surface (6) is always a *hyperbolic paraboloid.* If in the equation (6) we make z equal to any constant quantity, the equation will always be that of a straight line, being indeed necessarily one of the positions of the generatrix; also, if we put y equal to any constant quantity, we find that every section parallel to the plane of xz is a straight line, so that through every point on the surface of a hyperbolic paraboloid there may be drawn two straight lines, their assemblage constituting two distinct series situated in two distinct series of parallel planes, and hence there are two distinct ways in which the surface may be generated by the motion of a straight line, but not more than two ways, since the equation (6) represents a straight line only on the two hypotheses assumed above: and as no two of the positions of the same generatrix, however close, can be in the same plane, the hyperbolic paraboloid is a *twisted surface.*

(176.) We may show at once, by setting out with the equation of the hyperbolic paraboloid, that two straight lines pass through every

point on its surface, and, moreover, that these lines are both in the tangent plane at that point. Thus the equation of the surface is (*Anal. Geom.* p. 266,)

$$px^2 - p'y^2 = pp'z \quad \ldots\ldots (1),$$

and that of the tangent plane through $(x', y', z',)$

$$2pxx' - 2p'yy' = pp'(z + z') \quad \ldots\ldots (2),$$

the relation among the coordinates x', y', z', of the point of contact being of course

$$px'^2 - p'y'^2 = pp'z' \quad \ldots\ldots (3).$$

Adding together equations (1) and (3), and subtracting (2) from the sum, there results

$$p(x - x')^2 - p'(y - y')^2 = 0,$$

which is the condition necessary to be satisfied for every projected point $(x, y,)$ common to the surface (1) and the plane (2), seeing that it has resulted from the combination of their equations. Such condition being satisfied by every point in the lines represented by the equation

$$y - y' = \pm (x - x') \sqrt{\frac{p}{p'}},$$

it follows that the lines of which these are the projections are common to both surface and tangent plane, so that the tangent plane cuts the surface according to two straight lines passing through the point of contact.

PROBLEM II.

(177.) To determine the surface generated by the motion of a straight line along three others fixed in position, so that no two of them are in the same plane.

Let us first consider the case in which the three directrices are all parallel to the same plane.

Assume the axes of x and y in the plane passing through one of the

directrices (B), and parallel to the other two (B′), (B″). Let the axis of x coincide with (B), and the axis of y be parallel to (B′), and let the axis of z be drawn to pass through both (B′) and (B″); then the equations of the directrices will be

$$\text{(B)} \qquad y=0,\ z=0$$

$$\text{(B′)} \qquad x=0,\ z=h$$

$$\text{(B″)} \qquad y=ax,\ z=k,$$

and the equation of the generatrix in any position will be

$$x=mz+p,\ y=nz+q \ \ldots\ (1).$$

As this line has always a point in common with the directrices, all these equations exist together. Hence, eliminating x, y, z, we have, among the variable parameters m, n, p, q, the relations

$$q=0,\ mh+p=0,\ nk=a\,(mk+p) \ \ldots\ (2).$$

Eliminating the variable parameters from (1) and (2), we have

$$a\,(k-h)\,xz=ky\,(z-h)$$

for the equation of the surface sought, and which we find, by applying the same tests as in last problem, to be the same surface, viz. the hyperbolic paraboloid.

(178.) Suppose, now, that the three directrices are not all parallel to the same plane, then, taking any point in space for the origin, and parallels to the directrices for axes, the equations of these will be

$$\text{(B)} \qquad x=\alpha,\ y=\beta$$

$$\text{(B′)} \qquad z=\gamma,\ x=\alpha'$$

$$\text{(B″)} \qquad y=\beta',\ z=\gamma',$$

and the equation of the generatrix will be

$$x=mz+p,\ y=nz+q \ \ldots\ (1),$$

which, since it has a point in common with (B), gives rise to the condition

$$\frac{\alpha-p}{m}=\frac{\beta-q}{n} \quad \ldots\ldots (2),$$

and having, at the same time, a point in common with (B′), and another in common with (B″), we have the additional conditions

$$\alpha'=m\gamma+p,\ \beta'=n\gamma'+q \quad \ldots\ldots (3).$$

Eliminating now the arbitrary parameters m, n, p, q, by means of (1), and these equations of condition, we shall arrive at the equation of the surface. The equations (3) give, in conjunction with (1),

$$m=\frac{x-\alpha'}{z-\gamma},\quad n=\frac{y-\beta'}{z-\gamma'}$$

$$p=\alpha'-\gamma\,\frac{x-\alpha'}{x-\gamma},\quad q=\beta'-\gamma'\,\frac{y-\beta'}{z-\gamma'},$$

which values, substituted in (2), give

$$\left.\begin{array}{c}(\gamma-\gamma')\,xy+(\beta'-\beta)\,xz+(\alpha-\alpha')\,yz\\ +(\beta\gamma'-\beta'\gamma)\,x+(\alpha'\gamma'-\alpha\gamma)\,y+(\alpha'\beta-\beta\alpha')\,z\\ +\alpha\beta'\gamma-\alpha'\beta\gamma'\end{array}\right\}=0$$

for the equation of the surface. By applying the usual criteria, (*Anal. Geom.* pages 292, 294,) we find that the surface must be a hyperboloid, and as the squares of the variables are all absent from the equation, no intersection (*Anal. Geom.* p. 281) can possibly be an imaginary curve; hence the surface must be a *hyperboloid of a single sheet*, and it is obviously twisted, since the generatrix constantly changes the plane of its motion.

(179.) We may, as in the preceding problem, by commencing with the equation of this surface, show that through every point on it two straight lines may be drawn, and that they will both be in the tangent plane through the point. Thus the equation of the surface is

$$\frac{x^2}{a^2}+\frac{y^2}{b^2}-\frac{z^2}{c^2}=1 \quad \ldots\ldots (1),$$

and that of the tangent plane through (x', y', z')

$$\frac{xx'}{a^2} + \frac{yy'}{b^2} - \frac{zz'}{c^2} = 1 \ldots\ldots (2),$$

the relation among x', y', z' being fixed by the equation

$$\frac{x'^2}{a^2} + \frac{y'^2}{b^2} - \frac{z'^2}{c^2} = 1 \ldots\ldots (3).$$

Adding together equations (1) and (3), and subtracting twice equation (2) from the result, we have

$$\frac{(x-x')^2}{a^2} + \frac{(y-y')^2}{b^2} - \frac{(z-z')^2}{c^2} = 0 \ldots\ldots (4),$$

a relation which must have place for every point common to both the surface and the tangent plane.

Also, subtracting (3) from (2),

$$\frac{x'(x-x')}{a^2} + \frac{y'(y-y')}{b^2} - \frac{z'(z-z')}{c^2} = 0 \ldots\ldots (5).$$

Now, in order to ascertain whether the points fulfilling these conditions can lie in a straight line, let us combine them with the equations of a straight line through (x', y', z'), viz.

$$x - x' = a'(z - z'), \quad y - y' = b'(z - z') \ldots\ldots (6).$$

Substituting in the equations (4) and (5) these expressions for $x - x'$, $y - y'$, we have

$$(z-z')^2 \left(\frac{a'^2}{a^2} + \frac{b'^2}{b^2} - \frac{1}{c^2}\right) = 0,$$

$$(z-z') \left(\frac{a'x'}{a^2} + \frac{b'y'}{b^2} - \frac{z'}{c^2}\right) = 0,$$

$$\therefore \frac{a'^2}{a^2} + \frac{b'^2}{b^2} - \frac{1}{c^2} = 0,$$

$$\frac{a'x'}{a^2} + \frac{b'y'}{b^2} - \frac{z'}{c^2} = 0;$$

these relations, therefore, must exist among the constants in (6), for it to be possible for that line to belong to the surface. From the second of these we readily deduce a rational value of a', which, substituted in the first, b' will be given by the solution of the quadratic, which will furnish two values, so that two lines passing through the point of contact may be drawn, that shall be common to both the surface and the tangent plane.

Twisted Surfaces having but one Curvilinear Directrix.

(180.) In surfaces of this kind the generatrix moves along a straight line and a curve, remaining constantly parallel to a fixed plane, called the *directing plane.* Such surfaces are called *conoids;* and that they are twisted surfaces is plain, because a plane to pass through two positions of the generatrix must pass through the rectilinear directrix, and become, therefore, fixed, so that it cannot be moved round one position without ceasing to pass through two. The directing plane is usually taken for that of xy, the origin being at the point where the straight directrix pierces it.

PROBLEM III.

(181.) To determine the general equation of conoidal surfaces:

Let the equations of the straight directrix be

$$x = mz^2, \quad y = nz \quad . \quad . \quad . \quad . \quad (1),$$

and those of the curvilinear directrix,

$$F(x, y, z) = 0, \quad f(x, y, z) = 0 \quad . \quad . \quad . \quad . \quad (2).$$

The equation of the generatrix, being in every position parallel to the plane of xy, must always be of the form

$$z = \alpha, \quad y = \beta x + \gamma \quad . \quad . \quad . \quad . \quad (3),$$

α and β being variable parameters.

As this line has always a point in common with the line (1), their

equations exist together; hence, eliminating x, y, z, by means of these four equations, we have the condition

$$n\alpha = \beta m\alpha + \gamma \text{ or } \gamma = n\alpha - \beta m\alpha;$$

so that the equations (3) of the generatrix become

$$z = \alpha, \quad y - n\alpha = \beta(x - m\alpha) \ldots . (4);$$

but this same line has also a point in common with the curve (2); hence, eliminating x, y, z, by means of the four equations (2), (4), we have an equation containing only constants and the variable parameters α, β, which equation, solved for α, gives

$$\alpha = \phi(\beta) \ldots . (5).$$

But, by equations (4),

$$\alpha = z, \quad \beta = \frac{y - nz}{x - mz};$$

hence, by substitution in (5),

$$z = \phi\left(\frac{y - nz}{x - mz}\right),$$

which expresses the general relation among the coordinates of any point of the generatrix in any position; therefore this is the general equation of a conoidal surface.

(182.) If the straight directrix coincide with the axis of z, then $m = 0$, $n = 0$, and the conoid is represented by the general equation

$$z = \phi\left(\frac{y}{x}\right),$$

whether the axis of z, or the straight directrix, be perpendicular to the directing plane or not; if it is perpendicular, the conoid is called a *right conoid.* In these cases the equations of the generatrix are simply

$$z = \alpha, \quad y = \beta x.$$

(183.) As an example, let it be required to find the equation of the

inferior surface of a winding staircase, the aperture or column round which it winds being cylindrical.

To conceive the generation of this surface, let us suppose a rectangle to be rolled round a vertical column, which it just embraces, the line which was the diagonal of the rectangle will then become a winding curve called a *helix*, and it will make just one turn round the column, its horizontal projection being a circle; if immediately above this another equal rectangle be applied to the column, the vertical edges when brought together being in a line with those of the first, the diagonal of this will form a continuation of the helix, and in this way will be exhibited the trace of the edge of the surface in question on the vertical column, or the curvilinear directrix; the other directrix is the axis of the cylinder, the directing plane being horizontal.

Now for every point in the diagonal of a rectangle the abscissa has a constant ratio to the ordinate, the axes being the sides including the diagonal; so that, reckoning from the foot of the helix, the circular abscissas and vertical ordinates corresponding are in a constant ratio. Hence, taking the centre of the cylindrical base for the origin, and drawing the axis of y through the foot of the helix, calling h the height and $2\pi r$ the base of one of the rectangles, or of the cylinder, we shall have, for each point of the helix, these relations, viz.

$$x^2 + y^2 = r^2, \quad x = r \sin \frac{s}{r}, \quad \frac{z}{s} = \frac{h}{2\pi r} \ldots . \ (1),$$

and for the generating line, the equations

$$z = \alpha, \quad y = \beta x \ldots . \ (2).$$

If from the two last of (1) we eliminate the arc s we shall have the following equations of the projections of the curve

$$x^2 + y^2 = r^2, \ x = r \sin \left(\frac{2\pi z}{h}\right) \ldots . \ (3),$$

eliminating x, y, z, from the equations (2), (3) we have

$$\frac{1}{\sqrt{1+\beta^2}} = \sin \left(2\pi \frac{\alpha}{h}\right),$$

in which equation if we substitute for α and β the values z and $\frac{y}{x}$ given by (2), we shall obtain, finally,

$$\frac{x}{\sqrt{x^2+y^2}} = \sin\left(2\pi\frac{z}{h}\right), \text{ or } \frac{x}{y} = \tan\left(2\pi\frac{z}{h}\right),$$

which is the equation of the surface, that is, of the *twisted helixoid.*

(184.) It remains to determine the differential equation of conoidal surfaces. In order to this, we must eliminate the arbitrary function ϕ in the equation

$$z = \phi\left(\frac{y-nz}{x-mz}\right);$$

by differentiation, as in the several similar cases in Chapter II., we thus obtain the equation

$$\frac{p'}{q'} = \frac{p'(my-nx)-(y-nz)}{q'(my-nx)+(x-mz)},$$

which reduces to

$$p'(x-mz)+q'(y-nz)=0,$$

or, when the conoid is right, simply to

$$p'x+q'y=0,$$

because then $m=0$, $n=0$.

(185.) The same results may be at once obtained from the consideration of the tangent plane; for (x', y', z') being any point on the surface, the equation of the tangent plane is

$$z-z' = p'(x-x')+q'(y-y'),$$

which touches the surface along the generatrix through (x', y', z'), and this being every where at the same distance z' from the horizontal plane, it follows that if in the above equation we put $z=z'$, the result

$$p'(x-x')+q'(y-y')=0$$

will express the relation between the x, y of *every* point in this generatrix. But at that point where it cuts the straight directrix, the x, y have the relation

$$x=mz',\quad y=nz';$$

so that, by substitution, we have

$$p'(mz'-x')+q'(nz'-y')=0,$$

for the relation among the coordinates of every point (x', y', z') on the surface, which agrees with that deduced above.

Twisted Surfaces having Curvilinear Directrices only.

(186.) We now proceed to consider those surfaces which cannot have a rectilinear directrix, or rather those whose directrices may be any lines whatever. We shall first suppose two directrices.

PROBLEM IV.

(187.) To determine the general equation of surfaces generated by a straight line which moves along any two directrices (D) (D′) whatever, and continues at the same time parallel to a fixed plane.

Taking, as before, the directing plane for that of xy, the equation of the generatrix in any position will be

$$z=\alpha,\quad y=\beta x+\gamma \ldots . (1),$$

the parameters all varying with the varying positions of the generatrix. Let now the equations of the two fixed directrices be

(D) $\quad F(x, y, z)=0,\quad f(x, y, z)=0 \ldots . (2),$

(D′) $\quad F_1(x, y, z)=0,\quad f_1(x, y, z)=0 \ldots . (3).$

Then the condition is, first that the generatrix meets (D), or that their equations (1), (2) exist together; hence, by eliminating the coordinates of the common point from these four equations, we shall obviously obtain an equation containing only constants and the variable parameters α, β, γ; that is to say, we shall obtain among these parameters a relation

$$\Phi(\alpha, \beta, \gamma,) = 0.$$

Proceeding in the same manner with the equations (1), (3) which also exist together for a certain point, we obtain a second relation

$$\Psi(\alpha, \beta, \gamma,) = 0.$$

By means of these two equations we may eliminate any one of the parameters; therefore, eliminating first γ and then β, we have

$$\beta = \phi(\alpha), \quad \gamma = \psi(\alpha);$$

hence, substituting for these variable parameters their values in functions of the variable coordinates as furnished by equation (1), we have, for the general relation among these coordinates, the equation

$$y = x\phi(z) + \psi(z) \ldots . \quad (4).$$

This then is the general equation of all surfaces generated as announced, whatever be the form of the directrices; when these forms are given, the forms of ϕ and ψ become determinable by the above process, and then the general equation (4) takes the particular for' belonging to the individual surface.

(188.) Let us now determine the general equation of these surfaces in terms of the partial differential coefficients. Putting the equation (4) in the form

$$y - x\phi(z) = \psi(z),$$

the ratio of the partial coefficients of each side, taken relatively to x and y, will, by the principle in (58), be

$$\frac{-\phi(z)}{1} = \frac{p'}{q'}; \text{ that is, } \frac{p'}{q'} = -\phi(z),$$

an equation from which the arbitrary function $\psi(z)$ is eliminated. Applying the same principle to this last equation, we have

$$\frac{\mathrm{d}\left(\frac{p'}{q'}\right)}{\mathrm{d}x} \div \frac{\mathrm{d}\left(\frac{p'}{q'}\right)}{\mathrm{d}y} = \frac{p'}{q'};$$

that is, putting according to the usual notation

$$\frac{\mathrm{d}p'}{\mathrm{d}x} = r', \quad \frac{\mathrm{d}q'}{\mathrm{d}x} = \frac{\mathrm{d}p'}{\mathrm{d}y} = s', \quad \frac{\mathrm{d}q'}{\mathrm{d}y} = t',$$

$$\frac{q'r' - p's'}{q'^2} \div \frac{q's' - p't'}{q'^2} = \frac{p'}{q'},$$

whence

$$q'^2 r' - 2p'q's' + p'^2 t' = 0,$$

an equation from which both the arbitrary functions are eliminated, and which must be fulfilled for every point in every surface generated as in the problem, whatever be the directrices. We see that as two arbitrary functions were to be eliminated, the process has led to a partial differential equation of the second order.

PROBLEM V.

(189.) To determine the general equation of surfaces generated by the motion of a straight line along three curvilinear directrices (D), (D′), (D″).

We shall first remark that the motion of the generatrix is entirely governed by these conditions, for if we take any point on the first directrix (D) and conceive two cones whose bases are (D′), (D″) to have this point for their common vertex, these cones will obviously intersect each other in all the straight lines that can be drawn from the point to the curves (D′), (D″), the positions of these lines are therefore fixed by these intersecting cones, and these are fixed by their bases; hence, all the lines that can be drawn from the point to the lines (D′), (D″) are

determinate both in number and position, this being true for every point in (D), it follows that the surface generated by all these lines is determinate, and it is now required to find its equation.

As there is here no directing plane, the equation of the generatrix in any position will take the form

$$x = \alpha z + \gamma, \quad y = \beta z + \delta \ldots . (1);$$

and, since it always has a point in common with (D), we may eliminate by means of the equation of (D) combined with these, the coordinates of that point: the result will furnish a condition among the variable parameters. In like manner, employing the equation of (D'), we shall arrive at another equation of condition, and, lastly, the equation of (D'') will furnish a third equation. By means of these three equations any two of the parameters α, β, γ, δ, may be eliminated, and we shall obtain three equations of the form

$$\beta = \phi(\alpha), \quad \gamma = \psi(\alpha), \quad \delta = \pi(\alpha).$$

Substituting these expressions for β, γ, δ, in the equations (1), we shall have

$$x = \alpha z + \psi(\alpha), \quad y = z\phi(\alpha) + \pi(\alpha),$$

two equations which have place for every surface generated as proposed, the functions which fix the directrices being quite arbitrary. If these functions are known, or the directrices fixed, we may then eliminate the parameter α by means of these equations, and thus deduce the equation of the individual surface; but the general relations among the coordinates for all the surfaces of this family can be exhibited only by means of two equations as above. The general relation among the partial differential coefficients belonging to all this family of surfaces, may, however, be ascertained in a single equation by eliminating, as in last problem, all the arbitrary functions by successive differentiation; this will lead to a partial differential equation of the third order, for which see *Monge's Application de l'Analyse à la Géométrie*, p 195.

(190.) We shall terminate the present chapter with the following example:

On the opposite sides of the horizontal parallelogram ABDC are described two vertical semicircles, and perpendicular to their planes is drawn the straight line OY through the centre of the parallelogram; taking this straight line and the two semicircles as directrices, it is required to find the equation of the surface generated by a straight line moving along them.



Let the axes of coordinates be the perpendicular horizontal lines OX, OY, and the vertical OZ, then the equations of the three directrices will be

$$x = 0, \quad z = 0 \ldots . \ (1),$$

$$y = -b, \quad (x-a)^2 + z^2 = r^2 \ldots . \ (2),$$

$$y = +b, \quad (x+a)^2 + z^2 = r^2 \ldots . \ (3).$$

The equations of the generatrix, since it always passes through a point $(\beta, 0, 0)$ in the axis of y, will take the forms

$$x = \alpha(y-\beta), \quad z = \gamma(y-\beta) \ldots . \ (4),$$

and the condition to be fulfilled by this line is, that it rests on each of the semicircles; or that at certain points, x, y, z, are the same in the equations (2), (4) and (3), (4); hence, eliminating these first from (2), (4), and then from (3), (4), we have these relations among the variable parameters, viz.

$$\{\alpha(b+\beta)+a\}^2 + \gamma^2(b+\beta)^2 = r^2 \ldots . \ (5),$$

$$\{\alpha(b-\beta)+a\}^2 + \gamma^2(b-\beta)^2 = r^2 \ldots . \ (6),$$

which, by subtraction, give

$$\beta(b\alpha^2 + a\alpha + b\gamma^2) = 0.$$

This condition is satisfied by the value $\beta = 0$; but this is not admissible, since it would restrict the generatrix to pass always through the origin,

and have no motion along OY; hence, dividing by $\frac{\beta}{b}$, we have the relation

$$\alpha^2 + \gamma^2 + \frac{a\alpha}{b} = 0 \ldots\ldots (7),$$

between the parameters α, γ.

Substituting the value of γ^2, given by this equation in (5), it becomes

$$(b^2 - \beta^2)\, a\alpha = (r^2 - a^2) \ldots\ldots (8),$$

and by means of these equations, together with those of the generatrix, we may readily eliminate the parameters; thus the values of α and γ, given by (4), are

$$\alpha = \frac{x}{y - \beta}, \quad \gamma = \frac{z}{y - \beta},$$

and these, substituted in (7), give

$$\beta = y + \frac{b(x^2 + z^2)}{ax} \quad \therefore \alpha = -\frac{ax^2}{b(x^2 + z^2)},$$

and finally, these substituted in (8), give for the surface the equation

$$\{y + \frac{b(x^2 + z^2)}{ax}\}^2 - b^2 = b^2 (r^2 - a^2)\, (\frac{x^2 + z^2}{a^2 x^2}),$$

which is the same as

$$\{axy + b(x^2 + z^2)\}^2 = b^2 r^2 x^2 + b^2 (r^2 - a^2)\, z^2.$$

CHAPTER V.

ON DEVELOPABLE SURFACES AND ENVELOPES.

(191.) When in an equation between three variables

$$u = \mathrm{F}(x, y, z, a) = 0$$

there enters an arbitrary constant a, that equation, by giving different values to a, will represent so many different surfaces all belonging to the same family. If we fix one of these by any determinate value of a, another, intersecting this, will be represented by changing a into $a + h$, h being some finite value. If h be now continually diminished, the intersection will continually vary, and will become fixed only when the varying surface becomes coincident with the fixed surface. In this position the intersection is said to belong to consecutive surfaces, and it may be determined both in form and position by a process similar to that employed at (106). Thus a being the only variable concerned in the intersections, we have, by Taylor's theorem,

$$\mathrm{F}(x, y, z, a + h) = u + \frac{\mathrm{d}u}{\mathrm{d}a} h + \frac{\mathrm{d}^2 u}{\mathrm{d}a^2} \frac{h^2}{2} + \&\mathrm{c.} = 0,$$

but, since $u = 0$, therefore,

$$\frac{\mathrm{d}u}{\mathrm{d}a} + \frac{\mathrm{d}^2 u}{\mathrm{d}a^2} \frac{h}{2} + \&\mathrm{c.} = 0;$$

hence, when the surfaces are consecutive, that is, when $h = 0$, we have the following equations for determining the curve of intersection, viz.

$$\left.\begin{aligned} \mathrm{F}(x, y, z, a) &= 0 \\ \frac{\mathrm{dF}(x, y, z, a)}{\mathrm{d}a} &= 0 \end{aligned}\right\} \quad \ldots\ (1),$$

these, therefore, are the equations of the curve, which is the intersection of the surface (1) with its consecutive surface.

If from these two equations we eliminate α, the result will be the general relation among the coordinates of every point in every such consecutive intersection throughout the whole family of surfaces; this resulting equation will therefore represent the surface which is the locus of all these consecutive intersections. This locus, moreover, touches each of the variable surfaces throughout their intersections; for differentiating the equation $u = 0$ of any one of the variable surfaces, α being constant, we have

$$\frac{du}{dx} + \frac{du}{dz}p' = 0, \quad \frac{du}{dy} + \frac{du}{dz}q' = 0,$$

and, differentiating the equation $u = 0$ of the locus, α being variable, we have

$$\frac{du}{dx} + \frac{du}{dz}p' + \frac{du}{d\alpha}\frac{d\alpha}{dx} = 0, \quad \frac{du}{dy} + \frac{du}{dz}q' + \frac{du}{d\alpha}\frac{d\alpha}{dy} = 0,$$

which equations are identical to those above, since (equa. 1,)

$$\frac{du}{d\alpha} = 0,$$

and therefore each pair gives the same values for p' and q', consequently, at the points common to both surfaces they have common tangent planes. Hence the locus of the consecutive intersections touches, and envelopes all the variable surfaces; it is therefore called, by Monge, the *Envelope* of these surfaces.

(192.) If the envelope be formed by the consecutive intersections of *planes*, then, since from what has been just proved, the envelope is touched throughout each of the intersections by the corresponding plane; this envelope is such that the tangent plane at any point touches it throughout the rectilinear generatrix passing through that point; and this is the characteristic property of a *developable surface:* hence a developable surface may be considered as the envelope of a family of planes represented by the general equation

$$z = Ax + By + D \ldots . (1),$$

in which there enters a variable parameter α.

Now, to introduce this variable parameter in the most general manner possible into the equation (1), we ought to consider each of the coefficients A, B, C, to be functions of it, so that the general form will be

$$z = f(\alpha) + x\phi(\alpha) + y\psi(\alpha),$$

and therefore the line of contact (191) or generatrix of the surface will be represented by the equations

$$\left.\begin{aligned} z &= f(\alpha) + x\phi(\alpha) + y\psi(\alpha) \\ 0 &= f'(\alpha) + x\phi'(\alpha) + y\psi'(\alpha) \end{aligned}\right\} \ldots . (2).$$

When the forms of the functions f, ϕ, ψ, are fixed, the variable parameter α may be eliminated, and the resulting equation in x, y, z, will be that of the individual surface to which these particular forms belong. The equations (2), therefore, may be considered as representing the whole family of these surfaces, α in the first being a function of x and y, implied in the second.

(193.) 1. As an example, suppose it were required to determine the developable surface generated by the intersection of normal planes, at every point in a curve of double curvature.

Representing the proposed curve by the equations

$$y' = F(x'), \quad z' = f(x') \ldots . (1),$$

the general equation of the normal plane will be (131)

$$x - x' + \frac{dy'}{dx'}(y - y') + \frac{dz'}{dx'}(z - z') = 0 \ldots . (2),$$

in which the only variable parameter is x'; y' and z' being determinate functions of it given by the equations of the curve. Hence, differentiating with respect to x', we have

$$-\left(1 + \frac{dy'^2}{dx'^2} + \frac{dz'^2}{dx'^2}\right) + \frac{d^2y'}{dx'^2}(y - y') + \frac{d^2z'}{dx'^2}(z - z') = 0 \ldots (3).$$

Now the functions of x', which enter the equations (2), (3), being given by (1), we may eliminate this parameter from them, and the resulting equation in x, y, z, will be that of the developable surface required.

(194.) 2. As a second example, let it be required to determine the developable surface which touches and embraces two given curve surfaces.

If we suppose one of these surfaces to be a luminous surface enlightening the other, the surface which we seek will obviously embrace all the rays which proceed from the bright surface to the dark one, and the curve of contact on this latter will separate the illuminated and dark parts.

Let the equations of the two given surfaces be

$$F_1(x_1, y_1, z_1) = 0 \quad \ldots \quad (1),$$

$$F_2(x_2, y_2, z_2) = 0 \quad \ldots \quad (2),$$

then the equations to tangent planes to each will have the forms

$$z - z_1 = p_1 (x - x_1) + q_1 (y - y_1) \quad \ldots \quad (3)$$

and

$$z - z_2 = p_2 (x - x_2) + q_2 (y - y_2) \quad \ldots \quad (4),$$

and for these planes to belong to *both* surfaces, their equations must be identical; that is, we must have the conditions

$$p_1 = p_2, \quad q_1 = q_2 \quad \ldots \quad (5),$$

$$z_1 - p_1 x_1 - q_1 y_2 = z_2 - p_2 x_2 - q_2 y_2 \quad \ldots \quad (6).$$

By means of the six equations marked, five of the coordinates may be determined in terms of the sixth, x; hence, if these functions of x, be now substituted for their values in the remaining equation, we shall obtain a result containing only the variable parameter x_1, and which will consequently represent the family of planes which generate the developable surface sought. Calling this result $P = 0$, the generatrix of the surface will be given by the equations

$$P = 0, \quad \frac{dP}{dx_1} = 0,$$

from which, eliminating x_1, we have the equation of the surface sought; and this equation, combined with that of each surface separately, will give the two curves of contact.

PROBLEM.

(195.) To determine the differential equation of developable surfaces in general.

The general equation of the generating plane, arranged according to the variable coordinates x, y, z, of any point in it, is

$$z = p'x + q'y + z' - p'x' - q'y' \ldots\ldots (1),$$

and this plane remains the same for every point in the generatrix, as well as for the point (x', y', z'), so that the quantities

$$p', q', z', -p'x' - q'y' \ldots\ldots (2),$$

remain constant, although x', y', z' all vary, provided this variation is confined to the rectilinear generatrix, for which y is always a function of x, but not else; hence, the conditions which restrict the point x', y', z' to the generatrix on which it is first assumed, is, that the differential coefficients derived from (2), y being considered as a function of x, are all 0; and it is plain that if any two be 0, the third will be 0 also; hence, differentiating the two first, we have

$$r' + s'\frac{dy}{dx} = 0, \quad \text{and } s' + t'\frac{dy}{dx} = 0,$$

where $\frac{dy}{dx}$ fixes the position of the rectilinear directrix for which the expressions (2) remain constant. Eliminating, then, $\frac{dy}{dx}$, we obtain the following equation, which must hold for every directrix, viz.

$$r't' - s'^2 = 0 \ldots (3).$$

This, therefore, is the equation which the differential of the equation of every developable surface must accord with; or, in usual terms, it is the differential equation of developable surfaces in general.

(196.) We shall exhibit another method of obtaining the equation (3), from the general equations (2), art. (192). Differentiating the first of these, in which α is a function of x and y implied in the second, we have the two partial differential equations

$$p' = f'(\alpha)\frac{d\alpha}{dx} + \phi(\alpha) + x\phi'(\alpha)\frac{d\alpha}{dx} + y\psi'(\alpha)\frac{d\alpha}{dx}$$

$$q' = f'(\alpha)\frac{d\alpha}{dy} + x\phi'(\alpha)\frac{d\alpha}{dy} + \psi(\alpha) + y\psi'(\alpha)\frac{d\alpha}{dy},$$

but, in virtue of the second of the equations (2), these become

$$p' = \phi(\alpha), \quad q' = \psi(\alpha),$$

consequently, p' must be a function of q', and may therefore be represented by

$$p' = \pi(q').$$

Eliminating now the arbitrary function π by differentiation, as in (58), we have

$$\frac{r'}{s'} = \frac{s'}{t'} \quad \therefore r't' - s'^2 = 0.$$

as before.

(197.) We have as yet considered only the simplest class of surfaces, whose intersections, with their consecutive surfaces, are given by the general equations (1), art. (191), viz. plane surfaces, the intersections being straight lines. It is obvious, however, that whatever be the surfaces, the intersections are still given by the equations (1), and the envelope of these surfaces, found by eliminating from them the arbitrary parameter α. This parameter, however, may enter the equation of any particular family of surfaces in an infinite variety of different forms and ways; it may enter into only one of its terms, or be combined with several; a simple power only of it may enter, or a complicated function; and still, entering only as a parameter, the general equation, under all these changes, will still preserve the same character, and represent but one family of surfaces. With the envelope, however, it will be different; this depends as well on the arbitrary parameter, as on the variables which enter the general equation, since the value of this parameter must

be found from one of the equations (1), in terms of x, y, z, and this value substituted in the other for the equation of the envelope. Nevertheless, since, as just observed, the individual surfaces represented by the two equations (1), for every particular value of the parameter in whatever form it may enter, is always of the same degree, it follows that each individual intersection (1), will uniformly be a curve of the same order, and which will change its order only when the order of the surface changes. This curve of intersection, or of contact, common to all the envelopes of the same family of surfaces, is called, by *Monge*, the *Characteristic*.

Considering any of the characteristics (1) separately, we may enquire what are the points in which it is intersected by the consecutive characteristic; and the method of determining these intersections is analogous to that already explained in (107) and (186), that is, we must combine with the equations (1) of this curve their differentials taken relatively to α; hence, the consecutive intersections for any particular position of the characteristic, will be determined by the equations

$$F(x, y, z, \alpha) = 0$$

$$\frac{dF(x, y, z, \alpha)}{d\alpha} = 0$$

$$\frac{d^2F(x, y, z, \alpha)}{d\alpha^2} = 0,$$

each of these separately represents a surface, any two together a line common to both, and all three the point or points common to their intersection, α being considered constant in the results of the operations indicated. By solving these three equations for x, y, and z, we shall obviously obtain known values for the coordinates of the points of intersection required, which of course are all situated on the envelope.

Now, if from the three equations above, we eliminate α, we shall have two equations in x, y, z, existing together; which, being the same for the intersections of *every* pair of consecutive characteristics, must represent the locus of these intersections, and be situated on the envelope. It will therefore be a line which touches and encompasses all the characteristics, in the same manner as the envelope touches and embraces all the enveloped surfaces. It must then form an edge of the

envelope, or the line in which its sheets terminate, and it is therefore called, by *Monge*, the *edge of regression of the envelope*. In the developable surfaces, we have seen that the characteristic is a straight line, and the consecutive intersections of the characteristic, in every position, obviously form the edge which limits the locus of the characteristics, that is, the developable surface.

The consideration of envelopes, characteristics, and edges of regression, have been successfully employed by *Monge*, and succeeding writers, to remove several difficulties in the higher departments of the integral calculus, that do not appear to be otherwise clearly explicable; but it would be out of place here to do more than to hint at the importance of these researches; to pursue them to their fullest extent the advanced student must have recourse to the profound work of *Monge*, before referred to, viz. *Application de l'Analyse à la Géométrie*. We shall conclude the present chapter with one example on the determination of the envelope.

(198.) The centre of a sphere of given radius moves along a given plane curve, it is required to determine the surface which envelopes the sphere in every position.

Let the equation of the given curve, along which the centre moves, be

$$\beta = \phi(\alpha) \ldots . \quad (1),$$

so that for every abscissa α of this curve, the ordinate corresponding will be $\phi(\alpha)$; therefore, the variable coordinates of the centre of the sphere are $\alpha, \phi(\alpha)$; hence its equation, in any position, is

$$(x-\alpha)^2 + (y-\phi(\alpha))^2 + z^2 = r^2 \ldots (2),$$

so that the equations of the characteristic are,

$$\left.\begin{aligned} (x-\alpha)^2 + (y-\phi(\alpha))^2 + z^2 &= r^2 \\ x-\alpha + (y-\phi(\alpha))\,\phi'(\alpha) &= 0 \end{aligned}\right\} \ldots (3).$$

The last equation is that of a plane, passing through the point $(\alpha\ \phi(\alpha))$, or centre of the sphere; it is, moreover, perpendicular to the tangent to the curve (1) at this point, for the equation of this tangent is

$$(\beta' - \beta) = \phi'(\alpha)\,(\phi(\alpha') - \phi(\alpha)),$$

and that above is

$$y - \phi(\alpha) = -\frac{1}{\phi'(\alpha)}(x - \alpha),$$

so that whatever be the form of ϕ, the characteristic is always a great circle of the moveable sphere, of which the plane is normal to the curve. The species of the curve which is the characteristic being, however, constant, as observed in art. (197,) however ϕ and α may vary, the species may be at once determined by assuming $\alpha = 0$, $\phi(\alpha) = 0$, which reduces the equations of the characteristic to

$$\left.\begin{matrix} x^2 + y^2 + z^2 = r^2 \\ x = 0 \end{matrix}\right\} \text{ or } y^2 + z^2 = r^2,$$

which belongs to a circle; the species, therefore, is a curve of the second order.

To determine the equation of the envelope, we must eliminate α from (3), and the resulting equation in x, y, z, will belong to the envelope; thus, if the curve (1) be a circle of radius a, then

$$\phi(\alpha) = \sqrt{a^2 - \alpha^2} \therefore \phi'(\alpha) = \frac{-\alpha}{\sqrt{a^2 - \alpha^2}},$$

substituting these values in the equations (3), they become

$$(x - \alpha)^2 + (y - \sqrt{a^2 - \alpha^2})^2 + z^2 = r^2,$$

$$\alpha y = x\sqrt{a^2 - \alpha^2},$$

and determining, from this last equation, the expression for α, and substituting it in the preceding, we shall obtain, finally,

$$(a \pm \sqrt{x^2 + y^2})^2 = r^2 - z^2,$$

for the equation of the envelope.

CHAPTER VI.

ON CURVES OF DOUBLE CURVATURE.

(199.) In the preliminary chapter to the present Section, we investigated the expressions for the tangent lines and normal planes to these curves; we shall now discuss their general theory. As, however, in the course of this discussion, we shall sometimes have occasion to employ the differential expression for the arc of a curve of double curvature, we shall commence by seeking the form of this expression.

(200.) We know that the projecting surface of every curve of double curvature, is a cylindrical surface, (see *Anal. Geom.*, p. 311;) if, therefore, this cylindrical surface be developed, the curve will become plane, its length will be unaltered, and the curvilinear base of the projecting cylinder, which we shall here suppose to be vertical, will become a straight line on the plane of xy; hence, for the plane curve referred to this straight line t, and the axis of z, we shall have (87) the expression

$$(ds)^2 = (dz)^2 + (dt)^2,$$

but t being itself, in reality, the arc of a plane curve, we have

$$(dt)^2 = (dx)^2 + (dy)^2,$$

hence, by substitution,

$$(ds)^2 = (dx)^2 + (dy)^2 + (dz)^2,$$

which is the differential expression required.

Osculation of Curves of Double Curvature.

(201.) Let

$$y=f(x),\quad z=\mathrm{F}(x)\ \ldots\ (1),$$

and

$$\mathrm{Y}=\psi(x),\quad \mathrm{Z}=\Psi(x)\ \ldots\ (2),$$

be the equations of two curves of double curvature, or rather of the projections of these curves on the planes of xy, xz. Then, if we consider the constants, a, b, c, &c. which enter the first pair of equations as known, and the constants A, B, C, &c. belonging to the second pair as arbitrary, these latter may be determined so that the curve to which they belong may touch the proposed or fixed curve (1), in any given point, more intimately than any other curve of the family (2). For, giving to x any increment, h, we have, by Taylor's theorem,

$$y'=y+\frac{dy}{dx}h+\frac{d^2y}{dx^2}\frac{h^2}{1\cdot 2}+\&c.$$

$$\mathrm{Y}'=\mathrm{Y}+\frac{d\mathrm{Y}}{dx}h+\frac{d^2\mathrm{Y}}{dx^2}\frac{h^2}{1\cdot 2}+\&c.$$

and it has been shown (88), that if the constants which enter the first of the equations (2), be determined, all of them from the conditions

$$y=\mathrm{Y},\quad \frac{dy}{dx}=\frac{d\mathrm{Y}}{dx},\quad \frac{d^2y}{dx^2}=\frac{d^2\mathrm{Y}}{dx^2},\ \&c.\ \ldots\ (3),$$

the projection of the curve (2), on the plane of xy, shall touch more intimately the projection of (1) on that plane, than the projection of any other curve of the family (2).

In like manner, if the constants which enter the second of the equations (2), be all of them determined from the conditions

$$z=\mathrm{Z},\quad \frac{dz}{dx}=\frac{d\mathrm{Z}}{dx},\quad \frac{d^2z}{dx^2}=\frac{d^2\mathrm{Z}}{dx^2},\ \&c.\ \ldots\ (4),$$

the projection of the curve (2) on the plane of xz, will touch more in-

timately the projection of (1) on that plane, than the projection of any other curve of the family (2).

It is clear, therefore, that if all the constants in the equations (2) be determined conformably to the conditions (3), (4), the curve (2) will touch more intimately the curve (1) in space, than any other curve of the family (2), and that the contact will be the less intimate as the conditions (3), (4), satisfied by the arbitrary constants, are fewer.

The conditions for simple contact, or contact of the first order, are evidently

$$y = Y, \quad z = Z, \quad \frac{dy}{dx} = \frac{dY}{dx}, \quad \frac{dz}{dx} = \frac{dZ}{dx} \ldots\ldots (5),$$

and, for *contact of the second order*, we must have the two additional conditions

$$\frac{d^2y}{dx^2} = \frac{d^2Y}{dx^2}, \quad \frac{d^2z}{dx^2} = \frac{d^2Z}{dx^2},$$

and so on.

(202.) From these principles, we may very easily deduce the equations of the tangent at any point of a curve of double curvature.

Thus the equations of any straight line in space, are

$$y = Ax + B; \quad z = A'x + B' \ldots\ldots (1),$$

and these correspond to the equations (2) above; and as four arbitrary constants enter, the conditions (5) may be fulfilled by them; thus, taking the two last conditions, we have, by accenting the variables of the curve,

$$\frac{dy'}{dx'} = A, \quad \frac{dz'}{dx'} = A',$$

and, therefore, the two first require that

$$B = y' - \frac{dy'}{dx'}x', \quad B' = z' - \frac{dz'}{dx'}x',$$

so that the equations (1) of the tangent, through any point (x', y', z') are

$$\left.\begin{aligned} y - y' &= \frac{dy'}{dx'}(x - x') \\ z - z' &= \frac{dz'}{dx'}(x - x') \end{aligned}\right\} \quad \ldots\ (2).$$

PROBLEM I.

(203.) To determine the osculating circle, at any point in a curve of double curvature.

In finding the osculating circle, at any point of a plane curve, we had, of course, the plane of that curve given, but, in the present case, we have to determine both the plane of the circle and its radius. Now let us suppose that r is the radius of the osculating circle, and that α, β, γ, are the coordinates of its centre; then it is plain, that the circle will be a great circle of the sphere whose equation is

$$(x-\alpha)^2 + (y-\beta)^2 + (z-\gamma)^2 = r^2 \quad \ldots\ (1);$$

and, since the plane of the circle passes through the point (α, β, γ), its equation must be of the form

$$x - \alpha + m(y-\beta) + n(z-\gamma) = 0 \quad \ldots\ (2).$$

These two equations, combined with those of the proposed curve, give the values of x, y, z, common to all, and therefore belong to the point where the circle (1), (2) meets the curve. We have, therefore, to differentiate the equations (1), (2) successively, and to consider, agreeably to the conditions (3), (4), art. (201), that the resulting differential coefficients belong as well to the proposed curve at the point, as to this circle. For contact of the first order, we have

$$(x-\alpha) + (y-\beta)p' + (z-\gamma)q' = 0 \quad \ldots\ (3),$$

$$1 + mp' + nq' = 0 \quad \ldots\ (4);$$

and for contact of the second order we have, in addition,

$$1 + p'^2 + q'^2 + p''(y-\beta) + q''(z-\gamma) = 0 \quad \ldots\ (5).$$

$$mp'' + nq'' = 0 \quad \ldots\ (6).$$

All these six conditions, therefore, must exist for the contact at the point (x, y, z) in the proposed curve to be of the second order; and, as the equations (1), (2), of the touching curve, contain six disposable constants, viz. $\alpha, \beta, \gamma, r, m, n$, all these conditions may be fulfilled, but no more; hence the circle, determined agreeably to these conditions, will touch the proposed curve more intimately than any other, that is, it will be the *osculating circle*. From equations (4) and (6) we get

$$m = \frac{q''}{q'p'' - p'q''}, \quad n = -\frac{p''}{q'p'' - p'q''};$$

hence, equation (2) becomes

$$x - \alpha + \frac{q''}{q'p'' - p'q''}(y - \beta) - \frac{p''}{q'p'' - p'q''}(z - \gamma) = 0,$$

or

$$z - \gamma = \frac{q'p'' - p'q''}{p''}(x - \alpha) + \frac{q''}{p''}(y - \beta) \ldots . (7);$$

hence, the three conditions (2), (4), (6), determine the plane of the osculating circle, and which is called the *osculating plane*, through the proposed point (x, y, z). Equation (7) then represents this plane.

For the coordinates of the centre of the osculating circle we have, from equations (1), (2), (3),

$$x - \alpha = \frac{(np' - mq')r}{M}, \quad y - \beta = -\frac{(n - q')r}{M}, \quad z - \gamma = \frac{(m - p')r}{M},$$

where M is put for the expression

$$\sqrt{\{(np' - mq')^2 + (n - q')^2 + (m - p')^2\}}.$$

Substituting these values in (5), we have, for the radius of osculating circle,

$$r = \frac{(1 + p'^2 + q'^2)M}{(n - q')p'' - (m - p')q''}.$$

Hence, putting for m and n the values already deduced, and restoring the value of M, we have

$$r = -\frac{(1+p'^2+q'^2)^{\frac{3}{2}}}{\sqrt{\{p''^2+q''^2+(p'q''-q'p'')^2\}}},$$

$$\alpha = x - \frac{(1+p'^2+q'^2)(p'p''+q'q'')}{p''^2+q''^2+(p'q''-q'p'')^2},$$

$$\beta = y + \frac{(1+p'^2+q'^2)\{p''+q'(q'p''-p'q'')\}}{p''^2+q''^2+(p'q''-q'p'')^2},$$

$$\gamma = z - \frac{(1+p'^2+q'^2)\{q''-p'(q'p''-p'q'')\}}{p''^2+q''^2+(p'q''-q'p'')^2}.$$

(204.) The expression for r may be rendered more general, by considering the independent variables as arbitrary; in which case we have (66)

$$p'' = \frac{(d^2y)(dx)-(d^2x)(dy)}{(dx)^3}, \quad q'' = \frac{(d^2z)(dx)-(d^2x)(dz)}{(dx)^3}.$$

Also (199)

$$\frac{(ds)^2}{(dx)^2} = 1+p'^2+q'^2;$$

hence, making these substitutions in the above expression, we have

$$r = \frac{(ds)^3}{\sqrt{\{(dx)(d^2y)-(dy)(d^2x)\}^2+\{(dz)(d^2x)-(dx)(d^2z)\}^2+\{(dy)(d^2z)-(dz)(d^2y)\}^2}}.$$

(205.) If it were required to determine the circle having contact of the first order, merely with the proposed curve, only the conditions (1), (2), (3), (4) must be satisfied; the conditions (2), (4) determine the plane of this circle, that is, the tangent plane; as however the condition (4) leaves one of the constants, m, n, arbitrary, the tangent plane is not fixed, but may take an infinite variety of positions; yet as it must necessarily pass through the linear tangent, which is fixed, it follows that a plane through this, and revolving round it, is a tangent plane in every position; in one of which it touches the curve with a contact of the second order, and thus becomes the osculating plane.

(206.) There is another method of determining the equation of the osculating plane, very generally employed by French authors; they consider a curve of double curvature to have, at every point, two *consecutive elements*, or infinitely small contiguous arcs in the same plane, but not more, the plane of these elements being the osculating plane at the point. The process, then, is to assume the equation of a plane through the point

$$x - x' + m(y - y') + n(z - z') = 0 \quad \ldots \quad (1),$$

and to subject it to the condition of passing also through the points

$$(x + \mathrm{d}x', \quad y' + \mathrm{d}y', \quad z' + \mathrm{d}z')$$

and

$$x' + 2\mathrm{d}x' + \mathrm{d}^2x', \quad y' + 2\mathrm{d}y' + \mathrm{d}^2y', \quad z' + 2\mathrm{d}z' + \mathrm{d}^2z'.$$

Such a process the student will at once perceive to be exceedingly exceptionable; for, besides the vague notion attached to the infinitely small consecutive arcs, the expressions $x + \mathrm{d}x$, $y + \mathrm{d}y$, and the like, mean no more in the language of the *differential* calculus, than x, y, &c.; for $\mathrm{d}x$, $\mathrm{d}y$, &c. are not infinitely small, but absolutely 0, as we have all along been careful to impress on the mind of the student. The process is, however, susceptible of improvement, thus: suppose the plane (1) passing through one point (x', y', z') of the curve, passes also through a second point, of which the abscissa is $x' + \Delta x'$, where $\Delta x'$ means *the increment of* x; then, substituting $x' + \Delta x'$ for x', the equation (1) becomes

$$x - x' + m(y - y') + n(z - z') - (\Delta x' + m\Delta y' + n\Delta z') = 0 \ldots (2),$$

which, in virtue of (1), is the same as

$$\Delta x' + m\Delta y' + n\Delta z' = 0,$$

or

$$1 + m\frac{\Delta y'}{\Delta x'} + n\frac{\Delta z'}{\Delta x'} = 0.$$

Suppose now that these two points merge into one; that is, let $\Delta x' = 0$, then

$$1 + m\frac{dy'}{dx'} + n\frac{dz'}{dx'} = 0 \ldots\ldots (3);$$

hence the plane becomes determinable by the conditions (1), (3).

Again, let this plane pass through a third point, $x' + \Delta x'$, then substituting this for x' in both the equations (1), (3), they will furnish the additional condition

$$m\Delta\frac{dy'}{dx'} + n\Delta\frac{dz'}{dx'} = 0;$$

hence, dividing by $\Delta x'$ and supposing this third point to coincide with the former; that is, supposing $\Delta x' = 0$, we have the new condition

$$m\frac{d^2y'}{dx'^2} + n\frac{d^2z'}{dx'^2} = 0 \ldots\ldots (4).$$

The equations (3) and (4) determine m and n, and thence the plane (1), which is such as to pass through but one point of the curve, and at the same time to be so placed that the most minute variation from this position will cause it to pass through *three* points of the curve.

(207.) By whatever process the osculating plane is determined, the radius of the osculating circle may be easily found from considerations different from those at (203). For, as the linear tangent to the curve, must also be tangent to the osculating circle, it follows that the centre of this circle must be on the normal plane, as well as on the osculating plane; it must, therefore, lie in the line of intersection of this normal plane, with its consecutive normal plane; hence, if this line be determined, the combination of its equation with that of the osculating plane, will give the point sought. Now (188) the line of intersection of consecutive normal planes is

$$\left.\begin{aligned} x - x' + p'(y - y') + q'(z - z') &= 0 \\ p''(y - y') + q''(z - z') - p'^2 - q'^2 - 1 &= 0 \end{aligned}\right\},$$

therefore the centre is to be determined by combining these equations with that of the osculating plane, viz.

$$z-z'=\frac{q'p''-p'q''}{p''}(x-x')+\frac{q''}{p''}(y-y'),$$

being precisely the same equations as those employed before, for the same purpose. If the origin be at the point, and the tangent be the axis of x, then x', y', z', p', q', are each 0; therefore the equations of the line of intersection are

$$x=0,\quad y=-\frac{q''}{p''},\quad z+\frac{1}{p''},$$

and the equation of the osculating plane

$$p''z-q''y=0;$$

this, therefore, is perpendicular to the line of intersection, (*Anal. Geom.*, p. 234.)

(208.) The expressions in (203) for the coordinates of the centre of the osculating circle, will become very simple by introducing the substitutions furnished by art. (204); the results of these substitutions will be

$$\alpha=x+r^2\frac{d^2x}{ds^2},\quad \beta=y+r^2\frac{d^2y}{ds^2},\quad \gamma=z+r^2\frac{d^2z}{ds^2},$$

the independent variable being s. (*See Note* D.)

PROBLEM II.

(209.) To determine the centre and radius of spherical curvature at any point in a curve of double curvature.

We are here required to determine a sphere in contact with the proposed curve at a given point, such that a line on its surface in the direction of the proposed may, in the vicinity of the point, be closer to the curve than if any other sphere were employed. In the direction of the curve, the z and the y of the sphere must be both functions of x; so that the equation of the sphere is resolvable into two, corresponding

to the equations (2), art. (202), which two equations belong to the curve which osculates the proposed. The actual resolution of the equation into two is obviously unnecessary; it will be sufficient in that equation to consider x as the only independent variable.

The general equation of a sphere is

$$(x-\alpha)^2+(y-\beta)^2+(z-\gamma)^2=r^2 \quad \ldots\ldots (1),$$

and the particular sphere required will be that whose constants are determined from the following differential equations:

$$x-\alpha+p'(y-\beta)+q'(z-\gamma)=0 \quad \ldots\ldots (2),$$

$$p''(y-\beta)+q''(z-\gamma)+1+p'^2+q'^2=0 \quad \ldots\ldots (3),$$

$$p'''(y-\beta)+q'''(z-\gamma)+3(p'p''+q'q'')=0 \quad \ldots\ldots (4).$$

These four equations fix the values of the parameters α, β, γ, r, and, therefore, determine both the position and magnitude of the osculating sphere. If the origin of coordinates be at the proposed point, and the linear tangent be taken for the axis of x, the determination becomes easy, for x, y, z, being each $=0$, as also p', q', the foregoing equations (2), (3), (4), become

$$\alpha=0, \quad p''\beta+q''\gamma=1, \quad p'''\beta+q'''\gamma=0,$$

$$\therefore \beta=\frac{q'''}{p''q'''-p'''q''}, \quad \gamma=\frac{p'''}{q''p'''-q'''p''};$$

hence, by substitution in (1),

$$r=\frac{\sqrt{p'''^2+q'''^2}}{p''q'''-q''p'''}.$$

(210.) We already know that if to every point in a curve of double curvature normal planes be drawn, the intersections of these planes with the consecutive normal planes will be the characteristics of the developable surface which they generate, and the intersection of any characteristic with the consecutive characteristic will be a point in the edge of regression, corresponding to the given point on the proposed curve. Now

equation (2) above being that of the normal plane, this point is determined by precisely the same equations (2), (3), (4), as determine the centre of spherical curvature; these points, therefore, are one and the same, as might be expected; hence, the locus of the centres of spherical curvature forms the edge of regression of the developable surface generated by the intersections of the consecutive normals. If then, by means of one of the equations of the proposed curve and the three equations of condition mentioned, we eliminate x, y, z, and then perform the same elimination by means of the other equation of the curve and the same conditions, we shall obtain two resulting equations in α, β, γ, which will be the equations of the edge of regression.

PROBLEM III.

(211.) To determine the points of inflexion in a curve of double curvature.

Since a curve of double curvature, as its name implies, has curvature in two directions, if at any point its curvature in one direction changes from concave to convex, the point is called a point of *simple inflexion*. But if at the same point there is also a like change of curvature in the other direction, the point is then said to be one of *double inflexion*. In other words, if in but one projection the tangent cross the projected curve, the point is one of simple inflexion; but if the tangent cross the curve in both projections, then the point is one of double inflexion. As in plane curves the tangent line has contact one degree higher at a point of inflexion, so here the contact of the osculating plane is one degree higher. Hence, at such a point, besides the conditions in (203), which fix the osculating plane, we must, at a point of simple inflexion, have the additional condition arising from differentiating (6), viz.

$$mp''' + nq''' = 0.$$

Eliminating $\frac{m}{n}$ from this and equation (5), we have

$$p''q''' = q''p''',$$

which condition renders the expression for the radius of spherical

curvature at the point infinite, as it ought. Unless, therefore, this condition exist, the point cannot be one of inflexion; but the point for which the condition holds *may* be one of inflexion, yet to determine this the curve must be examined in the vicinity of the point.

As to points of double inflexion, it is evident, from what has been said (123) with respect to plane curves, that such points must fulfil the conditions

$$p'' = 0, \text{ or } \infty, \quad q'' = 0 \text{ or } \infty,$$

and these render the radius r, of absolute curvature, infinite or 0.

Evolutes of Curves of Double Curvature.

(212.) In speaking of the evolutes of plane curves we observed (105) that the evolute of any plane curve was such, that if a string were wrapped round it and continued in the direction of its tangent till it reached a point in the involute curve, the unwinding of this string would cause its extremity to describe the involute. But besides the *plane* evolute hitherto considered, there are numberless curves of double curvature, round which the string might be wound and continued in the direction of a tangent till it reached the involute, which would equally, by unwinding, describe this involute; and generally every curve, whether plane or of double curvature, has an infinite number of evolutes, as we are about now to show.

(213.) If through the centre of a circle, and perpendicular to its plane, an indefinite straight line be drawn, and any point whatever be taken in this line, then it is obvious that the point will be equally distant from every point in the circumference of the circle; so that, if a line be drawn from it to the circumference, this line, in revolving round the perpendicular under the same angle, will describe the circumference. Such a point is called a *pole* of the circle; so that every circle has an infinite number of poles, the locus of which is determined when the positions of any two are given.

(214.) Now, as respects curves of double curvature, we have seen

that the centre of the circle of absolute curvature corresponding to any point, is in the line where the normal at this point is intersected by its consecutive normal, the centre itself being that point in this line where it pierces the osculating plane, which (204) is the plane drawn through the tangent line perpendicular to this line of intersection, or characteristic; hence the characteristic corresponding to any point in the curve, is the locus of the poles of curvature at that point; and the intersection of this characteristic, with the perpendicular to it from the corresponding point of the curve, is that particular pole which is the centre of absolute curvature, the perpendicular itself being the radius.

As the locus of the poles corresponding to any point is no other than the characteristic, the locus of all the poles corresponding to all the points of the curve must be the locus of all the characteristics, and therefore (192) *a developable surface.*

(215.) Suppose now through any point, P, of the curve, a normal plane is drawn of indefinite extent, the characteristic or line of poles corresponding to the point will be in this plane; let, therefore, any straight line be drawn from P to intersect this line of poles in the point Q, and be continued indefinitely. If this normal plane be conceived to move, so that, while P describes the proposed curve, the plane continues to be normal, the characteristic will undergo a corresponding motion, and will generate the developable surface corresponding to the curve described by P; and this motion of the characteristic will cause a corresponding motion of the point Q, not only in space, but along the arbitrary line from P, which has no motion in the moving plane. As, therefore, Q moves along the characteristic, successive portions QQ′ of the line, PQ will apply themselves to the surface which the moveable characteristic generates, and there form a curve to which always the unapplied portion QP is a tangent. Now the normal plane being in every position tangent to the surface throughout the whole length of the characteristic, it is obvious that, in the above generation of this surface, nothing more in effect has been done than the bending of the original normal plane, supposed flexible, into a developable surface. If, therefore, we now perform the reverse operation, that is, if we unbend the normal plane, the point P will describe the curve of double

curvature, and the curve QQ′ traced on the developable surface will become the straight line PQ; so that the curve of double curvature may be described by the unwinding of a string wrapped about the curve Q′Q, and continued in the direction QP of its tangent, till it reaches the point P in the proposed curve. It follows, therefore, that the curve Q′Q is an evolute of the curve of double curvature proposed; and, moreover, that, as the line PQ originally drawn was quite arbitrary, *the proposed curve has an infinite number of evolutes, situated on the developable surface which is the locus of the poles of the proposed;* hence the locus of the poles is the locus of the evolutes.

If the original line PQ be perpendicular to the corresponding line of poles or characteristic, then, since this characteristic moves in the moving plane while PQ remains fixed, PQ cannot continue to be perpendicular to the characteristic; but the radius of absolute curvature is always perpendicular to the characteristic, this radius therefore cannot continue to intersect the characteristic in the point Q; so that *the locus of the centres of absolute curvature is not one of the evolutes of the proposed curve.*

(216.) Should the curve which we have all along considered of double curvature be plane, then, indeed, since the characteristics are all parallel, and perpendicular to the plane of the curve, the line PQ, once perpendicular, will be always perpendicular to the characteristic; so that then Q will coincide with the centre of curvature, PQ being no other than the radius of curvature, the locus of the centres being the plane evolute before considered. But when PQ is not drawn perpendicular to the original characteristic, but is inclined to it at an angle α, then it always preserves this inclination during the generation of the cylindrical surface, which is the locus of the poles; therefore every curvilinear evolute of a plane curve is a *helix* described on the surface of the cylinder, which is the locus of the poles of the plane curve.

Every curve traced on the surface of a sphere, has, for the locus of its evolutes, a conical surface whose vertex is at the centre of the sphere; because the normal planes to the curve being also normal planes to the spheric surface, all pass through the centre.

(217.) From what has now been said, it is obvious that if from any point in a curve a line be drawn to touch the developable surface which is the locus of its poles, and its prolongation be wound about the surface without *twisting*,* it will trace one of the evolutes; and, as the string may be drawn to touch the surface in every possible direction, it follows that every developable line on the surface will be an evolute. If the curve be plain, the evolutes are all on the cylindrical surface whose base is the plane evolute.

As obviously a developable line is the shortest on the surface that can join its extremities, it follows that the shortest distance between two points of an evolute measured on the surface, is the arc of that evolute between them.

PROBLEM IV.

(218.) Having given the equation of a curve of double curvature to determine those of any one of its evolutes.

All the evolutes of the curve being on the same developable surface, the equation of this surface must be common to them all; and we have already seen (192) how the equation of the surface is to be determined, so that it only remains to find for each evolute a particular equation which distinguishes it from all the others, and determines its course on the developable surface. In order to this, let us consider that each evolute must be such that the prolongation of its tangent at any point always cuts the involute, or, which is the same thing, the tangent to the projection of the evolute at any point passes through the corresponding point in the projection of the evolute; therefore, considering the plane of xy as that of projection, we have, for the tangent at any point (x', y')

---

* This is what I understand *Monge* to mean, when he says *(App. de l'Anal. de Géom.*, p. 348,) "si l'on plie *librement* sur cette surface le prolongement de cette tangente." It seems not improper to call such lines placed on a developable surface *developable lines*, and those which form curves on the developed surfaces *twisted lines*. Of these two species of lines all the former are evolutes, but none of the latter are.

in the projected evolute,

$$Y - y' = \frac{dy'}{dx'}(X - x');$$

and, since the same line passes through a point (x, y) in the projected involute, its equation is also

$$Y - y' = \frac{y' - y}{x' - x}(X - x'),$$

$$\therefore \frac{dy'}{dx'} = \frac{y' - y}{x' - x};$$

hence, combining this equation with that of the developable surface, determined agreeably to the process pointed out in art. (192), and eliminating x, y being a given function of x, we shall have two equations in x', y', z', of which one will contain partial differential coefficients of the first order, and which together will represent all the evolutes. To find that particular one which is fixed by any proposed condition, it will be necessary to discover, by the aid of the integral calculus, the primitive equation from which the differential equation mentioned is deducible; this primitive equation will involve an arbitrary constant, whose value may be fixed by the proposed condition, and thus the equations of the particular evolute will be determined.

We shall terminate this Section by subjoining a few miscellaneous propositions.

CHAPTER VII.

MISCELLANEOUS PROPOSITIONS.

PROPOSITION I.

(219.) To prove that the locus of all the linear tangents at any point of a curve surface is necessarily a plane.

This property we have hitherto assumed; it may, however, be demonstrated as follows:

Let the equation of any curve surface be

$$z = f(x, y) \ldots . (1),$$

x and y being the independent variables.

Through any given point on this surface let any curve be traced; then the projection of this curve on the plane of xy will be represented by

$$y = \phi(x) \ldots . (2),$$

which will equally represent the projecting cylinder; hence the combination of the equations (1), (2), completely determines the curve, and its projection on the plane of xz may be found by eliminating y from these equations; the result of this elimination will be the equation

$$z = f(x, \phi(x)) = \psi(x) \ldots . (3),$$

therefore, since the linear tangent in space is projected into tangents to these two curves (2), (3), its equations must be

$$\left.\begin{aligned} y - y' &= \frac{d\phi(x')}{dx'}(x - x') \\ z - z' &= \frac{d\psi(x')}{dx'}(x - x') \end{aligned}\right\}$$

where x', y', z', are the coordinates of the proposed point on the surface.

Now $\frac{d\psi(x)}{dx}$ is the total differential coefficient derived from the function $z = f(x, y)$, in which y is considered as a function of x, given by the equation (2), that is

$$\frac{d\psi(x)}{dx} = \left\{\frac{dz}{dx}\right\} = p' + q' \frac{d\phi(x)}{dx};$$

hence, by substitution, the equations of the tangent in space become

$$\left.\begin{aligned} y - y' &= \frac{d\phi(x')}{dx'}(x - x') \\ z - z' &= \left(p' + q' \frac{d\phi(x')}{dx'}\right)(x - x') \end{aligned}\right\} \quad \ldots\ldots (4).$$

Now, to obtain the locus of the tangents whatever be the curve through the point (x', y', z'), we must eliminate the function $\phi(x)$, on which alone the nature of the curve depends. Executing, then, this elimination by means of the equations (4) and there results for the required locus the equation

$$z - z' = p'(x - x') + q'(y - y'),$$

which is that of a plane.

PROPOSITION II.

(220.) Given the algebraic equation of a curve surface to determine whether or not the surface has a centre.

That point is called the centre which bisects all the chords drawn through it, so that if the equation of the surface is satisfied for any constant values x', y', z', it will equally be satisfied for the same values taken negatively, that is, for $-x'$, $-y'$, $-z'$, provided the origin of coordinates be placed at the centre; so that if no point exists for the origin of coordinates, in reference to which the equation

$$f(x, y, z) = 0$$

of the surface remains the same, whether the signs of the variables be assumed all + or all —, then we may conclude also that no centre exists.

The mode of proceeding, therefore, is to assume the indeterminates $x_{,}, y_{,}, z_{,}$, for the coordinates of the unknown centre, and to transport the origin of the axes to that point by substituting in the equation of the surface $x + x_{,}, y + y_{,}, z + z_{,}$ for x, y, z. This done we may readily deduce equations of condition, which will give the proper values of $x_{,}$, $y_{,}$, $z_{,}$, if a centre exists, or will show, by their incongruity, that the surface has no centre. Thus, suppose the equation of the surface is of an even degree, then we must equate to 0 the coefficients of all the *odd* powers and combinations of x, y, z, since the terms into which these enter would change signs when the variables change signs; we obtain in this way the equations of condition. If the equation of the surface be of an odd degree, then we must equate to zero the coefficients of all the *even* powers and combinations of x, y, z; so that only odd powers and combinations may effectively enter the equation, for then whether the variables be all + or all — the function $f(x, y, z)$ will still be 0.

Now the differential calculus furnishes us at once with the means of obtaining the several expressions which we must equate to zero without actually substituting $x + x_{,}, y + y_{,}, z + z_{,}$ for x, y, z, in the equation of the surface. For if we conceive these substitutions made in the function $f(x, y, z)$, we may consider the result as arising from $x_{,}, y_{,}, z_{,}$ taking the respective increments x, y, z, and we know that every such function may, by Taylor's theorem, be developed according to the powers and combinations of the increments; and that the several terms of the development consist each of the partial differential coefficients of the preceding term, the first being $f(x_{,}, y_{,}, z_{,})$. Hence, if the coefficients of the first powers of x, y, z, are to be respectively zero, then we have to equate to zero each of the partial coefficients derived from $u_{,} = f(x_{,}, y_{,}, z_{,}) = 0$, or, which is the same thing, from $u = f(x, y, z) = 0$ the proposed equation; if the coefficients of the second powers and combinations of x, y, z, are to be rendered each 0, then we shall have to equate to zero each partial coefficient derived from again differentiating, and so on.

As an illustration of this, let the general equation of surfaces of the second order

$$\left.\begin{aligned} Ax^2 + A'y^2 + A''z^2 + 2Byz + 2B'zx + 2B''xy \\ + 2Cx + 2C'y + 2C''z + E \end{aligned}\right\} = 0 = u \quad \ldots \quad (1)$$

be proposed, then the degree of the equation being even, the coefficients of the odd powers of the variables in the equation, arising from putting $x + x_{,}$, $y + y_{,}$, $z + z_{,}$, for x, y, z are to be equated to 0, and as the equation is but of the second degree, these odd powers will be of the first; hence we have merely to equate the first partial differential coefficients to 0, that is

$$\left.\begin{aligned} \frac{du}{dx} &= Ax + B'z + B''y + C = 0 \\ \frac{du}{dy} &= A'y + Bz + B''x + C' = 0 \\ \frac{du}{dz} &= A''z + B'x + By + C'' = 0 \end{aligned}\right\} \quad \ldots \quad (2).$$

The values of x, y, z, deduced from these equations are the coordinates $x_{,}$, $y_{,}$, $z_{,}$, of the centre. These values may be represented by

$$x_{,} = \frac{N}{D}, \quad y_{,} = \frac{N'}{D}, \quad z_{,} = \frac{N''}{D},$$

where

$$D = AB^2 + A'B'^2 + A''B''^2 - AA'A'' - 2BB'B'',$$

so that the surface has a centre if D is not 0; but if $D = 0$ and the numerators all finite, the surface has no centre; and, lastly, if $D = 0$ and either of the numerators also 0, then the surface has an infinite number of centres, and is, therefore, cylindrical.

The equations of condition (2) are the same as those at page 292 of the Analytical Geometry.

PROPOSITION III.

(221.) To determine the equation of the diametral plane in a surface of the second order which will be conjugate to a given system of parallel chords.

Let the inclinations of the chords to the axes be α, β, γ, then the equation of any one will be

$$x = mz + p, \quad y = nz + q \ldots . \ (1),$$

where

$$m = \frac{\cos \alpha}{\cos \gamma}, \quad n = \frac{\cos \beta}{\cos \gamma}.$$

For the points common to this line and the surface, we must combine this equation with equation (1), last proposition, and we shall have a result of the form

$$Rz^2 + Sz + T = 0 \ldots . \ (2),$$

which equation will furnish the two values of z corresponding to the two extremities of the diameter, and therefore half the sum of these values will be the z of the middle, that is, this z is

$$z = -\frac{S}{2R} \therefore 2Rz + S = 0 \ldots . \ (3),$$

which is obviously the differential coefficient derived from (2), or, which is the same thing, the total differential coefficient derived from (1) last proposition, in which x and y are functions of z given by the equations (1). This differential coefficient is, therefore,

$$\left\{\frac{du}{dz}\right\} = \frac{du}{dx}\frac{dx}{dz} + \frac{du}{dy}\frac{dy}{dz} + \frac{du}{dz} = 0$$

$$= m\frac{du}{dx} + n\frac{du}{dy} + \frac{du}{dz} = 0 \ . \ . \ . \ (4),$$

where p and q, the only quantities which vary with the chord, are eliminated; hence, this last equation represents the locus of the middle points of the chords or the diametral surface, and it is obviously a plane.

By actually effecting the differentiations indicated in equation (4) upon the equation (1), last proposition, we have, for the equation of the required diametral plane,

$$m(Ax + B'z + B''y + C) + n(A'y + Bz + B''x + C')$$
$$+ A''z + By + B'x + C'' = 0,$$

or

$$(Am + B' + B''n)x + (A'n + B + B''m)y +$$
$$(A'' + Bn + B'm)z + Cm + C'n + C'' = 0.$$

PROPOSITION IV.

(222.) A straight line moves so that three given points in it constantly rest on the same three rectangular planes; required the surface which is the locus of any other point in it.

Let the proposed planes be taken for those of the coordinates, and let the coordinates of the generating point be x, y, z, and the invariable distances of this point from the three points resting on the planes of yz, xz, and xy, X, Y, Z. The coordinates of these three points will be

In the plane of yz, 0, y', z'

xz, x'', 0, z''

xy, x''', y''', 0.

Then, since the parts of any straight line are proportional to their projections on any plane, each part having the same inclination to it, it follows that if we project successively each of the parts X, Y, Z, on the three coordinate planes, we shall have the relations

$$\left.\begin{aligned} \frac{x}{X} = \frac{x - x''}{Y} &= \frac{x - x'''}{Z} \\ \frac{y - y'}{X} = \frac{y}{Y} &= \frac{y - y'''}{Z} \\ \frac{z - z'}{X} = \frac{z - z''}{Y} &= \frac{z}{Z} \end{aligned}\right\} \quad \ldots\ (1).$$

But the part X of the moveable straight line comprised between the

generating point (x, y, z) and the point $(0, y', z')$ resting on the plane of y, z, has for its length the expression

$$X^2 = x^2 + (y - y')^2 + (z - z')^2,$$

or

$$1 = \frac{x^2}{X^2} + \frac{(y - y')^2}{X^2} + \frac{(z - z')^2}{X^2} \quad \ldots \quad (2),$$

but from the equations (1)

$$\frac{y - y'}{X} = \frac{y}{Y}, \quad \frac{z - z'}{X} = \frac{z}{Z};$$

hence, by substitution, (2) becomes

$$\frac{x^2}{X^2} + \frac{y^2}{Y^2} + \frac{z^2}{Z^2} = 1,$$

consequently, the surface generated is always of the second order. The surface would still be of the second order if the three *directing planes* were oblique instead of rectangular, as is shown by *M. Dupin*, in his *Developpements*, p. 342, whence the above solution is taken.

PROPOSITION V.

(223.) To determine the *line of greatest inclination* through any point on a curve surface.

The property which distinguishes the line of greatest inclination through any point is this, viz. that at every point of it the linear tangent makes with the horizon a greater angle than any other tangent to the surface drawn through the same point of the curve. Now, as all the linear tangents through any point are in the tangent plane to the surface at that point, that one which is perpendicular to the trace of the tangent plane will necessarily be the shortest, and therefore approach nearest to the perpendicular, that is, it will form a greater angle with the horizon than any of the others. We have, therefore, to determine the curve to which the linear tangent at every point is always perpendicular to the horizontal trace of the tangent plane to the surface through the same

point; or, which is the same thing, the projection of the linear tangent on the plane of xy must be perpendicular to the trace of the tangent plane.

Now the equation of the projection of the linear tangent at any point is

$$y - y' = \frac{dy'}{dx'}(x - x'),$$

and, by putting $z = 0$ in the equation of the tangent plane, we have, for the trace in the plane of xy, the equation

$$-z' = p'(x - x') + q'(y - y'),$$

and, since these two lines are to be *always* perpendicular to each other, we must have throughout the curve the general condition

$$\frac{dy}{dx} = \frac{q'}{p'} \therefore p' \frac{dy}{dx} - q' = 0,$$

p' and q' being derived from the equation of the surface; so that the values of these being obtained in terms of x and y, and substituted in the equation just deduced, the result will be the general differential equation belonging to the projection of every curve of greatest inclination that can be drawn on the proposed surface. To determine that passing through a particular point, or subject to a particular condition, we must, by help of the integral calculus, determine the general primitive equation from which the above is deducible; this primitive will involve an arbitrary constant, which may be fixed by the proposed condition, and thus the particular line be represented.

As an example, let us take the right conoid, of which the equation by (182) is

$$z = \phi\left(\frac{y}{x}\right).$$

Then for the ratio of the two partial differential coefficients p', q', which ratio is independent of the form ϕ (art. 58), we have

$$-\frac{\frac{1}{x}}{\frac{y}{x^2}} = \frac{dy}{dx}, \text{ by the condition above;}$$

hence

$$y\mathrm{d}y + x\mathrm{d}x = 0,$$

which equation is obviously the differential of

$$y^2 + x^2 = c,$$

where c is any constant quantity. The lines of greatest inclination are therefore such that their horizontal projections are concentric circles; the common centre being the point where the vertical axis, that is, the axis of z, pierces the plane of xy.

Again, let the surface be of the second order, and represented by

$$\frac{x^2}{\mathrm{A}} + \frac{y^2}{\mathrm{B}} + \frac{z^2}{\mathrm{C}} = 1;$$

then

$$\frac{\frac{y}{\mathrm{B}}}{\frac{x}{\mathrm{A}}} = \frac{\mathrm{d}y}{\mathrm{d}x} \therefore \mathrm{B}\frac{\mathrm{d}y}{y} - \mathrm{A}\frac{\mathrm{d}x}{x} = 0,$$

which equation is evidently the differential of

$$\mathrm{B}\log y - \mathrm{A}\log x = c;$$

and this leads to

$$y^{\mathrm{B}} \div x^{\mathrm{A}} = c, \text{ or } y^{\mathrm{B}} = c'x^{\mathrm{A}} \ldots\ldots (1),$$

where c' is the number whose logarithm is c, or $c' = \log^{-1} c$, and c is an arbitrary constant; hence the horizontal projections of the lines of greatest inclination will be parabolic curves when A and B have the same sign, and hyperbolic curves when the signs of A and B are different. (See ex. 6 and 7, page 148.)

Suppose the point of departure is in the plane of zx; then, since at that point $y = 0$, and $x = x'$, it follows that the equation of the line of greatest inclination through the point must be satisfied for these values; so that we must have

$$(2) \ldots\ldots 0 = c'x^{\mathrm{A}} \quad \therefore c' = 0.$$

Such, then, is the value of the arbitrary constant in the general equation (1) of these lines, for that particular line which we have selected from among them, so that this line coincides with the section made by the plane of xz; its projection being a straight line, which is a variety of the parabola.

PROPOSITION VI.

(224.) The six edges of any irregular tetraedron or triangular pyramid are opposed two by two, and the nearest distance of two opposite edges is called *breadth*; so that the tetraedron has *three* breadths and *four* heights. It is required to demonstrate that in every tetraedron the sum of the reciprocals of the squares of the *breadths* is equal to the sum of the reciprocals of the squares of the *heights*.

Let the vertex of the tetraedron be taken for the origin of the rectangular coordinates, and let also one of the faces coincide with the plane of xz, then the coordinates of the three corners of the base will be

$$0, 0, z', \mid x'', 0, z'', \mid x''', y''', z'''.$$

and the equations of the three edges terminating in the vertex will be

$$\begin{array}{l|l|l} x=0 & x=\frac{x''}{z''}z & x=\frac{x'''}{z'''}z \\ y=0 & y=0 & y=\frac{y'''}{z'''}z. \end{array}$$

Now the perpendicular distance between each of these edges and the opposite edge of the base will evidently be equal to the perpendicular demitted from the origin on a plane drawn through the latter edge and parallel to the former. Hence, denoting the three planes through the edges of the base by

$$Ax + By + Cz = 1 \mid Ex + Fy + Gz = 1 \mid Lx + Ky + Lz = 1,$$

they must be drawn so as to fulfil the conditions, (*See Anal. Geom.* p. 232,)

$$\begin{array}{rrr|rrr|rrr} & & Cz' = 1 & & & Gz' = 1 & Ix'' & +Lz'' = 1 \\ Ax'' & +Cz'' = 1 & & Ex''' + Fy''' + Gz''' = 1 & & & Ix''' + Ky''' + Lz''' = 1 & \\ Ax'''' + By'' + Cz''' = 0 & & & Ex'' & +Gz'' = 0 & & Lz' = 0 & \end{array}$$

These conditions fix the following values for A, B, C, &c. viz.

$$C = \frac{1}{z'}, \quad A = \frac{1}{x''} - \frac{z''}{x''z'}, \quad B = \frac{x''z''}{x''y'''z'} - \frac{x'''}{x''y'''} - \frac{z'''}{y'''z'};$$

$$G = \frac{1}{z'}, \quad E = -\frac{z''}{x''z'}, \quad F = \frac{1}{y'''} + \frac{x'''z''}{x''y'''z'} - \frac{z'''}{y'''z'};$$

$$L = 0, \quad I = \frac{1}{x'''}, \quad K = \frac{1}{y'''} - \frac{x'''}{x''y'''}.$$

Hence, calling the breadths B, B', B'', we have (*Anal. Geom.* p. 236,)

$$\frac{1}{B^2} = A^2 + B^2 + C^2 = \frac{(y'''z' - y'''z'')^2 + (x'''z'' - x'''z' - x''z''')^2 + (x''y''')^2}{(x''y'''z')^2}$$

$$\frac{1}{B'^2} = E^2 + F^2 + G^2 = \frac{(y'''z'')^2 + (x''z' + x'''z'' - x''z''')^2 + (x''y''')^2}{(x''y'''z')^2}$$

$$\frac{1}{B''^2} = I^2 + K^2 + L^2 = \frac{(y'''z')^2 + (x''z' - x'''z')^2}{(x''y'''z')^2}.$$

$$\text{Hence } \frac{1}{B^2} + \frac{1}{B'^2} + \frac{1}{B''^2} =$$

$$\left.\begin{array}{l} (z' - z'')^2 y''^2 + \{(z'' - z')x''' - x''z'''\}^2 + \\ \{(z' - z''')x'' + x'''z''\}^2 + (y'''z')^2 + (x'' - x''')^2 z'^2 \end{array}\right\} \div (x''y'''z')^2 \ldots\ldots (1).$$

Again, the expressions for the heights or perpendiculars demitted from each of the points

$$(0, 0, 0); \ (0, 0, z'); \ (x'', 0, z''); \ (x''', y''', z'''),$$

upon the plane which passes through the other three are, severally, (*Anal. Geom.* p. 236,)

$$H^2=\frac{(x''y'''z')^2}{(z''-z')^2y'''^2+\{(z'''-z')x''+(z'-z'')x'''\}^2+(y'''x'')^2}$$

$$H'^2=\frac{(x''y'''z')^2}{(z''y''')^2+(x''z'''-x'''z'')^2+(x''y''')^2},$$

$$H''^2=\frac{(x''y'''z')^2}{(y'''z')^2+(x'''z')^2},\quad H'''^2=\frac{(x''y'''z')^2}{(x''z')^2},$$

$$\therefore \frac{1}{H^2}+\frac{1}{H'^2}+\frac{1}{H''^2}+\frac{1}{H'''^2}=$$

$$\left.\begin{array}{l}(z''-z')^2y'''^2+\{(z'''-z')x''+(z'-z'')x'''\}^2+\\ 2(x''y''')^2+(y'''z'')^2+(x''z'''-x'''z'')^2+(y'''^2+x'''^2+x''^2)z'^2\end{array}\right\}\div(x''y'''z')^2\ldots(2).$$

which expression is the same as that before deduced, and thus the theorem is established by a process purely analytical. This remarkable property was discovered by *M. Brianchon*, and formed the subject of the prize question in the Ladies' Diary for 1830. Elegant solutions upon different principles by Messrs. Woolhouse and Godward may be seen in Professor Leybourn's Mathematical Repository, No. 25.

END OF THE DIFFERENTIAL CALCULUS.

NOTES.

NOTE (A), *page* 51.

On the Summation of slowly Converging Infinite Series.

LET

$$a - bx + cx^2 - dx^3 + \&c. = S,$$

$$\therefore\ -bx + cx^2 - dx^3 + \&c. = S - a$$

$$\therefore\ -b + cx - dx^2 + ex^3 - \&c. = \frac{S - a}{x}$$

Consequently, by adding these two last equations together, and representing the numerical differences, $b - c$, $c - d$, $d - e$, &c., by Δ, Δ', Δ'', &c., there results the equation

$$-b - \Delta x + \Delta' x^2 - \Delta'' x^3 + \&c. = \frac{x+1}{x}(S - a),$$

$$\therefore\ -bx - \Delta x^2 + \Delta' x^3 - \Delta'' x^4 + \&c. = (x+1)(S - a) = S'$$

$$\therefore\ S = \frac{S'}{x+1} + a;$$

that is,

$$S = a - \frac{bx}{x+1} - \frac{\Delta x^2}{x+1} + \frac{\Delta' x^3}{x+1} - \frac{\Delta'' x^4}{x+1} + \&c.$$

$$= a - \frac{bx}{x+1} + \frac{x}{x+1}\{0 - \Delta x + \Delta' x^2 - \Delta'' x^3 + \&c.\}.$$

Hence, treating the series within the brackets as we have treated the original, to which it is similar in form, we have

$$S = a - \frac{bx}{x+1} - \frac{\Delta x^2}{(x+1)^2} - \frac{\Delta^2 x^3}{(x+1)^2} + \frac{\Delta^{2'} x^4}{(x+1)^2} - \frac{\Delta^{2''} x^5}{(x+1)^2} + \&c.$$

Similarly,

$$S = a - \frac{bx}{x+1} - \frac{\Delta x^2}{(x+1)^2} - \frac{\Delta^2 x^3}{(x+1)^3} - \frac{\Delta^3 x^4}{(x+1)^3} + \frac{\Delta^{3'} x^5}{(x+1)^3}$$

$$\vdots\qquad\qquad - \&c.$$

$$S = a - \frac{bx}{x+1} - \frac{\Delta x^2}{(x+1)^2} - \frac{\Delta^2 x^3}{(x+1)^3} \cdots - \frac{\Delta^n x^{n+1}}{(x+1)^{n+1}} - \frac{\Delta^{n+1} x^{n+2}}{(x+1)^{n+1}}$$

$$+ \&c. \ldots . \text{(A)}.$$

These several expressions for S may be regarded as so many *differential theorems;* but the last is the form which corresponds most nearly to that under which the differential theorem is usually given. It is, however, more efficient, as it shows that if we stop at the term

$$\frac{\Delta^n x^{n+1}}{(x+1)^{n+1}},$$

we get one limit to the sum S; and if we stop at the immediately following term

$$\frac{\Delta^{n+1} x^{n+2}}{(x+1)^{n+1}},$$

we get another limit in the contrary sense. These limits, as we shall presently see, may be easily narrowed.

By considering the series to terminate at the former of the above terms, it will be the same as that investigated by the late Mr. Baron Maseres, in his *Scriptores Logarithmici*, vol. iii., p. 219; and which the Baron considered to be different from that deduced by Simpson, from a formula still more general, in his *Mathematical Dissertations*, p. 62. The two forms are however mutually interchangeable, regard being paid to the signs of the differences.

Suppose, in the proposed series S, that $a = 0$; then, dividing both sides of (A) by $-x$, we shall have

$$S = b - cx + dx^2 - ex^3 + \&c.$$

$$= \frac{b}{x+1} + \frac{\Delta x}{(x+1)^2} + \frac{\Delta^2 x^2}{(x+1)^3} \cdots + \frac{\Delta^n x^n}{(x+1)^{n+1}} + \frac{\Delta^{n+1} x^{n+1}}{(x+1)^{n+1}}$$

$$- \&c. \ldots . \text{(B)},$$

which, when $x = 1$, becomes

$$S = b - c + d - e + \&c.$$

$$= \frac{b}{2} + \frac{\Delta}{4} + \frac{\Delta^2}{8} \ldots\ldots + \frac{\Delta^n}{2^{n+1}} + \frac{\Delta^{n+1}}{2^{n+1}} - \&c. \ldots\ldots (C).$$

This series, supposing it to terminate at $\frac{\Delta^n}{2^{n+1}}$, furnishes the method proposed by Dr. Hutton in his valuable *Mathematical Tracts*, vol. i., p. 176. In Dr. Hutton's process the value of S is approached by determining, in succession, the values of

$$\frac{b}{2}, \quad \frac{b}{2} + \frac{\Delta}{4}, \quad \frac{b}{2} + \frac{\Delta}{4} + \frac{\Delta^2}{8}, \quad \&c.$$

which is done by finding first the successive sums of a few of the leading terms of the proposed series, taking the arithmetical mean between each pair of consecutive results, then the mean between each pair of these means, and so on. But simple as this operation undoubtedly is, yet, when the numbers consist of several places of figures, it cannot be safely performed mentally; so that Dr. Hutton's numerical examples, which exhibit the *results* only of these operations, do not fairly present the whole amount of numerical labour. Still his method is, upon the whole, more easy and convenient than any other which we have met with. Like all the ordinary processes, however, that method still leaves us in doubt as to the accuracy of a final decimal or two in the results determined by it, as indeed every process of approximation must necessarily do which does not furnish limits both above and below the value sought.

In the series above, marked (B), it is plain that if we stop at the term involving Δ^n, we shall obtain an *inferior* limit to the sum S; and, if we take in the term immediately following, we shall get a *superior* limit.

Now, to narrow these limits, we must observe that since

$$\Delta^{n+1} = \Delta^n - \Delta^{n'},$$

a superior limit will be obtained by adding to the inferior limit the quantity

$$\frac{\Delta^n x^{n+1}}{(x+1)^{n+1}}.$$

Moreover, the inferior limit cannot differ from the entire sum by so much as

$$\frac{\Delta^{n+1} x^{n+1}}{(x+1)^{n+1}},$$

whereas the superior limit differs from the truth by more than

$$\frac{\Delta^{n'} x^{n+1}}{(x+1)^{n+1}},$$

inasmuch as the succeeding term in the series (B) is also negative. The inferior limit will therefore be nearer to the truth than the superior, since $\Delta^{n'} > \Delta^{n+1}$; and consequently, half the sum of the two limits will be a superior limit still nearer. We may conclude, therefore, that if we multiply the final term in the inferior limit by $\frac{x}{2}$, and add the product to that limit, we shall thus obtain a near superior limit.

It is scarcely necessary to observe here, that in what has hitherto been said, the coefficients in the proposed series, S, are supposed continually to diminish, as also the several series of differences, which supposition is conformable to what usually occurs in practice, S being a *convergent* series.

As an example, let it be required to sum the converging series

$$1 - \frac{1}{3} + \frac{1}{5} - \frac{1}{7} + \frac{1}{9} - \frac{1}{11} + \&c.$$

It will be advisable to actually sum up first a few of the leading terms, and then to apply the formula (B), or rather (C), to the remainder of the series. The work may be arranged as follows:

+ 1					
− ·3					
+ ·2					
+ ·142857					
+ ·111111					
− ·090909	·744012 = sum of first six terms.				
+ ·076923	10256	2412	759	284	
− ·066667	7844	1653	475		
+ ·058823	6191	1178			
− ·052632	5013				
+ ·047619					

The numbers ·076923, ·010256, ·002412, ·000759, and ·000284, are the respective values of b, Δ^2, Δ^3, and Δ^4, in (C); and to deduce from these the value of S, it will be necessary merely to add half the last number, 284, to the preceding; half the sum to the next; and so on, to the last, adding half the final sum to the number ·744012, previously found. The remainder of the operation is therefore as follows:

·076923	10256	2412	759	284
·082767	11687	2862	901	142
·041384	5844	1431	450	
·744012				

·785396 = Inferior limit.

$$4 = \frac{142}{2^5}.$$

·785400 = Superior limit.

$\therefore$ ·785398 = $\frac{\pi}{4}$ very nearly.

This value of $\frac{\pi}{4}$ we may be assured, from the small interval between the limits, cannot possibly differ from the truth by more than a unit in the final decimal. It is in fact true even in the last figure.

We have separated the two parts of the process in this example for the purpose of clearer illustration; but they may be combined, as in the following example:

To find the value of the converging series

$$1 - \frac{1}{2^2} + \frac{3^2}{2^2 . 4^2} - \frac{3^2 . 5^2}{2^2 . 4^2 . 6^2} + \frac{3^2 . 5^2 . 7^2}{2^2 . 4^2 . 6^2 . 8^2} - \&c.,$$

which occurs in the expression for the time of vibration of a pendulum in a circular arc:



834625 = Inferior limit.

$$3 = \frac{106}{2^5}.$$

·834628 = Superior limit.

·834626 = S very nearly.

A value which cannot differ from the truth by more than a unit in the last decimal. It is in fact true in all its places.*

---

* For further particulars respecting this method of summing infinite series, see a paper by the author of the present work in the Philosophical Magazines for May and July, 1835.

NOTE (B), *page* 117.

Demonstration of the Theorems of Laplace and Lagrange.

Let it be required to develop the function

$$u = \Psi(z), \text{ where } z = F(y + xf(z)).$$

By differentiating the second of these equations, first relatively to x, and then relatively to y, we have

$$\frac{dz}{dx} = F'(y + xf(z))\left(f(z) + xf'(z)\frac{dz}{dx}\right)^*,$$

$$\frac{dz}{dy} = F'(y + xf(z))\left(1 + xf'(z)\frac{dz}{dy}\right).$$

Multiplying the first by $\frac{dz}{dy}$, and the second by $\frac{dz}{dx}$, and subtracting, there results

$$\frac{dz}{dx} - f(z)\frac{dz}{dy} = 0 \therefore \frac{dz}{dx} = f(z)\frac{dz}{dy} \quad \ldots\ (1);$$

but, since u or $\Psi(z)$ depends only on z, we shall have

$$\frac{du}{dx} = \Psi'(z)\frac{dz}{dx}, \quad \frac{du}{dy} = \Psi'(z)\frac{dz}{dy},$$

therefore, eliminating $\Psi(z)$, we get

$$\frac{du}{dx}\frac{dz}{dy} - \frac{du}{dy}\frac{dz}{dx} = 0,$$

---

* At page 88 we put $f'(z)$ to represent the differential coefficient of $f(z)$ relatively to x; here the same symbol denotes the coefficient relatively to z.

or putting for $\frac{dz}{dx}$ its value (1), and making for abridgment $f(z) = Z$, this last expression becomes divisible by $\frac{dz}{dy}$, and reduces to

$$\frac{du}{dx} = Z \frac{du}{dy} \ldots\ldots (A),$$

so that we may always substitute for $\frac{du}{dx}$ the quantity $Z \frac{du}{dy}$.

If we differentiate the preceding equation relatively to x, we shall obtain

$$\frac{d^2u}{dx^2} = \frac{dZ \frac{du}{dy}}{dx} \ldots\ldots (2),$$

but the expression $Z \frac{du}{dy}$ being no other than $f(z)\Psi'(z)\frac{dz}{dy}$, that is to say a function of z multiplied by $\frac{dz}{dy}$, we may consider it as the differential coefficient of some new function of z, which we may represent by u_1, and we shall then have

$$\frac{du_1}{dy} = Z \frac{du}{dy}, \quad \text{and} \quad \frac{dZ \frac{du}{dy}}{dx} = \frac{d^2u_1}{dx\,dy};$$

therefore (2), inverting the order of the differentiations in this last expression,

$$\frac{d^2u}{dx^2} = \frac{d^2u_1}{dy\,dx} = \frac{d \frac{du_1}{dx}}{dy} \ldots\ldots (3),$$

now, it must be observed that the relation (A) exists, whatever be the function u; it therefore exists for the function u_1; hence

$$\frac{du_1}{dx} = Z \frac{du_1}{dy}.$$

Substituting then in (3) for $\frac{du_1}{dx}$ its value here exhibited, and afterwards

for $\frac{du_1}{dy}$ its equal $Z\frac{du}{dy}$, there results

$$\frac{d^2u}{dx^2} = \frac{dZ\frac{du_1}{dy}}{dy} = \frac{dZ^2\frac{du}{dy}}{dy} \quad \ldots\ (B).$$

Differentiating this last equation relatively to x, we shall obtain

$$\frac{d^3u}{dx^3} = \frac{d^2Z^2\frac{du}{dy}}{dx\,dy},$$

and considering, as before, the function $Z^2\frac{du}{dy}$ to be the differential coefficient of some new function of z, viz. u_2, we shall have, by inverting the order of the differentiations,

$$\frac{du_2}{dy} = Z^2\frac{du}{dy}, \text{ and } \frac{d^3u}{dx^3} = \frac{d^2\frac{du_2}{dx}}{dy^2} \quad \ldots\ (4),$$

but the equation (A) subsisting for every function u, must have place for the function u_2, hence

$$\frac{du_2}{dx} = Z\frac{du_2}{dy},$$

therefore (4),

$$\frac{du_2}{dx} = Z^3\frac{du}{dy}, \quad \frac{d^3u}{dx^3} = \frac{d^2Z^3\frac{du}{dy}}{dy^2} \quad \ldots\ (C).$$

The analogy among the expressions (A), (B), (C), is obvious, and we shall now show that this analogy continues uninterrupted; that is, generally, if

$$\frac{d^{n-1}u}{dx^{n-1}} = \frac{d^{n-2}Z^{n-1}\frac{du}{dy}}{dy^{n-2}} \quad \ldots\ (M),$$

then

$$\frac{d^nu}{dx^n} = \frac{d^{n-1}Z^n\frac{du}{dy}}{dy^{n-1}} \quad \ldots\ (N).$$

For considering, as before, the function $Z^{n-1}\frac{du}{dy}$ to be the differential coefficient of some new function of z, viz. u_{n-1}, so that

$$Z^{n-1}\frac{du}{dy}=\frac{du_{n-1}}{dy} \quad \ldots\ldots (5),$$

we have, by differentiating (M),

$$\frac{d^n u}{dx^n}=\frac{d^{n-1}\frac{du_{n-1}}{dx}}{dy^{n-1}},$$

but the equation (A) subsisting for every function of z subsists for u_{n-1}, therefore

$$\frac{du_{n-1}}{dx}=Z\frac{du_{n-1}}{dy};$$

hence (5)

$$\frac{d^n u}{dx^n}=\frac{d^{n-1}Z^n\frac{du}{dy}}{dy^{n-1}}=\frac{d^{n-1}f(z)\frac{du}{dy}}{dy^{n-1}}.$$

Let now $x=0$ in the original function, and in each of the coefficients $\frac{du}{dx}$, $\frac{d^2u}{dx^2}$ $\ldots\ldots$ $\frac{d^n u}{dx^n}$; then we have

$$[u]=\Psi(F(y))=\phi(y), \quad [z]=F(y) \quad \therefore f[z]=f(F(y))=\psi(y),$$

and

$$[\frac{d^n u}{dx^n}]=\frac{d^{n-1}(\psi(y))^n\frac{d\phi(y)}{dy}}{dy^{n-1}}.$$

Consequently, by Maclaurin's theorem,

$$u=\phi(y)+\psi(y)\frac{d\phi(y)}{dy}\frac{x}{1}+\frac{d(\psi(y))^2\frac{d\phi(y)}{dy}}{dy}\frac{x^2}{1.2}+$$

$$\frac{d^2(\psi(y))^3\frac{d\phi(y)}{dy}}{dy^2}\frac{x^3}{1.2.3}+\text{&c.}$$

which is the theorem of *Laplace*.

The preceding investigation is taken with some slight variation from the large work of Lacroix, vol. I. p. 279.

In the particular case where

$$u = \Psi(z) \text{ and } z = y + xf(z),$$

we have

$$[u] = \Psi(y), \quad [z] = F(y) = y, \quad f[z] = \psi(y) = f(y);$$

hence the development is then

$$u = \Psi(y) + f(y)\frac{d\Psi(y)}{dy}\frac{x}{1} + \frac{d(f(y))^2\frac{d\Psi(y)}{dy}}{dy}\frac{x^2}{1.2} + \&c.$$

which corresponds with the theorem of *Lagrange*, given at page 117, Ψ being here put for F.

NOTE (C), *page* 160.

Suppose the function $f(x + h)$ fails to be developable according to Taylor's series for the value $x = a$, and let the true development be represented by

$$f(a + h) = f(a) + Ah^{\alpha} + Bh^{\alpha+\beta} + Ch^{\alpha+\beta+\gamma} + \&c. \ldots . (1),$$

the terms being arranged according to the powers of h. It is required actually to find these terms.

Let the difference $f(a + h) - f(a)$ be divided by such a power of h, that the quotient will become neither 0 nor ∞; when $h = 0$, such a power of h can be no other than h^{α}; or that which ought to appear in the first term of this difference, for if the developed difference were divided by a lower power of h than this, the quotient would evidently be 0, when $h = 0$ and if it were divided by a higher power, the quotient would be infinite when $h = 0$; hence the proper divisor h^{α} being found, if we put $h = 0$ in the quotient, the result will be simply A; having thus found the true first term of the difference, let it be transposed to the other side, and we shall then have the difference

$$\frac{f(a+h)-f(a)}{h^{\alpha}}-A=Bh^{\beta}+Ch^{\beta+\gamma}+\&c.$$

Now the first side of this equation being known, we have, as before, to find what power of h it must be divided by, so that the quotient may be neither 0 nor infinite, when $h=0$; this power will be h^{β}, and, putting $h=0$ in the quotient, the result is B; and in this manner it is plain that all the terms of the series are to be determined.

Let now $y=f(x)$ be the equation of a plane curve, and $Y=F(x)$ the equation of another having a common point with the former, at which $x=a$; then

$$F(a)=f(a);$$

beyond this point the ordinates of the two curves, for any abscissa $(a+h)$, will be $f(a+h)$ and $F(a+h)$, and their difference D will be

$$D=f(a+h)-F(a+h).$$

Let us now develop the function $F(a+h)$, as we have done the function $f(a+h)$, and we shall have

$$F(a+h)=F(a)+A'h^{\alpha'}+B'h^{\alpha'+\beta'}+C'h^{\alpha'+\beta'+\gamma'}+\&c.\ldots.(2);$$

and it may be shown precisely, as at page 161, that the greater the number of identical leading terms in the two developments (1), (2), the nearer will the developments themselves approach to identity; so that no curve passing through the point common to the two former can approach so closely to either, in the vicinity of that point, unless in the development of the ordinate the same number of leading terms agree with those in (1).

Lagrange observes (*Théorie des Fonctions Analytiques*, p. 184,) that we may call contact of the first order, contact of the second order, &c. the approximation of two curves, for which the two first terms, the three first terms, &c. are the same in the developments of the functions which represent the ordinates. All other authors who advert to this subject make the same remark, but it is erroneous, as a simple instance will show.

Let the developed ordinates be

$$A+Bh^{\frac{3}{2}}+Ch^{\frac{9}{4}}+Dh^{4}+\&c.$$

$$A+Bh^{2}+Ch^{\frac{9}{4}}+Mh^{\frac{11}{4}}+\&c.$$

According to Lagrange the contact here would be of the second order, but, by Taylor's theorem, these developments would be

$$A + 0h + Bh^2 + 0h^3 + 0h^4 + \&c.$$

$$A + 0h + Bh^2 + 0h^3 + 0h^4 + \&c.$$

and therefore, by the general principles established in Chap. II. Sect. II. the contact is of the fourth order, at least. If in the true developments the term next to $0h^4$ were the same in each, and the exponent of h a fraction between 4 and 5, while the term following differed in the two series, the contact might be properly said to be of the fifth order; but the sign of the difference of the two developments, when h is negative, will obviously depend on the fractional exponents of h, in the terms immediately beyond those which agree in the two series.

NOTE (D), *page* 277.

As the process from which the expressions for α, β, γ, given in article 208, are deduced, is not immediately obvious, we shall here exhibit it at length for the first of these expressions.

From the formulas for p'' and q'', at article 202, we immediately get for $p'p'' + q'q''$ the value

$$\frac{(d^2y)\,(dy)\,(dx) - (d^2x)\,(dy)^2 + (d^2z)\,(dz)\,(dx) - (d^2x)\,(dz)^2}{(dx)^4}$$

$$= \frac{(dx)\,\{(d^2y)\,(dy) + (d^2z)\,(dz)\} - (d^2x)\,\{(dy)^2 + (dz)^2\}}{(dx)^4}$$

$$= \frac{(dx)\,\{(ds)\,(d^2s) - (d^2x)\,(dx)^2\}^{*} - (d^2x)\,\{(ds)^2 - (dx)^2\}}{(dx)^4}$$

---

* Because from

$$(ds)^2 = (dx)^2 + (dy)^2 + (dz)^2$$

we get

$$(ds)\,(d^2s) = (dx)\,(d^2x) + (dy)\,(d^2y) + (dz)\,(d^2z).$$

$$= \frac{(dx)(ds)(d^2s) - (d^2x)(ds)^2}{(dx)^4} = \frac{(ds)\{(dx)(d^2s) - (d^2x)(ds)\}}{(dx)^4}.$$

Putting now for brevity $\frac{A}{(dx)^6}$ for what the denominator in the expression for α, article (201), becomes, when the independent variable is arbitrary, we have

$$\alpha = x + \frac{\frac{(ds)^2}{(dx)^2}}{\frac{A}{(dx)^6}} \cdot \frac{(ds)\{(d^2x)(ds) - (d^2s)(dx)\}}{(dx)^4}$$

$$= x + \frac{(ds)^3}{A} \cdot (d^2x)(ds) - (d^2s)(dx)$$

$$= x + \frac{(ds)^6}{A} \cdot \frac{(d^2x)(ds) - (d^2s)(dx)}{(ds)^3}$$

$$= x + r^2 \frac{d^2x}{ds^2}, \quad \text{art. (66)},$$

and by a similar process the expressions for β and γ are obtained.

The expressions $\alpha - x$, $\beta - y$, $\gamma - z$ are obviously the projections of the radius of curvature r on the axes of x, y, z. But, if we represent the inclinations of the radius to the axes by λ, μ, ν, the expressions for the projections will be

$$r\cos\lambda, \quad r\cos\mu, \quad r\cos\nu,$$

so that (208) we have, for the angles of inclination, the values

$$\cos\lambda = r\frac{d^2x}{ds^2}, \quad \cos\mu = r\frac{d^2y}{ds^2}, \quad \cos\nu = r\frac{d^2z}{ds^2}.$$

By employing these expressions *M. Cauchy* has arrived by rather a novel process at the theorem of *Meusnier*, given at p. 233.

Let the equation of any curve surface be

$$u = F(x, y, z) = 0,$$

upon which is traced any curve MG′, determined by the equation

$$y = \phi(x),$$

joined to the preceding.



If through the tangent MT to this curve, and also through the normal MN of the surface, we draw a plane, we shall be furnished with a normal section MG, of which the radius of curvature r, at M, will be some portion of the normal MN. Also the radius of curvature r' of the assumed curve MG′, at the same point, will be some portion of the line MN′, perpendicular to the tangent MT.

Now, considering s to be the independent variable, we have, for the inclinations of r' to the axes the expressions above, viz.

$$r' \frac{d^2x}{ds^2}, \quad r' \frac{d^2y}{ds^2}, \quad r' \frac{d^2z}{ds^2},$$

and the inclinations of r or of MN to the axes are (129)

$$v \frac{du}{dx}, \quad v \frac{du}{dy}, \quad v \frac{du}{dz}.$$

Hence, calling the angle N′MN, between the two radii, ω, we have (*Anal. Geom.* p. 239)

$$\cos \omega = v r' \left(\frac{du}{dx} \frac{d^2x}{ds^2} + \frac{du}{dy} \frac{d^2y}{ds^2} + \frac{du}{dz} \frac{d^2z}{ds^2}\right).$$

But the equation of the surface, considered as one of the equations of the curve MG′, gives after two successive differentiations, still regarding s as the independent variable,

$$\frac{du}{dx} \frac{d^2x}{ds^2} + \frac{du}{dy} \frac{d^2y}{ds^2} + \frac{du}{dz} \frac{d^2z}{ds^2} =$$

$$\left\{ \begin{array}{l} - \dfrac{d^2u}{dx^2} \dfrac{dx^2}{ds^2} - \dfrac{d^2u}{dy^2} \dfrac{dy^2}{ds^2} - \dfrac{d^2u}{dz^2} \dfrac{dz^2}{ds^2} \\ - 2 \dfrac{d^2u}{dx\,dy} \dfrac{dx\,dy}{ds^2} - 2 \dfrac{d^2u}{dx\,dx} \dfrac{dx\,dz}{ds^2} - 2 \dfrac{d^2u}{dy\,dz} \dfrac{dy\,dz}{ds^2}. \end{array} \right.$$

Now whatever be the curve MG′, provided only its tangent MT $= x'$ remains unchanged, the second member of this last equation will remain

unchanged, because the values of $\frac{dx}{ds}$, $\frac{dy}{ds}$, $\frac{dz}{ds}$, which are the same as those of $\frac{dx}{dx'}$, $\frac{dy}{dx'}$, $\frac{dz}{dx'}$, remain unchanged. Therefore this second member being substituted in the expression for cos ω leads to a result of the form

$$r' = K \cos \omega,$$

K being a constant expression for all the curves on the proposed surface which touch MT at the point M. Put now, in this expression, $\omega = 0$, then r' becomes r, therefore

$$r = K, \text{ consequently } r' = r \cos \omega,$$

which result comprehends the theorem of Meusnier; since, if the curve MG′ is plane, its plane will coincide with N′MT, and the angle ω of the two radii will become the angle formed by the plane N′MT of the oblique section with the plane NMT of the normal section passing through the same tangent MT.—*Leroy, Analyse Appliquée à la Géométrie*, p. 268

We may take this opportunity of remarking that, in our investigation of this theorem, at p. 233, it might easily have been shown, without referring to article (86), that

$$\frac{dx'^2}{dx^2} = 1 + \tan^2 \theta,$$

because, by the right-angled triangle,

$$x'^2 = x^2 \sec^2 \theta = x^2 (1 + \tan^2 \theta)$$

$$\therefore \frac{dx'^2}{dx^2} = 1 + \tan^2 \theta = \frac{ds^2}{dx^2}.$$

NOTE (E), *page* 136.

The erroneous doctrine adverted to at page 136 is laid down also by *Lacroix*, in his quarto treatise on the Calculus, vol. I. p. 340, from whom, indeed, Mr. Jephson seems to have adopted it. The principle as stated by Lacroix is "que la série de Taylor devient illusoire pour toute valeur qui rend imaginaire l'un quelconque de ces terms; et que cela peut arriver sans que la fonction soit elle-même imaginaire."

THE END.

Printed by J. and C. ADLARD, Bartholomew Close.

515
Y73